Surface Chemistry

INTRODUCTION :

Surface chemistry is that branch of chemistry which deals with study of the phenomena occuring at the surface or interface, i.e. at the boundary separating two bulk phases. In this chapter our main emphasis will be on three important topics related to surface chemistry, viz., adsorption, colloids and emulsions.

Section (A) : Adsorption

• Adsorption : The phenomenon of attracting and retaining the molecules of a substance on the surface of a liquid or a solid resulting into a higher concentration of the molecules on the surface is called adsorption. As a result of adsorption, there is a decrease of surface energy. The process of removal of an adsorbed substance from the surface on which it is adsorbed is called desorption. It is the reverse of adsorption and can be brought about by heating or by reducing the pressure.

• Adsorbent and adsorbate : The substance on the surface of which adsorption occurs is known as adsorbent. The substances that get adsorbed on the solid surface due to intermolecular attractions are called adsorbate. Charcoal, silica, gel, alumina gel are good adsorbents because they have highly porous structures and have large surface area. Colloids on account of their extremely small dimensions possess enoromous surface area per unit mass and are, therefore, also good adsorbents.

Examples of adsorption :

• Adsorption of a gas by charcoal : Finely divided activated charcoal has a tendency to adsorb a number of gases like ammonia, sulphur dioxide, chlorine, phosgene, etc. In this case, charcoal acts as an adsorbent while gas molecules act as adsorbate.

• Adsorption of a dye by charcoal : Animal charcoal is used for decolourising a number of organic substances in the form of their solutions. The discharge of the colour is due to the fact that the coloured component (generally an organic dye) gets adsorbed on the surface of the adsorbent (animal charcoal).

• **Sorption :** When both adsorption and absorption take place simultaneously.

Eg: Dyes get adsorbed as well as absorbed in the cotton fibre i.e. sorption takes place.

Difference between adsorption and absorption :

The important points of distinction between adsorption and absorption

Absorption	Adsorption
It is the phenomenon in which the particles of gas or liquid get uniformly distributed throughout the body of	It is the phenomenon of higher concentration of gas or liquid on the surface than in the bulk of the solid.
The concentration is the same throughout the material.	The concentration on the surface of the adsorbent is different (has higher concentration) from that in the bulk.
lt is a bulk phenomenon.	it is a surface phenomenon.
Absorption occurs at uniform rate.	Adsorption is rapid in the beginning and its rate slowly decreases
It is a slow process	It is a fast process

Thermodynamics of adsorption

1. Adsorption involves attracting molecules of adsorbate on surface of the adsorbent. Due to this energy is released and thus heat of adsorption is negative i.e. adsorption is always exothemic

2. The molecules of adsorbate are held on surface of the solid adsorbent due to this entropy decreases i.e.

∆S is negative.

3. $\Delta G = \Delta H - T\Delta S$, Therefore adsorption will occur only when ΔG is negative and this is possible only if $|\Delta H| > |T\Delta S|$

4. As adsorption proceeds ΔH becomes less and less negative hence ΔH becomes equal to TΔS and ΔG becomes zero This is the state at which equilibrium is attained.

Enthalpy of adsorption $\Delta H_{adsorption}$: It is the amount of the heat released when 1 mole of an adsorbate gets adsorbed on a particular adsorbent at adsorption equilibrium. It depends upon the nature of both the adsorbate as well as adsorbent.

Types of adsorption : The adsorption is classified into two types :

Physical adsorption (i.e. physisorption) : When the particles of the adsorbate are held to the surface (i) of the adsorbent by the physical forces such as van der Waal's forces, the adsorption is called physical adsorption or vanderwaals adsorption.

(ii) Chemical adsorption (i.e. chemisorption) :

When the molecules of the adsorbate are held to the surface of the adsorbent by the chemical forces, the adsorption is called chemical adsorption.

Differences between Physical and Chemical Adsorption		
Property	physical Adsorption	Chemical Adsorption
Nature of adsorption	Weak	Strong
Enthalpy of adsorption	Low	High
Reversibility of adsorption	Reversible and occur rapidly	Irreversible and occurs slowly
Temp. at which adsorption is more pronounced	Low temp.	High temp.
Effect of change in temp.	Decrease with rise in temp.	Increase with rise in temp.
	Not specific generally take place	Highly specific take place on
Specificity of adsorption	on all surface	specific surface
Nature of adsorbate layer	Multi-layered	Mono-layered
Effect of pressure on adsorption	Increase with pressure of adsorbate gas and attain limiting value	It is not appreciably affected by small changes in pressure. However, very high pressures are favourable
Energy of activation	Very low	significant relative high
Dependence on nature	depends on adsorbate (gas)	Depends on adsorbate as well
of adsorbate and	only easily liquefiable gases are	as adsorbent No correlation be
adsorbent	more readily adsorbed	can be achieved
Easy of desorption	easy since Vander Waal force involved	Not easy since chemical forces on involved
Graph	x m Temperature	x m Temperature

• **Competitive adsorption**: When an adsorbent is in contact with more than one species (adsorbate). There will be competition among them to get adsorbed on to the surface of the adsorbent. The one that is more strongly adsorbed gets deposited first in preference to the others. Further a strongly adsorbed substance may displace a weakly adsorbed substance.

Ex. NH_3 can displace O_2 or N_2 from the surface of charcoal.

Adsorption of gases on solids :

The extent of adsorption of a gas on a solid surface is affected by the following factors:

The nature of the gas (i.e. nature of the adsorbate). The easily liquefiable gases such as HCl, NH₃, Cl_2 etc. are adsorbed more than the permanent gases such as H₂, N₂ and O₂.

extent of adsorption critical temperature of gasease of liquification

Gas	H_2	CH₄	CO_2	SO2
Tc	33	190 K	304 K	330K

• **Nature of adsorbent.** The extent of adsorption of a gas depends upon the nature of adsorbent. Activated charcoal (i.e. activated carbon), metal oxides (silica gel and aluminium oxide) and clay can adsorb gases which are easily liquified. Gases such as H₂, N₂ and O₂ are generally adsorbed on finely divided transition metals Ni and Co.

extent of adsorption \propto surface area of solid.

• Activation of adsorbent :

(a) Metallic adsorbents are activated by mechanical rubbing or by subjecting it to some chemical reactions.

(b) To increase the adsorbing power of adsorbents, they are sub-divided into smaller pieces. As a results, the surface area is increased and therefore, the adsorbing power increases.

• Effect of temperature :Mostly the process of adsorption is exothermic and the reverse process or desorption is endothermic. If the above equilibrium is subjected to increase in temperature, then according to Le-Chaterlier's principle, with increase in temperature, the desorption will be favoured. Physical adsorption decreases continuously with increase in temperature whereas chemisorption increases initially, shows a maximum in the curve and then it decreases continuously.



The initial increase in chemisorption with increase in temperature is because of activation energy required.

This is why the chemical adsorption is also known as "Activated adsorption".

A graph between degree of adsorption (x/m) and temperature 't' at a constant pressure of adsorbate gas is known as **adsorption isobar**.

• Effect of pressure. The extent of adsorption of a gas per unit mass of adsorbent depends upon the pressure of the gas. The variation of extent of adsorption expressed as x/m (where x is the mole of adsorbate and m is the mass of the adsorbent) and the pressure is given as below. A graph between the amount of adsorption and gas pressure keeping the temperature constant is called an **adsorption isotherm.**



 $\frac{x}{m} \propto P$ Where x = mass of adsorbate M = mass of adsorbent

It is clear from the figure-1 that extent of adsorption (x/m) increases with pressure and becomes maximum corresponding to pressure P_s called saturation pressure.

Freundlich Adsorption isotherm

The variation of extent of adsorption (x/m) with pressure (P) was given mathematically by Freundlich. Where n can take any whole number value which depends upon the nature of adsorbate and adsorbent. The above relationship is also called Freundlich's adsorption isotherm.

$$\left(\frac{x}{m}\right) = kp^{1/n}$$

Pressure	Graph	Relation
At low pressure	straight line	$\frac{x}{m} = kP$
At intermediate pressure	dependent on power of pressure	$(x/m) = kp^{1/n}$ $(0 < \frac{1}{n} < 1)$
At high pressure	: Independent of pressure	$\frac{x}{m} = k$

where, x = Amount of gas adsorbed, m = Mass of adsorbent, K and n are adsorption constant, p is adsorption equilibrium pressure.

The constant k and n can be determined as explained below : Taking logarithms on both sides

Eq. $(x/m) = kp^{1/n}$

 $\log (x/m) = \log k + (1/n) \log p.$



One of the drawbacks of Freundlich isotherm is that it fails at high pressure of the gas.

we get

This equation applicable only when adsorbate substance form unimolecular layer on adsorbent surface.

Langmuir Adsorption Isotherm

Assumptions :

- 1. Gas is considered to behave ideally.
- 2. A solid surface is considered as homogenous but contains a fixed number of adsorption sites on surface.
- **3.** Each site adsorb a single molecule; means adsorption is monomolecular.
- **4.** Rate of adsorption = Rate of desorption.
- 5. There is no lateral overlap between adsorbed molecules.

x = aP

x = mass of gas adsorbed, m = mass of adsorbent, a & b are constants.

Let $\overline{m} = y = extent of adsorption$

$$\frac{ap}{1+bp}$$

$$\frac{P}{1} = \frac{1}{b}$$

ар

$$\frac{m}{x} = \frac{1+bp}{ap} = \frac{b}{a} + \frac{1}{ap}$$
.....(d)

A plot of m/x against 1/p gives a straight line with slope and intercept equal to 1/a and b/a respectively. At low pressure according to Eq. (c) x/m increases linearly with p. At high pressure according to Eq. (b) x/m becomes constant i.e. the surface is fully covered and change in pressure has no effect and no further adsorption takes place which is clear from the Figure-1.

Adsorption from solutions :

- 1. The extent of adsorption increases with increase in the surface area of adsorbent.
- 2. The extent of adsorption decreases with increase in temp.
- 3. The extent of adsorption is related to concentration of solution through this equation

Freundlich isotherm :
$$(x/m) = k(c)^{1/n}$$

where c is the equilibrium concentration of the solute in solution.

 $(n \ge 1)$

Highly specific in nature

Highly active under optimum Ph.

Influence of induce and poison

Temperature dependence here also is similar to that for adsorption of gases and in place of equilibrium pressure, we use equilibrium concentrations of the adsorbates in the solution.

Enzyme catalyst: Enzyme are complex nitrogenous organic compound which of emzyme CI produced by living plants and animal the chracterstics of enzyme catalyst are .

2.

4.

6.

- 1. Most highly efficient
- 3. Highly active under optimum temperature
- 5. Increase activity in presence of coenzyme
- Applications of adsorption :
 - In gas masks : Activated charcoal is generally used in gas masks to adsorb poisonous and toxic gases from air. These masks are commonly used by the miners because there are poisonous gases like CO, CH₄ etc. in the atmosphere in coal mines.
- 2. In dyeing of cloths : Mordants such as alums are used in dyeing of cloths. They adsorb the dye particles which, otherwise, do not stick to the cloths.
- 3. In dehumidizers : Silica gel is commonly used to adsorb humidity or moisture from air.
- 4. **Removal of colouring matter :** Many substances such as sugar, juice and vegetable oils are coloured due to the presence of impurities. They can be decolourised by placing them in contact with adsorbents like activated charcoal or fuller's earth.
- 5. Heterogeneous catalysis : The phenomenon of adsorption is useful in the heterogeneous catalysis. The metals such as Fe, Ni, Pt, Pd, etc. are used in the manufacturing processes such as Contact process, Haber process and the hydrogenation of oils. Their use is based upon the phenomenon of adsorption.
- 6. Refining Petroleum : Silica gel is used as adsorbent in petroleum refining.
- 7. **Chromatography :** It is a method for separation of component and is based on preferential adsorption column is very common device used.
- 8. Creating vacuum : High vacuum can be created by removing gas by adsorption.
- **9.** Adsorption Indicators : In volumetric analysis, adsorption indicator is used Surface of certain precipitates such as silver halide have the property of adsorbing some dye like eosin, fluorescein, etc In the case of precipitation titration (AgNO₃ vs NaCI) of the indicator is adsorbed at the end point producing a characteristic colour on the precipitate.
- **10.** In froth floatation process : (in metallurgy).
- 11. Softening of hard water : Ion exchange resins used for softening of hard water is based upon selective and competive adsorption of ions on resins.

$Na_2Z + Ca^{+2} \longrightarrow CaZ + 2 Na^+$

The organic polymers containing groups like -COOH, $-SO_3H$ and $-NH_2$ etc. possess the property of selective adsorption of ions from solution. These are quite useful in the softening of water.

Solved Examples

Example-1 A sample of charcoal weighing 6 g was brought into contact with a gas contained in a vessel of one litre capacity at 27°C. The pressure of the gas was found to fall from 700 to 400 mm. Calculate the volume of the gas (reduced to STP) that is adsorbed per gram of the adsorbent under the condition of the experiment (density of charcoal sample is 1.5 g cm³).
 Solution. The adsorption is taking place in a closed vessel hence if pressure falls there is correspondingly increase in volume constant, excess of the volume of the gas would be adsorbed.

 $\mathsf{P}_1\mathsf{V}_1=\mathsf{P}_2\mathsf{V}_2$



Section (B) : Catalysts

Catalysts : Berzillus in 1835 used the word catalyst first time for some substance which alter rate of chemical reaction and themselves remain chemically and quantitatively unchanged after the reaction and the phenomenon is known as catalysis.

Eg : Potassium chlorate when heated at 653K to 873K, it gives O₂, When MnO₂ is used in this reaction the O₂ is quickly at the low temperature hence MnO₂ is a catalyst

$$2\text{KCIO}_3 \rightarrow 2\text{KCI} + 3\text{O}_2$$

Homogeneous Catalysis : When catalysts and reactants are in same phase then the process is said to be homogeneous catalysis and

Eg: (i)
$$2SO_2(g) + O_2(g) \xrightarrow{NO(g)} 2SO_3(g)$$

(ii) $CH_3COOCH_3(\ell) \xrightarrow{HCl(\ell)} CH_3COOH(aq)$

(iii)
$$C_{12}H_{22}O_{11}(aq.) + H_2O(\ell) \xrightarrow{H_2SO_4(\ell)} C_6H_{12}O_6(aq.) + C_6H_{12}O_6(aq.)$$

Glucose Fructose

Heterogenous Catalysis : When catalysts and reactants are in different phases, then process a know as heterogenous catalysis and catalyst is called heterogeneous catalyst Pt(s)

Eg: (i)
$$2SO_2(g) + O_2(g \xrightarrow{Fr(s)} 2SO_3(g))$$

(ii) $N_2(g) + 3H_2(g) \xrightarrow{Fe(s)} 2NH_3(g)$
(iii) $4NH_3(g) + 5O_2(g) \xrightarrow{Pt(s)} 4NO(g) + 6H_2O(g)$
(iv) Vegetable oils $(\ell) + H_2(g) \xrightarrow{Ni(s)}$ Vegetable ghee (s).

Types of Catalysis

Type of Catalysis			
Positive	Negative	Auto	Induced
Substance which	Substance which decrease the	One of Product behave	When one reaction
increase the rate of	rate of chemical reaction	as catalyst for that	catalyst another
chemical reaction	$H_2O_2 \longrightarrow 2H_2O^+$	reaction and increase	reaction, than this
$2SO_2 + O_2 \xrightarrow{Pt} 2SO_2$	negative catalyse 2.120	the rate of reaction, than	phenomena is called
		this phenomena is called	as induced catalyses
		autocatalysis	$Na_2SO_3 \rightarrow Na_2SO_4$
		$CH_3COOC_2H_5 + H_2O$	
		\downarrow	
		CH3COOH + C2H5OH	

• Promotors/Activators : Substance which are not catalyst themselves but its presence can increase the catalytic activity of catalyst. A promotors increase the number of active sites on the surface Eg :

$$N_2 + 3H_2 \xrightarrow{Fe(catalyst)}{Mo(promotor)} 2NH_3$$

(iii)

(i)

 $CO + 2H_2 \xrightarrow{ZnO(catalyst)} CH_3OH$

• Catalytic Poisons/ Anti catalysts : Substance which are not catalyst themselves but whose presence decrease the activity of the catalyst. Poisoning is due to preferential adsorption of poison on the surface of the catalyst.

 $H_2 \xrightarrow[CO/H_2S(catalyst)]{} 2NH_3$ 2SO₂ + O₂ Platinised asbestos(catalyst) $\rightarrow 2SO_3$ As₂S₃(catalytic poisons)

(ii)

(iii)

$$\frac{\text{RCOCI} + \text{H}_2}{\text{BaSO}_4 (\text{ poisons catalyst})} \xrightarrow{\text{RCHO} + \text{HCI}}$$

Catalyst Inhibitor: Those substances which reduce the rate of a chemical reactions are known as . inhibitor.

e.g. H_3PO_4 and glycerol decrease the rate of decomposition of hydrogenperoxide.

Shape selective catalysis

Catalysis action depends upon.

- 1. Shape of the catalyst (Pore structure or cavities)
- Size of reactant and product. 2.

Zeolites

- 1. They are honey comb like structure i.e. shape selective catalysis.
- They are microporous aluminium silicate with 3D silicate in which silicon atom are replaced by 2. aluminum atoms i.e. Al - O - Si
- The pore size is generally in range 260-740 pm. 3.
- General formula of zeolite is Naxin [(AIO2)x (SiO2)y]. z H2O 4. Use of Zeolite
- 1. It is used as catalyst in petrolium industries for cracking hydrocarbon and isomerism i. e. ZSM-5.

 $R - OH \xrightarrow{ZSM-5} gasoline$

It convert alcohol directly into gasoline by dehydrating them to give a mixture of hydrocarbon. 2.

Characteristics of Catalysis :

- A Catalyst remains unchanged in mass and chemical compositions at the end of reactions. (i) However its physical state can be changed. Eg :
 - Granular MnO₂ during decomposition of KClO₃ is left as powder at the end of the reaction.
- (ii) Highly efficient : Finely devided state of catalyst is more efficient for the reactions because surface area increases and more adsorption take place.
- (iii) A catalyst cannot initiate the reaction. But some times the activation energy is so large that practically a reaction may not start until a catalyst lowers the activation energy significantly. For example, mixture of hydrogen and oxygen do not react at room temperature but the reaction occurs very rapid in presence of Pt black.

 $H_2 + O_2 \xrightarrow{\text{room temperature}}$ No reaction

$$H_2 + O_2 \xrightarrow{Pt black} H_2O_1$$

- Highly specific : Catalyst are generally specific in nature. A substance which act as a catalyst (iv) in a particular reaction, fails to catalyse other reaction.
- (vi) Catalyst cannot change equilibrium state but it helps to attain equilibrium quickly.
- A catalyst does not change the enthalpy, entropy and free energy of a reaction. (vii)
- Highly active under optimum temperature : There is a particular temperature at which the (viii) efficiency of a catalyst is maximum, this temperature is known as optimum temperature. On either side of the optimum temperature, the activity of catalyst decreases.(optimum range : 298 to 310 K).

- (ix) Highly active under optimum pH : Range is pH 5 to 7.
- (x) Influence of Inhibitor or poison.
- Adsorption Theory of Heterogeneous Catalyst : This theory explains the mechanism of heterogeneous catalyst. This theory is combination of two theory, intermediate compound formation theory and the old adsorption theory, the catalytic activity is localised on the surface on the catalyst. The mechanism involvs 5 steps.
 - (i) Diffusion of reactant to the surface of the catalyst of.
 - (ii) Adsorption of reactant molecules on the surface of the catalyst.
 - (iii) Formation of activated intermediate.
 - (iv) Formation of reactions product on the catalyst surface.
 - (v) Diffusion of reactions product from the catalyst surface or desorption.

Examples :

Let us consider addition of H_2 gas to ethlene in presence of Ni catalyst, the reaction takes places as follows.



• Factors Supporting Theory :

- (i) This theory explains the role of active centre, more free valency which provides the more space for more adsorption and concentration increases as a result of increase in rate of reaction.
- (ii) Rough surface has more active and pores, there will be more free valency so more will be rate of reaction.
- (iii) The theory explains the centre action of promotors which occupied the interstial void as a result surface area for the adsorption increases therefore rate of reaction increases



(iv) The theory explains the function of poisons or inhibitors. In poisoning preferential adsorption of poisons takes place on the catalyst, surface area for the adsorption on the catalyst decreases hence rate of reaction decreases.



Some Industrial Catalytic reactions

CHEMISTRY FOR JEE

S.N.	Process	Catalyst
1.	Haber's Process for the manufacture of ammonia $N_2 + 3H_2 \implies 2NH_3$	Finely divided iron, Molybdenum as promoter. Conditions : 200 atmospheric pressure and 450-500°C temperature.
2.	Ostwald's process for the manufacture of nitric acid. $4NH_3 + 5O_2 \rightarrow 4NO + 6H_2O$; $2NO + O_2 \rightarrow 2NO_2$ $4NO + 2H_2O + O_2 \rightarrow 4HNO_3$.	Platinised asbestos Temperature 300°C
3.	Lead chamber process for the manufacture of sulphuric acid $2SO_2 + O_2 \implies 2SO_3$; $SO_3 + H_2O \rightarrow H_2SO_4$.	Nitric oxide.
4.	Contact process for the manufacture of sulphuric acid. $2SO_2 + O_2 \implies 2SO_3$; $SO_3 + H_2SO_4 \rightarrow H_2S_2O_7$	Platinised asbestos or vanadium pentoxide(V_2O_5) Temperature 400–450°C.

Section (C) : Classification, Preparation & Purification of Colloid

• Colloid Solution :

Colloid State : A substance is said to be in colloidal state when the size of the particle of disperse phase is greater than particle of true solution and less than that of suspension solution particle, their range of diameters lie between 1 and 1000 nm $(10^{-9} \text{ to } 10^{-6} \text{ m})$.

Colloid solution : It is a heterogeneous system consisting of 2 phase :

(1) Disperse Phase (D.P) : The phase which is dispersed through the medium is called dispersed phase or discontinuous phase or internal phase.

(2) Dispersion Medium (D.M) : A medium in which colloidal particles are dispersed is called dispersion medium. It is also known as continuous phase or outer phase or external phase.

Colloidal solution = D.P. + D.M.

Ex. In Gold sol, Gold is D.P and water is D.M.

• Differentiating point of colloids :

(1) A colloid is a **heterogeneous system** in which one substance is dispersed (dispersed phase) as very fine particles in another substance called dispersion medium.

(2) The solution and colloid essentially differ from one another by particle size.

- * In a solution, the particles are ions or small molecules.
- * In a colloid, the dispersed phase may consist of particles of a single macromolecule (such as protein or synthetic polymer) or an aggregate of many atoms, ions or molecules.

(3) Colloidal particles are larger than simple molecules but small enough to remain suspended. They have a range of diameters between 1 and 1000 nm $(10^{-9} \text{ to } 10^{-6} \text{ m})$.

Classification of colloids :

1. On the basis of physical state of D.P. and D.M.

On the bases of physical state of D.P. and D.M. colloidal solution may be divided into eight system.

nples
e coloured glasses, and gem stones
s, cell fluids
ke, dust
ese, butter, jellies
hair cream
mist, cloud, insecticide sprays
ice stone, foam rubber
n, whipped cream, soap lather.

* Solution of gas in gas is not a colloidal system because it forms homogeneous mixture.

2. On the basis of D.M. : Colloidal solution are classified as

D.M.	Name of colloidal system
Water	Hydro sol or aqua sol
Alcohol	Alco sols
Benzene	Benzo sols

Air

Aero sols

- * Aquadag & oildag are colloidal solution of graphite in water & oil respectively.
- * Colloidal solution are often termed as sol.

3. On the Basis of interaction of D.P. for D.M. : There are two types-

(i) Lyophilic colloids / liquid loving sols / intrinsic colloid. The colloidal solution in which the particles of the dispersed phase have a great affinity (or love) for the dispersion medium, are called lyophilic colloids. These solutions are easily formed and the lyophilic colloids are reversible in nature. In case when water acts as the dispersion medium, the lyophilic colloid is called hydrophilic colloid. The common examples of lyophilic colloids are glue, gelatin, starch, proteins, egg albumin, rubber, etc.

(ii) Lyophobic colloids / solvent hating colloid / extrinsic colloid. The colloidal solutions in which there is no affinity between particles of the dispersed phase and the dispersion medium are called lyophobic colloids. Such solutions are formed with difficulty only by special methods. These sols are readily precipitated (or coagulated) upon addition of small amounts of electrolytes, by heating or by shaking and hence are not stable. Further, once precipitated, they do not give back the colloidal sol by simple addition of the dispersion medium. Hence these sols are also called irreversible sols. They need stabilising agents for their preservation. In case the dispersion medium is water, the lyophobic sol is called hydrophobic colloid. For example, the solution of metals like Ag and Au, hydroxides like AI (OH)₃, Fe(OH)₃, metal sulphides like As₂S₃ etc.

* **Lyophilic sols are more stable than lyophobic sols,** the additional stability is due presence of an envelope of the solvent layer (say water) around the colloidal particle, the process is known as hydration, To coagulate a hydrophilic sols we have to add a dehydrating agent in addition to electrolyte.

S.No.	Property	Lyophilic colloids	Lyophobic colloids
1	Ease of preparation	There are easily formed by direct mixing.	These are formed only by special methods
2	Reversible or irreversible nature	These are reversible in nature	These are irreversible in nature.
3	Particles nature	The particles of colloids are true molecules but are big in size	The particles are aggregates of many molecules
4	Stability	These are very stable	These are unstable and require traces of stabilizers
5	Action of electrolytes	No effect	The addition of small amount of electrolytes causes precipitation (called coagulation) of colloidal solution.

DISTINCTION BETWEEN LYOPHILIC AND LYOPHOBIC COLLOIDS

S.No.	Property	Lyophilic colloids	Lyophobic colloids
6	Charge on particles	The particles do not carry any charge.	The particles move in a specific direction either towards anode or cathode depending upon their charge in an electric field
7	Hydration	The particles of colloids are heavily hydrated due to the attraction for the solvent.	The particles of colloids are not appreciably hydrated.
8	Viscosity	It is higher than that of dispersion medium	It is nearly same as that of dispersion medium
9	Tyndall effect	They do not show tyndall effect	They show tyndall effect.
10	Surface tension	Lower than dispersion medium	It is nearly same as that of dispersion medium
11	Coagulation of	Precipitated by high concentration of	Precipitated by low concentration of
	precipitation	electrolyte	electrolyte
12	Migration in electric	May or may not migrate as they may	Migrate toward anode or cathode as these
12	field	not carry charge	particles carry charge
13	Example	Mostly organic nature; Starch and Gelatin	Inorganic nature; Transition metal salt in water, gold etc., Metal solution

4. On the basis of chemical composition:

Inorganic Colloids

(i) Metal sols : Cu, Ag, Au, Pt Sols.

(ii) Non Metal sols : S, I₂, Graphite

(iii) Sol of oxide and hydroxide : SnO₂, TiO₂, Fe₂O₃, Fe(OH)₃, Al(OH)₃, Cr(OH)₃

(iv) Salt Sol - AgBr, AgI, As₂ S₃, etc.

Organic Colloids

(i) Homopolar sol - In this type of colloid, particles carry similar type of charge. eg. Sol of rubber in benzene which contain - ve charge colloidal particle of latex.

(ii) Hydroxy Sol- Starch sol

5. On the basis of charge on particles

(i) positive Sol

(a) Metal Oxide & Hydroxide - SnO₂, TiO₂, Fe₂O₃, AI(OH)₃, Fe(OH)₃, Cr(OH)₃.

(b) Basic Dyes Methylene blue, vismark brown.

(ii) negative Sol -

(a) Metal sol - Ag, Au, Pt, Cu (b) Acidic dye - congo red, eosin

(c) Sulphide Sol- CdS, HgS, As₂S₃, Sb₂S₃.

(d) Natural sol - Blood, clay, charcoal, latex rubber, dust particle in water, starch, carbon particle in smoke, gum.

6. Other type of colloids : Multimolecular, macromolecular and associated colloids

- **Multimolecular colloids :** In this type, the particles consist of an aggregate of atoms or small molecules size less than 1 nm. For example, sols of gold atoms and sulphur (S₈) molecules. In these colloids, the particles are held together by van der Waal's forces.
- **Macromolecular colloids :** In this types, the particles of the dispersed phase are sufficiently big in size (macro) to be of colloidal dimensions. These macromolecules forming the dispersed phase are generally polymers having very high molecular masses. These colloids are quite stable and resemble true solutions in many respects. Naturally occuring macromolecules are starch, cellulose, proteins, enzymes, gelatin, etc.
- **Associated colloids (Micelles):** These are the substances which behave as normal strong electrolytes at low concentration but behave as colloidal particles at higher concentration. These associated particles are also called micelles. Ex. Soap.

Micelles :

 \Rightarrow Micelles are relatively small, spherical structures composed of any where for a few to few thousand molecule that attract one another to reduce surface tension within a membrane of cell. The formation of micelles takes place only at particular temp, that temperature is called **Kraft**

temperature.

The concentration above which micelle formation becomes appreciable is termed is **critical micelles concentration**. its value depend upon natures of D.P. and D.M. eg. Surface active agent (surfactants, which decrease the surface tension) like soaps and detergents form micelle beyond CMC ($\sim 10^{-3}$ mol/litre for soaps).

* Usually longer the hydrophobic chain, smaller is its CMC.

* Also CMC increase with decreasing polarity of the D.M.

* The micelles 'formation takes place only above a particular temperature called as Kraft Temperature (T_k) .

*At CMC, the micelles are spherical in shape, but that start flattening with increase in concentration and ultimately form sheet or film like structures which have a thickness of two molecules. These are called **lamelar micelles** or **McBain Micelles**.

Mechanism of micelle formation: Let us take the example of soap solutions. Soap is sodium salt of a higher fatty acid and may be represented as $RCOO^-Na^+$ e.g., sodium stearate viz.CH₃(CH₂)₁₆COO⁻ Na⁺ which is a major component of many bar soaps. When dissolved in water, it dissociates into RCOO- and Na+ ions. The RCOO⁻ ions, however, consist of two parts i.e., long hydrocarbon chain R (also called non-polar 'tail') which is hydrophobic (water repelling) and the polar group COO⁻ (also called polar-ionic 'head') which is hydrophobic (water loving). The RCOO⁻ lons are, therefore, present on the surface with their COO⁻ groups in water and the hydrocarbon chains R staying away from it, and remain at the surface, but at higher concentration these are pulled into the bulk of the solution and aggregate in a spherical form with their hydrocarbon chains pointing towards the centre with COO⁻ part remaining outward on the

surface. An aggregate thus formed is known as '**lonic micelle**'. These micelles may contain as many as upto 100 such ions.

Aggregation of RCOO- ions to form an ionic micelle. Similarly, in case of detergents, e.g., sodium lauryl sulphate viz. $CH_3(CH_2)_{11}SO_4^-$ Na⁺, the polar group is $-SO_4^-$ along with the long hydrocarbon chain. Hence, the mechanism of micelle formation is same as that of soaps.



Example of micelles :

(i) Sodium stearate C₁₇H₃₅COO⁻Na⁺(Soap).

(ii) Sodium lauryl sulphate CH₃ [CH₂]₁₁ SO₄⁻ Na⁺ (Detergent).

(iii) Cetyl trimethyl ammonium bromide (Detergent). CH₃(CH₂)₁₅N⁺(CH₃)₃Br⁻.

The Cleansing Action of Soaps : It has been mentioned earlier that a micelle consists of a hydrophobic hydrocarbon like central core. The cleansing action of soap is due to these micelles, because oil and grease can be solubilised in their hydrocarbon, like centres which are not otherwise soluble in water. This is shown diagrammatically in Figure The dirt goes out along with the soap micelles.



(a) A sodium stearate molecule

(b) The simplified representation of the molecule that shows a hydrophilic head and a hydrophobic tail

- (c) Grease (oily substance) is not soluble in water
- (d) When soap is added to water, the non-polar tails of soap molecules dissolve in grease

(e) Finally, the grease is removed in the form of micelles containing grease.

*Surfactants : They can be ionic as well as non-ionic. The ionic are soaps and detergent. The surfactant gets adsorbed at the interface between the dispersed droplets and dispersion medium in the form of mono molecular layer and lowers the interfacial tension between oil and water so as to facilitate the mixing of two liquids.

Preparation of lyophobic colloidal sols :

[A] Condensation methods :

In these methods particles of atomic or molecular size are induced to combine to form aggregates having colloidal dimensions. For this purpose chemical as well as physical methods can be applied.

(a) Chemical methods. Colloidal solutions can be prepared by chemical reactions leading to formation of molecules by double decomposition, oxidation, reduction or hydrolysis. These molecules then aggregate leading to formation of sols.

(i) Double decomposition : When a hot aqueous dilute solution of arsenous oxide (As_2O_3) is mixed with a saturated solution of H_2S in water, a colloidal sol of arsenous sulphide (As_2S_3) is obtained.

As₂O₃(in hot water) + $3H_2S$ (saturated solution in H₂O) CH(OH)COOK CH(OH)COOK As₂S₃(sol) + $3H_2O$

 $\dot{C}H(OH)COO(SbO)_{+3H_2S} \rightarrow \dot{C}H(OH)COOH_{+Sb_2S_3} \text{ (orange sol) + 2H_2O}$

(ii) Oxidation : A colloidal sol of sulphur is obtained by passing H₂S into a solution of sulphur dioxide.

 $SO_2 + 2H_2S(\text{saturated solution in } H_2O) \xrightarrow{\text{Oxidation}} 3S(\text{sol}) + 2H_2O$

Sulphur sol can also be obtained when H₂S is bubbled through Br₂ water or nitric acid (oxidizing agent).

$$2H_2S$$
 (aq.) + Br₂ (aq.) \longrightarrow $2HBr$ (aq.) + S (sol).

or by bubbling O_2 (g) through a solution of H_2S : $2H_2S$ (aq.) + O_2 (g) $\longrightarrow 2H_2O$ (l) + 2S (sol). (iii) **Reduction :** Colloidal sol of metals like gold, silver solution are obtained by following method.

 $2 \text{AuCl}_3 + 3 \text{HCHO} + 3\text{H}_2\text{O} \longrightarrow 2\text{Au(sol)} + 3\text{HCOOH} + 6\text{HCl}.$

(purple of cassius)

 $2AuCl_3 + 3SnCl_2 \xrightarrow{\text{Reduction}} 3SnCl_4 + 2Au(sol)$

AgNO₃ + tannic acid $\xrightarrow{\text{Reduction}}$ silver sol.

 NH_2NH_2 can also be used as reducing agent. *Sol of gold is also known as purple of cassius. (iv) Hydrolysis : A colloidal sol of metal hydroxides like Al(OH)₃ or Cr(OH)₃ is obtained by boiling a dilute solution of FeCl₃, AlCl₃ or CrCl₃.

 $FeCI_3 + 3H_2O \longrightarrow Fe(OH)_3$ (sol) + 3HCI; $AICI_3 + 3H_2O \longrightarrow AI(OH)_3$ (sol) + 3HCI The colloidal sol of sillicic acid is also obtained by hydrolysis of dilute solution of sodium silicate with

hydrochloric acid. Na₄SiO₄ + 4HCl \longrightarrow Si(OH)₄ (sol) + 4NaCl.

(b) Physical methods : The following physical methods are used to prepare the colloidal solutions.

(i) By Exchange of solvent : When a true solution is mixed with an excess of the other solvent in which the solute is insoluble but solvent is miscible, a colloidal sol is obtained. For example,

when a solution of sulphur in alcohol is poured in excess of water, a colloidal sol of sulphur is obtained.
when a solution of phenolphthalein in alcohol is poured in excess of water a white sol of phenolphthalein is found.

•Phenolphthalein, I₂, sulphur sol can be prepared by this methods.

(ii) Excessive cooling : Molecules of certain substance condense together on excess cooling to form colloidal size particle. The colloidal sol of ice in an organic solvent such as CHCl₃ or ether can be obtained by freezing a solution of water in the solvent. The molecules of water which can no longer be held in solution separately combine to form particles of colloidal size.

(iii) By condensing vapours of a substance into solvent : Substance like sulphur and Hg in water are prepared by passing their vapours in cold water containing small amount of stabilising agent like ammonium nitrate.

[B] Dispersion Methods : In these methods large particles of the substance are broken into particles of colloidal dimensions in the presence of dispersion medium. These are stabilized by adding some suitable stabilizer. Some of the methods employed are given below :

(a) Mechanical dispersion (By colloidal mill):

J

Substance is first finely powdered.

It is shaken with the D.M. to form a suspension. \downarrow This suspension is passed through a colloidal mill.

The simplest type of colloidal mill is disc mill which consists of two metal discs nearly touching each other & rotating in opposite.

 \downarrow

J

Direction at a high speed (7,000 revolutions per min.).

Suspension Hollow shaft Discharge Discharge Fig. : A colloid mill

The suspended particles are broken to produce colloidal size particle.

* This method is used to prepare printing ink.

(b) Electrical disintegration or Bredig's Arc method : This

process involves dispersion as well as condensation. Colloidal

sols of less reactive metals such as gold, silver, platinum,

copper, lead etc., can be prepared by this method. In this

method, electric arc is struck between electrodes of the metal

immersed in the dispersion medium as shown in fig. The intense

heat produced vaporises the metal, which then condenses to

form particles of



colloidal size by surrounding cooling mixture (ice).

*A slight trace of KOH is added in water to stabilized colloidal solutions.

(c) Ultrasonic dispersion : Ultrasonic vibration (having frequency larger than audible range) can bring about the transformation of coarse suspension or liquids like oil, mercury etc. into colloidal range.

*This is the latest method for preparation of metal oxides and metal sulphide sols from their coarse suspension.

*It is a suitable technique for oils also. This method also comprises both dispersion and condensation.

(d) Peptization: The term has originated from the digestion of proteins by the enzyme pepsin. Peptization may be defined as (the process of converting a precipitate into colloidal sol by shaking it with dispersion medium in the presence of a small amount of electrolyte).

The electrolyte used for this purpose is called peptizing agent. This method is applied, generally, to convert a freshly prepared precipitate into a colloidal sol. During peptization, the precipitate adsorbs one of the ions of the electrolyte on its surface.

The ion adsorbed on the surface is common either with the anion or cation of the electrolyte. This causes the development of positive or negative charge on precipitates which ultimately break up into smaller particles having the dimensions of colloids.



For example :

(i) When freshly precipitated Fe(OH)₃ is shaken with aqueous solution of FeCl₃ (peptizing agent) it adsorbs Fe³⁺ ions and thereby breaks up into small-sized particles.



Charge Colloidal particals of Fe $(OH)_3$

(ii) Freshly prepared stannic oxide on treatment with a small amount of dilute hydrochloric acid forms a stable colloidal sol of stannic oxide, SnO₂; Sn⁴⁺.

 $SnO_2 + 4HCI \rightarrow Sn^{4+} + 2H_2O + 4CI^-$; $SnO_2 + Sn^{4+} \rightarrow SnO_2 / Sn^{4+}$.

(iii) Freshly precipitated silver chloride can be converted into a colloidal sol by adding a small amount of hydrochloric acid, AgCl : Cl^{-} .

(iv) Cadmium sulphide can be peptised with the help of hydrogen sulphide, CdS : S²⁻.

Purification of Colloidal Sols : The colloidal sols obtained by various methods are impure and contain impurities of electrolytes and other soluble substances. These impurities may destabilise the sol. Hence, they have to be removed. A very important method of removal of soluble impurities from sols by a semipermeable membrane is known as dialysis.

A. Dialysis : It is a process of removing a dissolved substance from a colloidal solution by means diffusion through suitable membrane. Since particles in true solution (ions or smaller molecules) can pass through

animal membrane or parchment paper or cellophane sheet but colloidal particle do not, the appratus used for this purpose is called Dialyser.

A bag of suitable membrane containing the colloidal solutions is suspended in a vessel through which fresh water continously flow. The molecules and ions (crystalloids) diffuse through membrane into the outer water & pure colloidal solution is left behind.



Fig. : An apparatus for electrodialysis.

Movement of ions across the membrane can be expedited by applying electric potential through two electrodes as shown in fig.

This method is faster than simple dialysis and is known as **Electrodialysis.**

*The most important applications of dialysis is in the purification of blood in the artificial kidney machine. In case of kidney failure, blood cannot be purified. Under such condition, the blood is separated from dissolved toxic impurities by dialysis and re-introduced in the bloods stream.

*Dialysis is not applicable for non-electrolytes like glucose, sugar, etc.

B. Ultra Filtration : In this method, colloidal sols are purified by carrying out filtration through special type of graded filters called ultra-filters. These filter papers allow only the electrolytes to pass through. These filter papers are made of particular pore size by impregnating with colloidal solution and subsequently hardened by soaking in formaldehyde collodion. In order to accelerate the filtration through such filter papers, increased pressure or section is employed.

Solved Examples.

Example-2	When SO ₂ is bubbled into H_2S gas, colloidal sol is formed. What type of colloidal sol is it ?	
Solution.	$2H_2S + SO_2 \longrightarrow 2H_2O + 3S$ (colloidal). Lyophobic colloidal sol of sulphur is formed.	
Example-3 Solution.	A reddish brown positively charged sol is obtained by adding small quantity of FeCl ₃ solution to freshly prepared and well washed $Fe(OH)_3$ precipitate. How does it take place ? It is due to adsorptions of Fe^{3+} ions on the surface of $Fe(OH)_3$ which gives colloidal sol.	
	$Fe(OH)_3$ (ppt.) + Fe^{3+} (ions adsorbed) \longrightarrow [$Fe(OH)_3$] Fe^{3+} (colloidal sol).	
Example-4	Suppose we have a cube of 1.00 cm length. It is cut in all three directions, so as to produce eight cubes, each 0.50 cm on edge length. Then suppose these 0.50 cm cubes are each subdivided into eight cubes 0.25 cm on edge length, and so on. How many of these successive subdivisions	
Solution.	We find that every division in two equal halves also reduces the size of edge lengths to one half. 1	
	In first subdivision 1 cm is reduceds to 0.5 cm = $\overline{2}$ cm.	
	In second subdivision 0.5 cm is reduced to 0.25 cm = $\frac{1}{4}$ cm = $\left(\frac{1}{2}\right)^2$ cm	
	$\left(\frac{1}{2}\right)^n$	
	In n subdivision 1 cm is reduced to $\lfloor 2 \rfloor$. Size of colloidal particles lies between 1 to 1000 mm. Thus to make n subdivision required particle size may be attained.	
	$\left(\frac{1}{2}\right)^n$ = 100 nm = 100 × 10 ⁻⁹ m = 100 × 10 ⁻⁷ cm.	
	$\therefore \qquad n \log 2 = 5$	
	n × 0.3010 = 5. 5	
	$n = \frac{1000}{0.3010} = 16.61 = 17$ subdivisions are required for dimension of 100 nm.	

Important properties of colloidal sols :

• Heterogeneous character :

- Colloidal sols are heterogeneous in character as they consist of two phases. (a) dispersed phase and (b) dispersion medium.
- Visibility : Due to scattering caused by the colloidal particles, it will appear as a bright spot moving randomly.

• **Filterability :** Colloidal particles pass through an ordinary filter paper. However, the particle do not pass through other fine membranes.

• **Colligative Properties :** Colloidal sols show the colligative properties viz. relative lowering of vapour pressure, elevation in boiling point, depression in freezing point and osmotic pressure. However, due to high average molecular masses of colloidal particles, mole fraction of the dispersed phase is very low. Hence, the values of the colligative properties observed experimentally are very small. Only osmotic pressure measurements are used in determining the molecular mass of polymers.

• Optical Properties-Tyndall effect : Tyndall, in 1869, observed that if a strong beam of light is passed through a colloidal sol placed in a dark place, the path of the beam gets illuminated. This phenomenon is called Tyndall effect, which is due to the scattering of light by the colloidal particles. The illuminated path of beam is called Tyndall cone. This phenomenon is due to scattering of light from the surface of colloidal particles. In a true solution there are no particles of sufficiently large diameter to scatter light & hence the beam is invisible.



*The intensity of scattered light depends on the difference between the refractive indice of the D.P and D.M., In lyophobic colloids, this difference is appreciable and therefore the tyndal effect is quite well defined but in lyophilic sols the difference is very small and the tyndal effect is very weak. Thus in sols of silicic acid, blood serum, albumin, etc. there is little or no tyndal effect.

Example of Tyndall Effect

- \rightarrow Blue colour of sky and sea water.
- \rightarrow Visibility of tail of comets.
- \rightarrow Light thrown from a projector in cinema hall.
- \rightarrow Appearance of dust particle in a semi darked room.

• Application of Tyndall Effect :

- (i) In making ultramicroscopes.
- (ii) In finding heterogenity of solution.

Solved Examples

Example-5 Solution.	Under what conditions is Tyndal effect observed ? Tydalls effect is applicable when :
	(a) The diameter of the dispersed particles is not much smaller than the wavelength of the light used.
	(b) The refractive indices of the dispersed phase and the dispersion medium must differ greatly in magnitude.
Example-6	In the lower layer of the atmosphere, there is a great deal of dust. When the weather is fine, it is possible to see the magnificent red colour of the setting sun. What have these observation to do with colloids ?
Solution.	Dust in the atmosphere is often colloidal. When the sun is low down on the horizon, light from it has to pass through a great deal of dust to reach your eyes. The blue part of the light is scattered

away from your eyes. You see the red part of the spectrum, which remains. Red sunsets are the Tyndall effect on a large scale.

• Mechanical Properties :

Brownian movement: Robert Brown, a botanist, discovered in 1827 that pollen grains placed in water do not remain at rest but move about continuously and randomly. Later on, this phenomenon was observed in case of colloidal particles when they were seen under an ultramicroscope. The particles were seen to be in constant zig-zag motion as shown in fig. This zig-zag motion is

called Brownian movement.



Brownian movement arises because of the impact of the molecules of the dispersion medium with the colloidal particles.

Dependence: (i) Size of colloidal particles

Mobility ∝ Size of the particle (*ii*) Viscosity of solution

Mobility ∝ Viscosity

(iii) Temperature

Mobility < Temperature

• Factors Affecting Brownian Movement :

(i) If particles is large then brownian movement becomes less.

(ii) Brownian movement increases with increasing temperature.

(iii) The brownian movement does not change with time & remains same for months or even for a year.

• Important :

(i) In confirmation of kinetic energy.

- (ii) Determination of Avogadro numbers.
 - (iii) Stability of colloidal solution : Brownian movement does not allow the colloidal particles to settle down to gravity & thus is responsible for their stability.

• Electrical Properties (Electrophoresis) :

The existence of charge on colloidal particles is confirmed by electrophoresis experiment. When electric potential is applied across two platinum electrodes dipping in a colloidal solution, the colloidal particles move towards one or the other electrode. The movement of colloidal particles under on applied electric potential is called electrophoresis.

Positively charged particles move towards the cathode while negatively charged particles move towards the anode. Depending upon the direction a movement of particles towards cathode or anode electrophoresis can be called 'cataphoresis' or 'Anaphoresis'

Arsenious sulphide, gold, silver and platinum particles in their respective colloidal sole are negatively charged while particles of Fe(OH)₃, Al(OH)₃ are positively charged.



(Fig. : A set up for electrophoresis.)

The colloidal solution is placed in a U-tube fitted with platinum electrodes. On passing an electric current, the charged colloidal particles move towards the oppositely charged electrode. Thus, if arsenic sulphide sol is taken in the U-tube, in which negatively charge particle of arsenic sulphide move towards the anode. *Earlier this process was called cataphoresis because most of the colloidal sols studied at that time were positively charged and moved towards cathode.

- Electro osmosis: When movement of colloidal particles is prevented by some suitable means (porous diaphragm or semi permeable membrenes), it is observed that the D.M. begins to move in an electric field. This phenomenon is termed electrosmosis.
- Sedimentation potential or Dorn potential : When the charged colloidal particles are made to settle down under centrifugal field, there occurs a charge separation and a potential difference is developed. This effect is called **Dorn effect** and the potential difference thus developed is called Dorn potential or sedimentation potential. This process is **reverse of electrophoresis**.
- **Isoelectric point :** The H⁺ concentration at which the colloidal particles have no charge is known as the isoelectric point. At this point stability of colloidal particles becomes very less & do not move under influence of electric field.
- Streaming potential : A potential difference is developed across a porous partition when the dispersion medium of a charged colloid is forced through it. This is called Streaming potential. This process is reverse of electro-osmosis.
- **Charge on colloidal particles :** Colloidal particles are either positively charged or negatively charged. This charge is due to preferential adsorption of either positive or negative ions on their surface. There is adsorption of common ion present in excess.
 - Fe(OH)₃ sol prepared by the hydrolysis of FeCl₃ solution adsorbs Fe³⁺ and this is positively charged.

FeCl₃ + 3H₂O
$$\longrightarrow$$
 Fe(OH₃) + 3HCl; Fe(OH)₃ + FeCl₃ \rightarrow Fe(OH)₃ $\stackrel{!}{\downarrow}$ Fe³⁺ : 3Cl⁻
Fixed part Diffused part

Positive charge on colloidal sol is due to adsorption of Fe^{3+} ion (common ion between $Fe(OH)_3$ and $FeCl_3$).

 \bullet As_2S_3 colloidal sol is obtained when As_2O_3 is saturated with H_2S :

 $\mathsf{As}_2\mathsf{O}_3 + \mathsf{3H}_2\mathsf{S} \to \mathsf{As}_2\mathsf{S}_3 + \mathsf{3H}_2\mathsf{O}.$

 As_2S_3 adsorbs S^{2-} ions (common between H_2S and As_2S_3 and thus is negatively charged).

 $As_2S_3 + H_2S \rightarrow As_2S_3$ | $S^{2-}: 2H^+$.

Agl in contact with AgNO₃ forms positively charged colloidal sol due to adsorption of Ag⁺ ion.
 Agl + AgNO₃ → [Agl]Ag⁺ : NO₃⁻, Agl in contact with KI forms negatively charged colloidal sol due to

adsorption of I⁻ ion AgI + KI \rightarrow AgI \downarrow I⁻ : K⁺.

• SnO₂ in acidic medium forms positively charged colloidal sol due to adsorption of Sn⁴⁺ formed.

 $SnO_2 + 4H^+ \rightarrow Sn^{4+} + 2H_2O$ $SnO_2 + Sn^{4+} \rightarrow SnO_2^{\dagger}$ Sn^{4+} SnO_2 in alkaline medium forms negatively charged colloidal sol due to adsorption of SnO_3^{2-} formed.

 $SnO_2 + 2OH^- \rightarrow SnO_3^{2-} + H_2O$ $SnO_2 + SnO_3^{2-} \rightarrow SnO_2^{-1}$ $SnO_2^{-1} + SnO_3^{2-} \rightarrow SnO_2^{-1}$

Charge on colloidal particle may be due to some other reasons also e.g. *Due to electron capture by collidal particle during electro dispersion of metal. *Due to self dissociation.

• Electric Double Layer Theory or Helm-holtz Electric double layer : Electric double layer Theory



CHEMISTRY FOR JEE

- The combination of two layer of opposite charges (+ve and –ve charge) around colloidal particle is called Hebnholtz electrical double layer.
- First layer of ions is firmly held and is termed as fixed layer.
- Secondary layer is mobile and is termed as diffused layer.
- The charges of opposite sign on fixed and diffused parts of double layer result in difference in potential between these layers.
- This potential difference between the fixed layer and diffused layer of opposite charges is called electrokinetic potential or zeta potential.

Section (D) : Coagulation, Protection & Application of colloid

• Coagulation/Flocculation : This process of aggregation of colloidal particles into an insoluble precipitate by the addition of some suitable electrolyte is known as coagulation. At lower concentration of electrolytes, the aggregation of particles is called flocculation that can be reversed on shaking while at higher concentration of electrolyte, coagulation takes place and the same cannot be reversed simply by shaking. The stability of the lyophobic colloids is due to presence of charge on colloidal particles. If, somehow, the charge is removed, the particles will come near to each other to form aggregates and settle down under the force of gravity.

Coagulation value or Flocculation value : It needs to be noted that the coagulation of a colloidal solution by an electrolyte does not take place until the added electrolyte has certain minimum concentration in the solution. The minimum concentration of electrolyte in millimoles required to cause coagulation of one litre of colloidal solution is called coagulation value. It is express in terms of millimoles/litre.

millimoles of electrolyte volume of sol in litre

Coagulation value = volume of sol in lite

Coagulation of lyophobic sols can be carried out by the following methods.

(i) By electrophoresis

(ii) By mutual precipitation : It is a process in which oppositely charged sol are mixed in proper proportion to neutralise the charge of each other causing coagulation of both the sol.

Example : Positively charged Fe(OH)₃ and negatively charged As₂S₃ colloidal particle containing sol on mixing get coagulated.

(iii) By Prolonged Dialysis : On prolonged dialysis, traces of the electrolyte present in the sol are removed almost completely and the colloidal unstable and ultimately coagulate.

(iv) By Boiling : Sols such as sulphur and silver halides dispersed in water may be coagulated by boiling because increased collisions between sol particle and the water molecule removed the adsorbed electrolytes. This takes away the charge from the particles and helps them to coagulate.

(v) By cooling : Certain sol can also be coagulated by lowering temperature. For example, accumulation of cream on the surface of milk on cooling. This is because at lower temperature the dispersion medium molecules do not exert sufficient force on to the dispersed particles and hence the Brownian motion becomes less effective.

(vi) By the addition of electrolyte : When excess of an electrolyte is added, the colloidal particles are precipitated.

• Comparision of relative coagulating power of two electrolyte for the same colloidal solution :

The coagulation value decrease with increase in charge of the coagulating ion.

			. I				
Coagulating power \propto coagulation value .							
coagulating	power	of	electrolyte	А	_ coagulation	value	В
coagulating	power	of	electrolyte	В	coagulation	value	Α

• Factor-Affecting Coagulations : (i) Nature of sols : The lyophobic colloid can easily coagulate because it is a less stable colloid, but lyophilic colloids coagulate hardly by the addition of electrolyte due to protective layer of D.M. surrounding the colloidal particle.

(ii) Nature of electrolyte : In equimolar electrolyte, strong electrolyte have greater coagulating power than weak electrolyte. Example : 0.1M NaCl > 0.1M CH₃COOH.

• Hardy-Schulze Rule : According to this rule greater is the valency of coagulating ion, greater its power to cause precipitation. This is known as Hardy-Schulze rule.

In case of positive charged sol, the coagulating power of anion is in the order of $[Fe(CN)_6]^{4-} > PO_4^{3-} > SO_4^{2-} > Cl^-$

In case of negative charged sol, the coagulating power of cation is in the order of $AI^{3+} > Ba^{2+} > Na^+$.

• The coagulating power of bivalent ion is 20-80 times higher than monovalent ion and coagulating power of trivalents is many times more than bivalent.

Solved Examples.

Example-9 The particles of a particular colloidal solution of arsenic trisulphide (As₂S₃) are negatively charged. Which 0.0005 M solution would be most effective in coagulating this colloidal solution. KCl, MgCl₂, AlCl₃ or Na₃PO₄? Explain.
 Solution. Since As₂S₃ is a negatively charged colloidal sol hence positively charged ion will cause its

Since As₂S₃ is a negatively charged colloidal sol hence positively charged ion will cause its coagulation. By Hardy-Schulze rule "greater the charge on ion, greater the coagulating power to coagulate oppositely charged colloidal sol", hence out of K⁺, Mg²⁺, Al³⁺ and Na⁺, Al³⁺ would be most effective.

Protective colloidal sols : Lyophilic colloidal sols are much more stable than lyophobic colloidal sols. This is due to the extensive solvation of lyophilic colloidal sols, which forms a protective layer outside it and thus prevents it from forming associated colloids. Lyophobic sols can easily precipitate by addition of small amount of an electrolyte. They can be prevented from coagulation by previous addition of some lyophilic colloid. This is due to formation of a protective layer by lyophilic sols outside lyophobic sols.

Process of protecting the lyophobic colloid solution from precipitation by an electrolyte due to previous addition of some lyophilic colloid is called protection of colloid and lyophilic colloidal sols are called protective sols.

Eg : Gelatin, Sodium caseinate, Egg albumin, Gum arabic, Potato starch etc.,

Gelatin (lyophilic) protects gold sol (lyophobic) colloids is expressed in terms of gold number.

Gold Number : Zsigmondy (1901) introduce a term called gold number it is defined as "the minimum amount of the protective colloid in milligrams which when added to 10 ml of a standard gold sol is just sufficient to prevent a colour change from red to blue on the addition of 1 ml of 10% sodium chloride solution. It may be noted that smaller of the gold number, greater will be protecting power of the protective colloid.

Protecting power $\propto \frac{1}{\text{gold number}}$

Gold Number =

Uses of protective action :

(i) Gelatin is added in the preparation of ice cream to protect the particle of ice.

(ii) Protargol and Argyrol, is a silver sol protected by organic material used as eye drop.

Applications of Colloids : Colloids including emulsions find a number of uses in our daily life and industry. Some of the uses are given below.

- **1. In medicines :** A wide variety of medicinal and pharmaceutical preparations are emulsions. Colloidial medicines are easily adsorbed by the body tissue because of large surface area.
 - * Colloidal antimony is used in curing kalaazar.
 - * Milk of magnesia, an emulsion, is used for stomach disorder.
 - * Colloidal gold is used for intramuscular injection.
 - * Colloidal sulphur are used as Germicides.
 - * Argyrol is a silver sol used as an eye lotion.
 - * Colloidal Fe(OH)₃ is given to arsenic poisoning patients as it adsorbs arsenic and then gets omited out.
- 2. Tanning
- 3. Photographic plate & Film
- 4. Rubber plating
- 5. Sewage disposal
- 6. Cottrell smoke precipitator :

Smoke is a dispersion of negatively charged colloidal particles of carbon in air and can be made free of these colloidal particles by passing it through cottrell precipitator as shown in fig. installed in the chimney of an industrial plant. It consists of two metal discs charged to a high potential. The carbon particles get discharged and precipitate, while gases come out from the chimney.

7. Formation of deltas: The river water contains colloidal particles of sand and clay which carry negative charge. The sea water contains +ve ions such as Na⁺, Mg²⁺, Ca²⁺, etc. As the river water meets sea water, these ions discharge the sand or clay particle which are precipitated in the form of delta.

8. Artificial rain

- 9. Stop bleeding from a cut
- 10. Stop Screen
- 11. Preparation of nano-materials
- **12.** In disinfectants : The disinfectants such as dettol and lysol give emulsions of the oil-in-water type when mixed with water.

STEM TECHNOLOGY : The size and shape of the colloidal particles is determined with the help of an electron microscope which has much more resolving power (of the order of 10^{-12} m.) The different techniques used to study the colloidal particles are :

(i) Scanning electron microscope (SEM)

- (ii) Transmission electron microscope (TEM) and
- (iii) Scanning transmission electron microscope (STEM).

Section (E) : Emulsion, Micelle & Gel

• **Emulsions :** Pair of immiscible liquid is called emulsion. Emulsion are unstable and some time they are separated into two layers on keeping still, for the stabilising of an emulsion, a third component is added called emulsifying-Agent form an interfacial film between D.P. and D.M.

Emulsion droplets are bigger than sol particles and can be seen under an ordinary microscope and sometimes even with a magnifying glass.

Example : Milk is an emulsion in which liquid fat is D.P. and liquid water is D.M. and **casein** is emulsifying agent.

Demulsification : The separation of an emulsion into its constituent liquids is called demulsification. Various techniques employed for this are freezing, boiling, centrifugation, electrostatic precipitation or chemical methods which destroys the emulsifying agents.

Demulsification can be brought about by :

(i) Freezing (ii) Heating

(iii) Centrifugal action (Separation of cream of milk done by centrifugation).

(iv) Removal of emulsifiers by adding a better solvent for them like alcohol, phenol etc, called demulsifiers. **Types of emulsions :** Depending on the nature of the dispersed phase, the emulsions are classified as :

- (a) Oil in water emulsions (b) Water in oil emulsions
- **Inversion of phase :** The conversion of emulsion of oil in water (o/w) into water in oil (w/o) or vice versa is called the inversion of phase.

[A] Identification of the type of emulsion : These two types may be identified by :

- **Dilution test :** An emulsion can be diluted with any amount of the dispersion medium, while the dispersed liquid, if added, forms a separate layer. Thus if a few drops of water added to the emulsion are soluble in it, it is oil in water type and if immiscible, it is water in oil type.
- **Dye test :** If a small amount of oil soluble dye gives a uniform colour to the emulsion, it is water in oil type otherwise it is oil in water type.
- Electrical conductivity test : If conductivity of emulsion increases significantly by adding a very small amount of electrolyte, it is oil in water type and if there is no significant increase in conductivity, it is water in oil type.

[B] Applications of emulsions :

- Disinfactants like phenyl, dettol when mixed with water form emulsion.
- Digestion of fat in small intestine occurs easily due to emulsion.
- In metallurgical process the concentration of ore by froath floatation method is based upon emulsion.
- Milk is an emulsion of liquid fat in water in which casein emulsifying agent.
- Cleansing action of soap is due to formation of emulsions. Soaps and detergents emulsify the grease along with the adhering dirt and carry them away in the wash water.
- For concentrating ores, the finely powdered ore is treated with an oil. Oil forms emulsion with the ore particles. When air is bubbled into the mixture, emulsion containing the particles of the mineral are carried to the surface.

Gelation :

- Ex: Gelatin dissolved in water forming a colloidal. Sol Which when cooled sets into a gelly. *Gel have honey-comb structure :
- **Ex :** Sillicic acid, Gum arabic, Sodium oleate, Gelatin, Solid alcohol, etc.

Types of Gel :

- (i) Elastic gel : Those gel which have elastic properties. Eg : Gelatin, Starch, Agar-Agar etc.
- (ii) Non- elastic gel : Those gel which are rigid. Eg : Silica gel.

Properties of Gel :

- 1. Syneresis/weeping of gel: The spontaneous liberation of liquid from a gel is called syneresis or weeping of gels. It is reverse of swelling. Eq : Gelatin, Agar-Agar show syneresis at low concentration while sillicic acid shows it at high concentration. 2. Imbibition or swelling of gel: When gel is kept in a suitable liquid (water) it absorb large volume of liquid. The phenomenon is called, imbibition or swelling of gel. 3. Thixotropic : Some gels when shaken to form a sol, on keeping changes into gel are termed as thixotropic gel and phenomenon is called thixotropy. Eg: Gelatin and silica liquify on shaking changing into corresponding sol and the sol on keeping changes back into gel. MISCELLANEOUS SOLVED PROBLEMS (MSPs) Which of the following is (are) lyophobic colloids ? 1. (1) Gold sol (2) As₂S₃ sol (3) Starch sol (4) Fe(OH)₃ sol
- **Sol.** Gold sol, As_2S_3 and $Fe(OH)_3$ are lyophobic colloid. Therefore, (1, 2, 4) are correct options.
- The presence of colloidal particles of dust in air imparts blue colour to the sky. This is due to

 (1) Absorption of the light
 (2) Scattering of the light
 (3) Reflection of the light
 (4) None of these
- **Sol.** Due to scattering of the light. Therefore, (2) is correct option.
- 3. The volume of nitrogen gas U_m (measured at STP) required to cover a sample of silica gel with a monomolecular layer is 129 cm³ g⁻¹ of gel. Calculate the surface area per gram of the gel if each nitrogen molecule occupies 16.2 x 10⁻²⁰ m².
- Ans. 561.8 cm³
- **Sol.** 22400 cm³ of N₂ at STP contain = 6.022×10^{23} molecules
 - \therefore 129 cm³ of N₂ at STP will contain =
 - Area occupied by a single molecule = $16.2 \times 10^{-20} \text{ m}^2$
 - : Area occupied by 3.468 × 10²¹ molecules of nitrogen = $(16.2 \times 10^{-20}) \times (3.468 \times 10^{21}) \text{ m}^2 = 561.8 \text{ m}^2$.

 $= 3.468 \times 10^{21}$ molecules

- 4. Which of the following has minimum gold number ?
 (1) Potato starch (2) Gum arabic (3) Gelatin (4) Albumin
 Colotin has minimum gold number. Therefore (2) is correct ention
- **Sol.** Gelatin has minimum gold number. Therefore, (3) is correct option.
- 5. Which of the following are correctly matched ?
 (1) Butter-gel
 (2) Milk-emulsion
 (3) Fog-aerosol
 (4) Dust-solid sol
- **Sol.** (1,2,3) are correct matches.
- 6. Explain the adsorption of nitrogen on iron.
- **Sol.** When nitrogen gas is brought in contact with iron at 83 K, it is physisorbed on iron surface as nitrogen molecules, N₂.As the temperature is increased the amount of nitrogen adsorbed decreases rapidly and at room temperature, practically there is no adsorption of nitrogen on iron. At 773 K and above, nitrogen is chemisorbed on the iron surface as nitrogen atoms.
- **7.** How do size of particles of adsorbent, pressure of gas and prevailing temperature influence the extent of adsorption of a gas on a solid ?
- **Sol.** (a) Smaller the size of the particles of the adsorbent, greater is the surface area and hence greater is the adsorption
 - (b) At constant temperature, adsorption first increases with increase of pressure and then attains equilibrium.

(c) In physical adsorption, it decreases with increase of temperature but in chemisorption, first it increases and then decreases.

8. How is adsorption of a gas is related to its critical temperature ?

- **Sol.** Higher is the critical temperature of a gas, greater the van der Waal's forces of attraction and hence greater is the adsorption.
- 9. Physical adsorption is essentially quite appreciable :
 - (1) at room temperature (2) at higher temperature
 - (3) at lower temperature (4) none of these
- **Sol.** Rate of physical adsorption decreases with increase in temperature (exothermic process). Therefore, (3) is correct option.
- 10. What type of colloidal sols are formed in the following ?
 (i) Through cooled water, vapours of sulphur are passed.
 (ii) White of an egg is mixed with water.
- Sol. (i) Sulphur molecules associate together to form molecular sols.
 (ii) Macromolecular sol because protein molecules present in the white of the egg are macromolecules & are soluble in water.
- **11.** What happens when persistent dialysis of a colloidal solution is carried out.

Sol. The stability of a colloidal sol is due to the presence of a small amount of the electrolyte. On persistent dialysis, the electrolyte is completely removed. As a result, the colloidal sol becomes unstable and gets coagulated.