

## Exercise-1

▶ Marked Questions may have for Revision Questions.

### PART - I : SUBJECTIVE QUESTIONS

- How much time (in years) would it take to distribute one Avogadro number of wheat grains if  $10^{10}$  grains are distributed each second ?
- The weight of one atom of Uranium is 238 amu. Its actual weight is ..... g.
- ▶ Calculate the weight of  $12.044 \times 10^{23}$  atoms of carbon.
- How many grams of silicon is present in 35 gram atoms of silicon (Given at. wt. of Si = 28).
- Find the total number of nucleons present in 12 g of  $^{12}\text{C}$  atoms.
- Find (i) the total number of neutrons, and (ii) the total mass of neutrons in 7 mg of  $^{14}\text{C}$ . (Assume that the mass of a neutron = mass of a hydrogen atom)
- ▶ Calculate the number of electrons, protons and neutrons in 1 mole of  $^{16}\text{O}^{2-}$  ions.
- How many atoms are there in 100 amu of He ?
- The density of liquid mercury is  $13.6 \text{ g/cm}^3$ . How many moles of mercury are there in 1 litre of the metal? (Atomic mass of Hg = 200.)
- Calculate the atomic mass (average) of chlorine using the following data :

	% Natural Abundance	Molar Mass
$^{35}\text{Cl}$	75	35.0 g
$^{37}\text{Cl}$	25	37.0 g

- ▶ Average atomic mass of Magnesium is 24.31 amu. This magnesium is composed of 79 mole % of  $^{24}\text{Mg}$  and remaining 21 mole % of  $^{25}\text{Mg}$  and  $^{26}\text{Mg}$ . Calculate mole % of  $^{26}\text{Mg}$ .
- ▶ The number of molecules in 16 g of methane is :
- Calculate the number of molecules in a drop of water weighing 0.09 g.
- A sample of ethane has the same mass as 10.0 million molecules of methane. How many  $\text{C}_2\text{H}_6$  molecules does the sample contain?
- The number of neutrons in 5 g of  $\text{D}_2\text{O}$  (D is  $^2\text{H}$ ) are :
- ▶ Calculate the weight of  $6.022 \times 10^{23}$  formula units of  $\text{CaCO}_3$ .
- From 200 mg of  $\text{CO}_2$ ,  $10^{21}$  molecules are removed. How many moles of  $\text{CO}_2$  are left ?

### PART - II : OBJECTIVE QUESTIONS

#### Single Correct Questions (SCQ)

- Which is not a basic postulate of Dalton's atomic theory ?  
 (1) Atoms are neither created nor destroyed in a chemical reaction.  
 (2) Different elements have different types of atoms.  
 (3) Atoms of an element may be different due to presence of isotopes.  
 (4) Each element is composed of extremely small particles called atoms.

2. The modern atomic weight scale is based on :  
 (1)  $^{12}\text{C}$  (2)  $^{16}\text{O}$  (3)  $^1\text{H}$  (4)  $^{18}\text{O}$
3. 1 amu is equal to  
 (1)  $\frac{1}{12}$  of C-12 (2)  $\frac{1}{14}$  of O-16 (3) 1 g of  $\text{H}_2$  (4)  $1.66 \times 10^{-23}$  kg
4. If the atomic mass of sodium is 23, the number of moles in 46 g of sodium is :  
 (1) 1 (2) 2 (3) 2.3 (4) 4.6
5. How many grams are contained in 1 gram-atom of Na ?  
 (1) 13 g (2) 23 g (3) 1 g (4)  $\frac{1}{23}$  g
6. 1.0 g of hydrogen contains  $6 \times 10^{23}$  atoms. The atomic weight of helium is 4. It follows that the number of atoms in 1 g of He is :  
 (1)  $\frac{1}{4} \times 6 \times 10^{23}$  (2)  $4 \times 6 \times 10^{23}$  (3)  $6 \times 10^{23}$  (4)  $12 \times 10^{23}$
7. The atomic weights of two elements A and B are 40u and 80u respectively. If x g of A contains y atoms, how many atoms are present in 2x g of B?  
 (1)  $\frac{y}{2}$  (2)  $\frac{y}{4}$  (3) y (4) 2y
8. A sample of aluminium has a mass of 54.0 g. What is the mass of the same number of magnesium atoms? (At. wt. Al = 27, Mg = 24)  
 (1) 12 g (2) 24 g (3) 48 g (4) 96 g.
9. The number of atoms in 558.5 g of Fe (at wt. = 55.85) is :  
 (1) Twice that in 60 g carbon (2)  $6.022 \times 10^{22}$   
 (3) Half in 8 g He (4)  $558.5 \times 6.023 \times 10^{23}$
10. Which of the following has the Maximum mass ?  
 (1) 1 g-atom of C (2)  $\frac{1}{2}$  mole of  $\text{CH}_4$   
 (3) 10 mL of water (4)  $3.011 \times 10^{23}$  atoms of oxygen
11. The total number of protons, electrons and neutrons in 12 g of  $^{12}_6\text{C}$  is :  
 (1)  $1.084 \times 10^{25}$  (2)  $6.022 \times 10^{23}$  (3)  $6.022 \times 10^{22}$  (4) 18
12. 1 mole of element X has mass, 3/10 times the mass of 1 mole of element Y. One average atom of element X has mass, 2 times the mass of one atom of  $^{12}\text{C}$ . What is the atomic weight of Y ?  
 (1) 80 (2) 15.77 (3) 46.67 (4) 40.0
13. The charge on 1 gram ions of  $\text{Al}^{3+}$  is : ( $N_A$  = Avogadro number, e = charge on one electron)  
 (1)  $\frac{1}{27} N_A e$  coulomb (2)  $\frac{1}{3} \times N_A e$  coulomb (3)  $\frac{1}{9} \times N_A e$  coulomb (4)  $3 \times N_A e$  coulomb
14. It is known that an atom contains protons, neutrons and electrons. If the mass of neutron is assumed to half of its original value whereas that of proton is assumed to be twice of its original value, then the atomic mass of  $^{14}_6\text{C}$  will be :  
 (1) same (2) 114.28 % less (3) 14.28 % more (4) 28.56 % less

15. The isotopic abundance of C-12 and C-14 is 98% and 2% by mass respectively. What would be the number of C-14 isotope in 12 g carbon sample ?  
 (1)  $1.032 \times 10^{22}$  (2)  $3.01 \times 10^{23}$  (3)  $5.88 \times 10^{23}$  (4)  $6.02 \times 10^{23}$
16. In chemical scale, the relative mass of the isotopic mixture of X atoms ( $X^{20}$ ,  $X^{21}$ ,  $X^{22}$ ) is approximately equal to : ( $X^{20}$  has 99 percent abundance)  
 (1) 20.002 (2) 21.00 (3) 22.00 (4) 20.00
17. Indium (atomic weight = 114.8) has two naturally occurring isotopes, the predominant one form has isotopic weight 115 and abundance of 95.00%. Which of the following isotopic weights is the most likely for the other isotope ?  
 (1) 111 (2) 112 (3) 113 (4) 114
18. The number of molecules of  $\text{CO}_2$  present in 44 g of  $\text{CO}_2$  is :  
 (1)  $6.0 \times 10^{23}$  (2)  $3 \times 10^{23}$  (3)  $12 \times 10^{23}$  (4)  $3 \times 10^{10}$
19. The number of mole of ammonia in 4.25 g of ammonia is :  
 (1) 0.425 (2) 0.25 (3) 0.236 (4) 0.2125
20. Which one of the following pairs of gases contains the same number of molecules :  
 (1) 16 g of  $\text{O}_2$  and 14 g of  $\text{N}_2$  (2) 8 g of  $\text{O}_2$  and 22 g of  $\text{CO}_2$   
 (3) 28 g of  $\text{N}_2$  and 22 g of  $\text{CO}_2$  (4) 32 g of  $\text{O}_2$  and 32 g of  $\text{N}_2$
21. The weight of a molecule of the compound  $\text{C}_{60}\text{H}_{22}$  is :  
 (1)  $1.09 \times 10^{-21}$  g (2)  $1.24 \times 10^{-21}$  g (3)  $5.025 \times 10^{-23}$  g (4)  $16.023 \times 10^{-23}$  g
22. Number of electrons in 1.8 mL of  $\text{H}_2\text{O}(\ell)$  is about :  
 (1)  $6.02 \times 10^{23}$  (2)  $3.011 \times 10^{23}$  (3)  $0.6022 \times 10^{21}$  (4)  $60.22 \times 10^{20}$
23. One mole of  $\text{P}_4$  molecules contain :  
 (1) 1 molecule (2) 4 molecules  
 (3)  $\frac{1}{4} \times 6.022 \times 10^{23}$  atoms (4)  $24.088 \times 10^{23}$  atoms
24. A sample of ammonium phosphate  $(\text{NH}_4)_3\text{PO}_4$  contains 3.18 mole of H atoms. The number of mole of O atoms in the sample is :  
 (1) 0.265 (2) 0.795 (3) 1.06 (4) 3.18
25. Torr is unit of :  
 (1) Temperature (2) Pressure (3) Volume (4) Density
26. The atmospheric pressure on Mars is 0.61 kPa. What is the pressure in mm Hg ?  
 (1) 0.63 (2) 4.6 (3) 6.3 (4) 3.2
27. Centigrade and Fahrenheit scales are related as :  
 (1)  $\frac{C}{5} = \frac{F - 32}{9}$  (2)  $\frac{C}{9} = \frac{F - 32}{5}$  (3)  $\frac{C}{8} = \frac{F - 32}{5}$  (4) None of these
28. At what temperature, both Celsius and Fahrenheit scale read the same value :  
 (1)  $100^\circ$  (2)  $130^\circ$  (3)  $60^\circ$  (4)  $-40^\circ$
29. The value of universal gas constant R depends on :  
 (1) temperature of gas (2) volume of gas  
 (3) number of moles of gas (4) units of volume and pressure
30. The value of gas constant in calorie per degree temperature per mol is approximately :

- (1) 1 cal                      (2) 2 cal                      (3) 3 cal                      (4) 4 cal
31. The value of R in SI unit is :  
 (1)  $8.314 \times 10^{-7} \text{ erg K}^{-1} \text{ mol}^{-1}$                       (2)  $8.314 \text{ JK}^{-1} \text{ mol}^{-1}$   
 (3)  $0.082 \text{ litre atm K}^{-1} \text{ mol}^{-1}$                       (4)  $2 \text{ cal K}^{-1} \text{ mol}^{-1}$
32. The pressure of sodium vapour in a 1.0 L container is 9.5 torr at  $927^\circ\text{C}$ . How many atoms are in the container ?  
 (1)  $9.7 \times 10^7$                       (2)  $7.5 \times 10^{19}$                       (3)  $4.2 \times 10^{17}$                       (4)  $9.7 \times 10^{19}$
33. The pressure of a gas having 2 mole in 44.8 litre vessel at 546 K is :  
 (1) 1 atm                      (2) 2 atm                      (3) 3 atm                      (4) 4 atm
34. According to the ideal gas laws, the molar volume of a gas is given by :  
 (1) 22.4 litre                      (2)  $RT / P$                       (3)  $8RT / PV$                       (4)  $RT / PV$
35. Equal volumes of oxygen gas and a second gas weigh 1.00 and  $19/8$  grams respectively under the same experimental conditions. Which of the following is the unknown gas?  
 (1) NO                      (2)  $\text{SO}_2$                       (3)  $\text{CS}_2$                       (4) CO
36. A high altitude balloon contains 6.0 g of helium in  $10^4 \text{ L}$  at 240 K. Assuming ideal gas behaviour, how many grams of helium would have to be added to increase the pressure to  $4.0 \times 10^{-3} \text{ atm}$  ?  
 (1) 1                      (2) 1.2                      (3) 1.5                      (4) 2.0
37. Four 1-litre flasks are separately filled with the gases  $\text{H}_2$ , He,  $\text{O}_2$  and  $\text{O}_3$  at the same temperature and pressure. The ratio of total number of atoms of these gases present in different flask would be :  
 (1) 1 : 1 : 1 : 1                      (2) 1 : 2 : 2 : 3                      (3) 2 : 1 : 2 : 3                      (4) 3 : 2 : 2 : 1
38. Under the same conditions, two gases have the same number of molecules. They must  
 (1) be noble gases                      (2) have equal volumes  
 (3) have a volume of  $22.4 \text{ dm}^3$  each                      (4) have an equal number of atoms
39. 16 g of an ideal gas  $\text{SO}_x$  occupies 5.6 L. at STP. The value of x is  
 (1)  $x = 3$                       (2)  $x = 2$                       (3)  $x = 4$                       (4) none of these
40. The ratio of the weight of one litre of a gas to the weight of 1.0 L oxygen gas both measured at S.T.P. is 2.22. The molecular weight of the gas would be :  
 (1) 14.002                      (2) 35.52                      (3) 71.04                      (4) 55.56
41. Avogadro number is :  
 (1) Number of atoms in one gram of the element  
 (2) Number of millilitre which one mole of a gaseous substance occupies at NTP (1 atm &  $0^\circ\text{C}$ )  
 (3) Number of molecules present in one gram molecular mass of a substance.  
 (4) All are correct
42. The weight of  $1 \times 10^{22}$  molecules of  $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$  is :  
 (1) 41.59 g                      (2) 415.9 g                      (3) 4.159 g                      (4) None of these
43. How many moles of electron weigh one kilogram :  
 (1)  $6.023 \times 10^{23}$                       (2)  $\frac{1}{9.108} \times 10^{31}$                       (3)  $\frac{6.023}{9.108} \times 10^{54}$                       (4)  $\frac{1}{9.108 \times 6.023} \times 10^8$
44. Number of atoms in 560 g of Fe (atomic mass  $56 \text{ gmol}^{-1}$ ) is :  
 (1) Twice that in 70 g N    (2) Half that in 20 g H    (3) Both (1) and (2)                      (4) None of these
45. Which has maximum number of atoms :

- (1) 24 g of C (12)      (2) 56 g of Fe (56)      (3) 27 g of Al (27)      (4) 108 g Ag (108)

46. If we consider that  $1/6$ , in place of  $1/12$  mass of carbon atom is taken to be the relative atomic mass unit, the mass of one mole of a substance will :  
 (1) decrease twice  
 (2) increase two fold  
 (3) remain unchanged  
 (4) be a function of the molecular mass of the substance
47. How many moles of magnesium phosphate,  $\text{Mg}_3(\text{PO}_4)_2$  will contain 0.25 mole of oxygen atoms ?  
 (1) 0.02      (2)  $3.125 \times 10^{-2}$       (3)  $1.25 \times 10^{-2}$       (4)  $2.5 \times 10^{-2}$
48. Given that the abundances of isotopes  $^{54}\text{Fe}$ ,  $^{56}\text{Fe}$  and  $^{57}\text{Fe}$  are 5%, 90% and 5% respectively, the atomic mass of Fe is :  
 (1) 55.85      (2) 55.95      (3) 55.75      (4) 56.05

### Comprehension #

A vessel of 25 L contains 20 g of ideal gas X at 300K. The pressure exerted by the gas is 1 atm. 20 g of ideal gas Y is added to the vessel keeping the same temperature. Total pressure became 3 atm. Upon further addition of 20 g ideal gas Z the pressure became 7 atm. Answer the following questions. (Hint: Ideal gas equation is applicable on mixture of ideal gases) [Take,  $R = 1/12 \text{ L.atm / mol K}$ ]

49. Find the molar mass of gas X.  
 (1) 20 g      (2) 10 g      (3) 30 g      (4) 5 g
50. Identify the correct statement(s) :  
 I. gas Y is lighter than gas X  
 II. gas Z is lighter than gas Y  
 (1) I only      (2) II only      (3) Both I and II      (4) None of the statements
51. Find the average molar mass of the mixture of gases X, Y and Z.  
 (1) 40/7      (2) 50/7      (3) 20      (4) 60/7
52. Match the column :

	Column-I				Column-II
	(Atomic mass (M))				(% composition of heavier isotope)
	Isotope-I	Isotope-II	Average		
(1)	$(z - 1)$	$(z + 3)$	$z$	(p)	25% by moles
(2)	$(z + 1)$	$(z + 3)$	$(z + 2)$	(q)	50% by moles
(3)	$z$	$3z$	$2z$	(r)	% by mass dependent on $z$
(4)	$(z - 1)$	$(z + 1)$	$z$	(s)	75% by mass



# Answers

## PART – I

1.  $1.9 \times 10^6$  years (approx.)
2.  $3.95 \times 10^{-22}$
3. 24 g
4. 980 g of Si
5.  $12 \times 6.022 \times 10^{23}$
6.  $24.088 \times 10^{20}$ , 0.004 g.
7.  $10 \times 6.022 \times 10^{23}$ ,  $8 \times 6.022 \times 10^{23}$ ,  $8 \times 6.022 \times 10^{23}$ .
8. 25
9. 68 mole
10. 35.5
11. 10
12.  $6.02 \times 10^{23}$
13.  $3.01 \times 10^{21}$  molecules of  $H_2O$
14.  $5.33 \times 10^6$
15.  $2.5 N_A$
16. 100 g
17. 0.00288

## PART – II

1. (3)
2. (1)
3. (1)
4. (2)
5. (2)
6. (1)
7. (3)
8. (3)
9. (1)
10. (1)
11. (1)
12. (1)
13. (4)
14. (3)
15. (1)
16. (1)
17. (1)
18. (1)
19. (2)
20. (1)
21. (2)
22. (1)
23. (4)
24. (3)
25. (2)
26. (2)
27. (1)
28. (4)
29. (4)
30. (2)
31. (2)
32. (2)
33. (2)
34. (2)
35. (3)
36. (4)
37. (3)
38. (2)
39. (2)
40. (3)
41. (3)
42. (3)
43. (4)
44. (3)
45. (1)
46. (3)
47. (2)
48. (2)
49. (1)
50. (3)
51. (4)
52. (1) - (p,r) ; (2) - (q,r) ; (3) - (q,s) ; (4) - (q,r)