

# Ordinary Thinking

## Objective Questions

### Significant figures, Units for measurement, Matter and Separation of mixture

- One fermi is [Haryana CEET 1994; DPMT 2004]
  - $10^{-13} \text{ cm}$
  - $10^{-15} \text{ cm}$
  - $10^{-10} \text{ cm}$
  - $10^{-12} \text{ cm}$
- A picometre is written as
  - $10^{-9} \text{ m}$
  - $10^{-10} \text{ m}$
  - $10^{-11} \text{ m}$
  - $10^{-12} \text{ m}$
- One atmosphere is equal to
  - 101.325 K pa
  - 1013.25 K pa
  - $10^5 \text{ Nm}$
  - None of these
- Dimensions of pressure are same as that of [CBSE PMT 1995]
  - Energy
  - Force
  - Energy per unit volume
  - Force per unit volume
- The prefix  $10^{18}$  is [Kerala MEE 2002]
  - Giga
  - Nano
  - Mega
  - Exa
- Given the numbers : 161cm, 0.161cm, 0.0161 cm. The number of significant figures for the three numbers are [CBSE PMT 1998]
  - 3, 4 and 5 respectively
  - 3, 3 and 3 respectively
  - 3, 3 and 4 respectively
  - 3, 4 and 4 respectively
- Significant figures in 0.00051 are
  - 5
  - 3
  - 2
  - 4
- Which of the following halogen can be purified by sublimation
  - $F_2$
  - $Cl_2$
  - $Br_2$
  - $I_2$
- Difference in density is the basis of [Kerala MEE 2002]
  - Ultrafiltration
  - Molecular sieving
  - Gravity Separation
  - Molecular attraction
- Which of the following elements of matter would best convey that there is life on earth
  - Oxygen
  - Hydrogen
  - Carbon
  - Iron
- The compound which is added to table salt for maintaining proper health is
  - KCl
  - KBr
  - NaI
  - $MgBr_2$
- Which of the following contains only one element
  - Marble
  - Diamond
  - Glass
  - Sand
- In known elements, the maximum number is of [CPMT 1985]
  - Metals
  - Non-metals
  - Metalloids
  - None of these
- Which one of the following is not an element
  - Diamond
  - Graphite
  - Silica
  - Ozone
- A mixture of  $ZnCl_2$  and  $PbCl_2$  can be separated by [AFMC 1989]
  - Distillation
  - Crystallization
  - Sublimation
  - Adding acetic acid
- A mixture of methyl alcohol and acetone can be separated by
  - Distillation
  - Fractional distillation
  - Steam distillation
  - Distillation under reduced pressure
- In the final answer of the expression  $\frac{(29.2 - 20.2)(1.79 \times 10^5)}{1.37}$ . The number of significant figures is [CBSE PMT 1994]
  - 1
  - 2
  - 3
  - 4
- 81.4 g sample of ethyl alcohol contains 0.002 g of water. The amount of pure ethyl alcohol to the proper number of significant figures is
  - 81.398 g
  - 71.40 g
  - 91.4 g
  - 81 g
- The unit  $J Pa^{-1}$  is equivalent to
  - $m^3$
  - $cm^3$
  - $dm^3$
  - None of these
- From the following masses, the one which is expressed nearest to the milligram is
  - 16 g
  - 16.4 g
  - 16.428 g
  - 16.4284 g
- The number of significant figures in  $6.02 \times 10^{23}$  is [Manipal PMT 2001]
  - 23
  - 3
  - 4
  - 26
- The prefix zepto stands for [DPMT 2004]
  - $10^9$
  - $10^{-12}$
  - $10^{-15}$
  - $10^{-21}$
- The significant figures in 3400 are [BHU 2004]
  - 2
  - 5
  - 6
  - 4
- The number of significant figures in 6.0023 are [Pb.CET 2001]
  - 5
  - 4
  - 3
  - 1
- Given  $P = 0.0030m$ ,  $Q = 2.40m$ ,  $R = 3000m$ , Significant figures in  $P, Q$  and  $R$  are respectively [Pb. CET 2002]
  - 2, 2, 1
  - 2, 3, 4
  - 4, 2, 1
  - 4, 2, 3

26. The number of significant figures in 60.0001 is [Pb. CET 2000]
- (a) 5 (b) 6  
(c) 3 (d) 2
27. A sample was weighted using two different balances. The result's were (i) 3.929 g (ii) 4.0 g. How would the weight of the sample be reported
- (a) 3.929 g (b) 3 g  
(c) 3.9 g (d) 3.93 g

### Laws of chemical combination

- Which of the following pairs of substances illustrate the law of multiple proportions [CPMT 1972, 78]
 

(a)  $CO$  and  $CO_2$  (b)  $H_2O$  and  $D_2O$   
(c)  $NaCl$  and  $NaBr$  (d)  $MgO$  and  $Mg(OH)_2$
- 1.0 g of an oxide of A contained 0.5 g of A. 4.0 g of another oxide of A contained 1.6 g of A. The data indicate the law of
 

(a) Reciprocal proportions (b) Constant proportions  
(c) Conservation of energy (d) Multiple proportions
- Among the following pairs of compounds, the one that illustrates the law of multiple proportions is
 

(a)  $NH_3$  and  $NCl_3$  (b)  $H_2S$  and  $SO_2$   
(c)  $CuO$  and  $Cu_2O$  (d)  $CS_2$  and  $FeSO_4$
- The percentage of copper and oxygen in samples of  $CuO$  obtained by different methods were found to be the same. This illustrates the law of [AMU 1982, 92]
 

(a) Constant proportions (b) Conservation of mass  
(c) Multiple proportions (d) Reciprocal proportions
- Two samples of lead oxide were separately reduced to metallic lead by heating in a current of hydrogen. The weight of lead from one oxide was half the weight of lead obtained from the other oxide. The data illustrates [AMU 1983]
 

(a) Law of reciprocal proportions  
(b) Law of constant proportions  
(c) Law of multiple proportions  
(d) Law of equivalent proportions
- Chemical equation is balanced according to the law of [AMU 1984]
 

(a) Multiple proportion (b) Reciprocal proportion  
(c) Conservation of mass (d) Definite proportions
- Avogadro number is
 

(a) Number of atoms in one gram of element  
(b) Number of millilitres which one mole of a gaseous substances occupies at NTP  
(c) Number of molecules present in one gram molecular mass of a substance  
(d) All of these
- Different proportions of oxygen in the various oxides of nitrogen prove the [MP PMT 1985]
 

(a) Equivalent proportion (b) Multiple proportion  
(c) Constant proportion (d) Conservation of matter
- Two elements X and Y have atomic weights of 14 and 16. They form a series of compounds A, B, C, D and E in which the same amount of element X, Y is present in the ratio 1 : 2 : 3 : 4 : 5. If the compound A has 28 parts by weight of X and 16 parts by weight of Y, then the compound of C will have 28 parts weight of X and [NCERT 1971]
 

(a) 32 parts by weight of Y (b) 48 parts by weight of Y  
(c) 64 parts by weight of Y (d) 80 parts by weight of Y
- Carbon and oxygen combine to form two oxides, carbon monoxide and carbon dioxide in which the ratio of the weights of carbon and oxygen is respectively 12 : 16 and 12 : 32. These figures illustrate the
 

(a) Law of multiple proportions  
(b) Law of reciprocal proportions  
(c) Law of conservation of mass  
(d) Law of constant proportions
- A sample of calcium carbonate ( $CaCO_3$ ) has the following percentage composition : Ca = 40%; C = 12%; O = 48%. If the law of constant proportions is true, then the weight of calcium in 4 g of a sample of calcium carbonate obtained from another source will be
 

(a) 0.016 g (b) 0.16 g  
(c) 1.6 g (d) 16 g
- n g of substance X reacts with m g of substance Y to form p g of substance R and q g of substance S. This reaction can be represented as,  $X + Y = R + S$ . The relation which can be established in the amounts of the reactants and the products will be
 

(a)  $n - m = p - q$  (b)  $n + m = p + q$   
(c)  $n = m$  (d)  $p = q$
- Which of the following is the best example of law of conservation of mass [NCERT 1975]
 

(a) 12 g of carbon combines with 32 g of oxygen to form 44 g of  $CO_2$   
(b) When 12 g of carbon is heated in a vacuum there is no change in mass  
(c) A sample of air increases in volume when heated at constant pressure but its mass remains unaltered  
(d) The weight of a piece of platinum is the same before and after heating in air
- The law of multiple proportions is illustrated by the two compounds [NCERT 1972]
 

(a) Sodium chloride and sodium bromide  
(b) Ordinary water and heavy water  
(c) Caustic soda and caustic potash  
(d) Sulphur dioxide and sulphur trioxide
- In compound A, 1.00 g nitrogen unites with 0.57 g oxygen. In compound B, 2.00 g nitrogen combines with 2.24 g oxygen. In compound C, 3.00 g nitrogen combines with 5.11 g oxygen. These results obey the following law [CPMT 1971]
 

(a) Law of constant proportion  
(b) Law of multiple proportion  
(c) Law of reciprocal proportion  
(d) Dalton's law of partial pressure

16. Hydrogen combines with oxygen to form  $H_2O$  in which 16 g of oxygen combine with 2 g of hydrogen. Hydrogen also combines with carbon to form  $CH_4$  in which 2 g of hydrogen combine with 6 g of carbon. If carbon and oxygen combine together then they will do show in the ratio of  
(a) 6 : 16 or 12 : 32 (b) 6 : 18  
(c) 1 : 2 (d) 12 : 24
17. 2 g of hydrogen combine with 16 g of oxygen to form water and with 6 g of carbon to form methane. In carbon dioxide 12 g of carbon are combined with 32 g of oxygen. These figures illustrate the law of  
(a) Multiple proportions (b) Constant proportions  
(c) Reciprocal proportions (d) Conservation of mass
18. An element forms two oxides containing respectively 53.33 and 36.36 percent of oxygen. These figures illustrate the law of  
(a) Conservation of mass (b) Constant proportions  
(c) Reciprocal proportions (d) Multiple proportions
19. After a chemical reaction, the total mass of reactants and products [MP PMT 1989]  
(a) Is always increased (b) Is always decreased  
(c) Is not changed (d) Is always less or more
20. A sample of pure carbon dioxide, irrespective of its source contains 27.27% carbon and 72.73% oxygen. The data support [AIIMS 1992]  
(a) Law of constant composition  
(b) Law of conservation of mass  
(c) Law of reciprocal proportions  
(d) Law of multiple proportions
21. The law of definite proportions is not applicable to nitrogen oxide because [EAMCET 1981]  
(a) Nitrogen atomic weight is not constant  
(b) Nitrogen molecular weight is variable  
(c) Nitrogen equivalent weight is variable  
(d) Oxygen atomic weight is variable
22. Which one of the following pairs of compounds illustrates the law of multiple proportion [EAMCET 1989]  
(a)  $H_2O, Na_2O$  (b)  $MgO, Na_2O$   
(c)  $Na_2O, BaO$  (d)  $SnCl_2, SnCl_4$
4. 1 amu is equal to  
(a)  $\frac{1}{12}$  of C - 12 (b)  $\frac{1}{14}$  of O - 16  
(c) 1g of  $H_2$  (d)  $1.66 \times 10^{-23}$  kg
5. Sulphur forms the chlorides  $S_2Cl_2$  and  $SCl_2$ . The equivalent mass of sulphur in  $SCl_2$  is [EAMCET 1985; Pb. CET 2001]  
(a) 8 g/mole (b) 16 g/mole  
(c) 64.8 g/mole (d) 32 g/mole
6. The sulphate of a metal M contains 9.87% of M. This sulphate is isomorphous with  $ZnSO_4 \cdot 7H_2O$ . The atomic weight of M is [IIT 1991]  
(a) 40.3 (b) 36.3  
(c) 24.3 (d) 11.3
7. When 100 ml of 1 M NaOH solution and 10 ml of 10 N  $H_2SO_4$  solution are mixed together, the resulting solution will be [DPMT 1982]  
(a) Alkaline (b) Acidic  
(c) Strongly acidic (d) Neutral
8. In chemical scale, the relative mass of the isotopic mixture of oxygen atoms ( $O^{16}, O^{17}, O^{18}$ ) is assumed to be equal to [Bihar MADT 1981]  
(a) 16.002 (b) 16.00  
(c) 17.00 (d) 11.00
9. For preparing 0.1 N solution of a compound from its impure sample of which the percentage purity is known, the weight of the substance required will be [MP PET 1996]  
(a) More than the theoretical weight  
(b) Less than the theoretical weight  
(c) Same as the theoretical weight  
(d) None of these
10. 1 mol of  $CH_4$  contains  
(a)  $6.02 \times 10^{23}$  atoms of H  
(b) 4 g atom of Hydrogen  
(c)  $1.81 \times 10^{23}$  molecules of  $CH_4$   
(d) 3.0 g of carbon
11. In the reaction  $2Na_2S_2O_3 + I_2 \rightarrow Na_2S_4O_6 + 2NaI$ , the equivalent weight of  $Na_2S_2O_3$  (mol. wt. = M) is equal to  
(a) M (b)  $M/2$   
(c)  $M/3$  (d)  $M/4$
12. When potassium permanganate is titrated against ferrous ammonium sulphate, the equivalent weight of potassium permanganate is [CPMT 1988]  
(a) Molecular weight / 10 (b) Molecular weight / 5  
(c) Molecular weight / 2 (d) Molecular weight
13. Boron has two stable isotopes,  $^{10}B$  (19%) and  $^{11}B$  (81%). The atomic mass that should appear for boron in the periodic table is [CBSE PMT 1990]  
(a) 10.8 (b) 10.2  
(c) 11.2 (d) 10.0

### Atomic, Molecular and Equivalent masses

1. Which property of an element is always a whole number [MP PMT 1986]  
(a) Atomic weight (b) Equivalent weight  
(c) Atomic number (d) Atomic volume
2. Which one of the following properties of an element is not variable [Bihar MADT 1981]  
(a) Valency (b) Atomic weight  
(c) Equivalent weight (d) All of these
3. The modern atomic weight scale is based on [MP PMT 2002]  
(a)  $C^{12}$  (b)  $O^{16}$   
(c)  $H^1$  (d)  $C^{13}$
4. 1 amu is equal to  
(a)  $\frac{1}{12}$  of C - 12 (b)  $\frac{1}{14}$  of O - 16  
(c) 1g of  $H_2$  (d)  $1.66 \times 10^{-23}$  kg
5. Sulphur forms the chlorides  $S_2Cl_2$  and  $SCl_2$ . The equivalent mass of sulphur in  $SCl_2$  is [EAMCET 1985; Pb. CET 2001]  
(a) 8 g/mole (b) 16 g/mole  
(c) 64.8 g/mole (d) 32 g/mole
6. The sulphate of a metal M contains 9.87% of M. This sulphate is isomorphous with  $ZnSO_4 \cdot 7H_2O$ . The atomic weight of M is [IIT 1991]  
(a) 40.3 (b) 36.3  
(c) 24.3 (d) 11.3
7. When 100 ml of 1 M NaOH solution and 10 ml of 10 N  $H_2SO_4$  solution are mixed together, the resulting solution will be [DPMT 1982]  
(a) Alkaline (b) Acidic  
(c) Strongly acidic (d) Neutral
8. In chemical scale, the relative mass of the isotopic mixture of oxygen atoms ( $O^{16}, O^{17}, O^{18}$ ) is assumed to be equal to [Bihar MADT 1981]  
(a) 16.002 (b) 16.00  
(c) 17.00 (d) 11.00
9. For preparing 0.1 N solution of a compound from its impure sample of which the percentage purity is known, the weight of the substance required will be [MP PET 1996]  
(a) More than the theoretical weight  
(b) Less than the theoretical weight  
(c) Same as the theoretical weight  
(d) None of these
10. 1 mol of  $CH_4$  contains  
(a)  $6.02 \times 10^{23}$  atoms of H  
(b) 4 g atom of Hydrogen  
(c)  $1.81 \times 10^{23}$  molecules of  $CH_4$   
(d) 3.0 g of carbon
11. In the reaction  $2Na_2S_2O_3 + I_2 \rightarrow Na_2S_4O_6 + 2NaI$ , the equivalent weight of  $Na_2S_2O_3$  (mol. wt. = M) is equal to  
(a) M (b)  $M/2$   
(c)  $M/3$  (d)  $M/4$
12. When potassium permanganate is titrated against ferrous ammonium sulphate, the equivalent weight of potassium permanganate is [CPMT 1988]  
(a) Molecular weight / 10 (b) Molecular weight / 5  
(c) Molecular weight / 2 (d) Molecular weight
13. Boron has two stable isotopes,  $^{10}B$  (19%) and  $^{11}B$  (81%). The atomic mass that should appear for boron in the periodic table is [CBSE PMT 1990]  
(a) 10.8 (b) 10.2  
(c) 11.2 (d) 10.0

14. What is the concentration of nitrate ions if equal volumes of 0.1  $M$   $AgNO_3$  and 0.1  $M$   $NaCl$  are mixed together  
[CPMT 1983; NCERT 1985]  
(a) 0.1  $M$  (b) 0.2  $M$   
(c) 0.05  $M$  (d) 0.25  $M$
15. Total number of atoms represented by the compound  $CuSO_4 \cdot 5H_2O$  is  
[BHU 2005]  
(a) 27 (b) 21  
(c) 5 (d) 8
16. 74.5 g of a metallic chloride contain 35.5 g of chlorine. The equivalent weight of the metal is  
[CPMT 1986]  
(a) 19.5 (b) 35.5  
(c) 39.0 (d) 78.0
17. 7.5 grams of a gas occupy 5.8 litres of volume at STP the gas is  
(a)  $NO$  (b)  $N_2O$   
(c)  $CO$  (d)  $CO_2$
18. The number of atoms in 4.25 g of  $NH_3$  is approximately  
[CBSE PMT 1999; MH CET 2003]  
(a)  $1 \times 10^{23}$  (b)  $2 \times 10^{23}$   
(c)  $4 \times 10^{23}$  (d)  $6 \times 10^{23}$
19. One litre of a gas at STP weight 1.16 g it can possible be  
[AMU 1992]  
(a)  $C_2H_2$  (b)  $CO$   
(c)  $O_2$  (d)  $CH_4$
20. The vapour density of a gas is 11.2. The volume occupied by 11.2 g of the gas at ATP will be  
[Bihar CET 1995]  
(a) 11.2 L (b) 22.4 L  
(c) 1 L (d) 44.8 L
21. Equivalent weight of crystalline oxalic acid is  
[MP PMT 1995]  
(a) 30 (b) 63  
(c) 53 (d) 45
22. The equivalent weight of an element is 4. Its chloride has a V.D 59.25. Then the valency of the element is  
[BHU 1997]  
(a) 4 (b) 3  
(c) 2 (d) 1
23. 1.25 g of a solid dibasic acid is completely neutralised by 25 ml of 0.25 molar  $Ba(OH)_2$  solution. Molecular mass of the acid is  
(a) 100 (b) 150  
(c) 120 (d) 200
24. The oxide of a metal has 32% oxygen. Its equivalent weight would be  
[MP PMT 1985]  
(a) 34 (b) 32  
(c) 17 (d) 8
25. The mass of a molecule of water is  
[Bihar CEE 1995]  
(a)  $3 \times 10^{-26}$  kg (b)  $3 \times 10^{-25}$  kg  
(c)  $1.5 \times 10^{-26}$  kg (d)  $2.5 \times 10^{-26}$  kg
26. 1.24 gm P is present in 2.2 gm  
(a)  $P_4S_3$  (b)  $P_2S_2$   
(c)  $PS_2$  (d)  $P_2S_4$
27. The atomic weights of two elements A and B are 40 and 80 respectively. If x g of A contains y atoms, how many atoms are present in 2x g of B  
(a)  $\frac{y}{2}$  (b)  $\frac{y}{4}$   
(c) y (d) 2y
28. Assuming fully decomposed, the volume of  $CO_2$  released at STP on heating 9.85 g of  $BaCO_3$  (Atomic mass of Ba=137) will be  
[CBSE PMT 2000]  
(a) 0.84 L (b) 2.24 L  
(c) 4.06 L (d) 1.12 L
29. If  $N_A$  is Avogadro's number then number of valence electrons in 4.2 g of nitride ions ( $N^{3-}$ )  
(a)  $2.4 N_A$  (b)  $4.2 N_A$   
(c)  $1.6 N_A$  (d)  $3.2 N_A$
30. The weight of  $1 \times 10^{22}$  molecules of  $CuSO_4 \cdot 5H_2O$  is  
[IIT 1991]  
(a) 41.59 g (b) 415.9 g  
(c) 4.159 g (d) None of these
31. Rearrange the following (I to IV) in the order of increasing masses and choose the correct answer from (a), (b), (c) and (d) (Atomic mass: N=14, O=16, Cu=63).  
I. 1 molecule of oxygen  
II. 1 atom of nitrogen  
III.  $1 \times 10^{-10}$  g molecular weight of oxygen  
IV.  $1 \times 10^{-10}$  g atomic weight of copper  
(a) II<I<III<IV (b) IV<III<II<I  
(c) II<III<I<IV (d) III<IV<I<II
32. 1.520 g of the hydroxide of a metal on ignition gave 0.995 gm of oxide. The equivalent weight of metal is  
[DPMT 1984]  
(a) 1.520 (b) 0.995  
(c) 19.00 (d) 9.00
33. How much coulomb charge is present on 1g ion of  $N^{3-}$   
(a)  $5.2 \times 10^6$  Coulomb (b)  $2.894 \times 10^5$  Coulomb  
(c)  $6.6 \times 10^6$  Coulomb (d)  $8.2 \times 10^6$  Coulomb
34. Ratio of  $C_p$  and  $C_v$  of a gas X is 1.4, the number of atom of the gas 'X' present in 11.2 litres of it at NTP will be  
[CBSE 1999]  
(a)  $6.02 \times 10^{23}$  (b)  $1.2 \times 10^{23}$   
(c)  $3.01 \times 10^{23}$  (d)  $2.01 \times 10^{23}$
35. If we consider that 1/6, in place of 1/12, mass of carbon atom is taken to be the relative atomic mass unit, the mass of one mole of a substance will  
[AIEEE 2005]  
(a) Decrease twice  
(b) Increase two fold  
(c) Remain unchanged  
(d) Be a function of the molecular mass of the substance

36. What should be the equivalent weight of phosphorous acid, if  $P=31$ ;  $O=16$ ;  $H=1$   
(a) 82 (b) 41  
(c) 20.5 (d) None of these
37. The number of molecule at NTP in 1 ml of an ideal gas will be  
(a)  $6 \times 10^{23}$  (b)  $2.69 \times 10^{19}$   
(c)  $2.69 \times 10^{23}$  (d) None of these
38. The specific heat of a metal is 0.16 its approximate atomic weight would be  
(a) 32 (b) 16  
(c) 40 (d) 64
39. The weight of a molecule of the compound  $C_{60}H_{122}$  is  
[AIIMS 2000]  
(a)  $1.4 \times 10^{-21}$  g (b)  $1.09 \times 10^{-21}$  g  
(c)  $5.025 \times 10^{23}$  g (d)  $16.023 \times 10^{23}$  g
40. What is the weight of oxygen required for the complete combustion of 2.8 kg of ethylene [CBSE PMT 1989]  
(a) 2.8 kg (b) 6.4 kg  
(c) 9.6 kg (d) 96 kg
41. What volume of  $NH_3$  gas at STP would be needed to prepare 100ml of 2.5 molal (2.5m) ammonium hydroxide solution  
(a) 0.056 litres (b) 0.56 litres  
(c) 5.6 litres (d) 11.2 litres
42. If the density of water is  $1 \text{ g cm}^{-3}$  then the volume occupied by one molecule of water is approximately [Pb. PMT 2004]  
(a)  $18 \text{ cm}^3$  (b)  $22400 \text{ cm}^3$   
(c)  $6.02 \times 10^{-23} \text{ cm}^3$  (d)  $3.0 \times 10^{-23} \text{ cm}^3$
43. Caffeine has a molecular weight of 194. If it contains 28.9% by mass of nitrogen, number of atoms of nitrogen in one molecule of caffeine is  
(a) 4 (b) 6  
(c) 2 (d) 3
44. A 400 mg iron capsule contains 100 mg of ferrous fumarate,  $(CHCOO)_2Fe$ . The percentage of iron present in it is approximately  
(a) 33% (b) 25%  
(c) 14% (d) 8%
45. The element whose a atom has mass of  $10.86 \times 10^{-26} \text{ kg}$  is  
(a) Boron (b) Calcium  
(c) Silver (d) Zinc
46. The number of gram atoms of oxygen present in 0.3 gram mole of  $(COOH)_2 \cdot 2H_2O$  is  
(a) 0.6 (b) 1.8  
(c) 1.2 (d) 3.6
47. A gaseous mixture contains  $CH_4$  and  $C_2H_6$  in equimolecular proportion. The weight of 2.24 litres of this mixture at NTP is  
(a) 4.6 g (b) 1.6 g  
(c) 2.3 g (d) 23 g
48. Vapour density of a metal chloride is 66. Its oxide contains 53% metal. The atomic weight of the metal is  
[Bihar MADT 1982]  
(a) 21 (b) 54  
(c) 27.06 (d) 2.086
49. One gram of hydrogen is found to combine with 80g of bromine one gram of calcium valency=2 combines with 4g of bromine the equivalent weight of calcium is  
(a) 10 (b) 20  
(c) 40 (d) 80
50. The equivalent weight of  $MnSO_4$  is half its molecular weight when it is converted to [IIT 1988; CPMT 1994]  
(a)  $Mn_2O_3$  (b)  $MnO_2$   
(c)  $MnO_4$  (d)  $MnO_4^{2-}$
51. 100 mL of  $PH_3$  on decomposition produced phosphorus and hydrogen. The change in volume is [MNR 1986]  
(a) 50 mL increase (b) 500 mL decrease  
(c) 900 mL decrease (d) Nil.
52. 12g of Mg (at. mass 24) on reacting completely with acid gives hydrogen gas, the volume of which at STP would be [CPMT 1978]  
(a) 22.4 L (b) 11.2 L  
(c) 44.8 L (d) 6.1 L
53. Which of the following has least mass [Pb. PET 1985]  
(a) 2 g atom of nitrogen (b)  $3 \times 10^{23}$  atoms of C  
(c) 1 mole of S (d) 7.0 g of Ag
54. How many mole of helium gas occupy 22.4 L at  $0^\circ C$  at 1 atm. pressure [Kurukshetra CEE 1992; CET 1992]  
(a) 0.11 (b) 0.90  
(c) 1.0 (d) 1.11
55. Volume of a gas at STP is  $1.12 \times 10^{-7}$  cc. Calculate the number of molecules in it [BHU 1997]  
(a)  $3.01 \times 10^{20}$  (b)  $3.01 \times 10^{12}$   
(c)  $3.01 \times 10^{23}$  (d)  $3.01 \times 10^{24}$
56. 4.4 g of an unknown gas occupies 2.24L of volume at standard temperature and pressure. The gas may be [MP PMT 1995]  
(a) Carbon dioxide (b) Carbon monoxide  
(c) Oxygen (d) Sulphur dioxide
57. The number of moles of oxygen in 1 L of air containing 21% oxygen by volume, in standard conditions, is [CBSE PMT 1995; Pb. PMT 2004]  
(a) 0.186 mol (b) 0.21 mol  
(c) 2.10 mol (d) 0.0093 mol
58. The number of molecules in 8.96 L of a gas at  $0^\circ C$  and 1 atmosphere pressure is approximately [BHU 1993]  
(a)  $6.02 \times 10^{23}$  (b)  $12.04 \times 10^{23}$   
(c)  $18.06 \times 10^{23}$  (d)  $24.08 \times 10^{22}$

59. The equivalent weight of a metal is 9 and vapour density of its chloride is 59.25. The atomic weight of metal is [Pb. CET 2002]  
 (a) 23.9 (b) 27.3  
 (c) 36.3 (d) 48.3
60. The molecular weight of a gas is 45. Its density at STP is [Pb. PMT 2004]  
 (a) 22.4 (b) 11.2  
 (c) 5.7 (d) 2.0
61. Equivalent weight of a bivalent metal is 37.2. The molecular weight of its chloride is [MH CET 2003]  
 (a) 412.2 (b) 216  
 (c) 145.4 (d) 108.2
62. On reduction with hydrogen, 3.6 g of an oxide of metal left 3.2 g of metal. If the vapour density of metal is 32, the simplest formula of the oxide would be [DPMT 2004]  
 (a)  $MO$  (b)  $M_2O_3$   
 (c)  $M_2O$  (d)  $M_2O_5$
63. The number of molecules in 4.25 g of ammonia are [Pb. CET 2000]  
 (a)  $0.5 \times 10^{23}$  (b)  $1.5 \times 10^{23}$   
 (c)  $3.5 \times 10^{23}$  (d)  $1.8 \times 10^{32}$
- (c)  $6 \times 10^{23}$  (d)  $12 \times 10^{23}$
6. The volume occupied by 4.4 g of  $CO_2$  at STP is [AFMC 1997, 2004; Pb. CET 1997, 2002]  
 (a) 22.4 L (b) 2.24 L  
 (c) 0.224 L (d) 0.1 L
7. The number of water molecules present in a drop of water (volume 0.0018 ml) at room temperature is [DCE 2000]  
 (a)  $6.023 \times 10^{19}$  (b)  $1.084 \times 10^{18}$   
 (c)  $4.84 \times 10^{17}$  (d)  $6.023 \times 10^{23}$
8. One mole of calcium phosphide on reaction with excess of water gives [IIT 1999]  
 (a) One mole of phosphine  
 (b) Two moles of phosphoric acid  
 (c) Two moles of phosphine  
 (d) One mole of phosphorus pentoxide
9. 19.7 kg of gold was recovered from a smuggler. How many atoms of gold were recovered ( $Au = 197$ ) [Pb. CET 1985]  
 (a) 100 (b)  $6.02 \times 10^{23}$   
 (c)  $6.02 \times 10^{24}$  (d)  $6.02 \times 10^{25}$
10. The total number of protons in 10 g of calcium carbonate is ( $N_0 = 6.023 \times 10^{23}$ )  
 (a)  $1.5057 \times 10^{24}$  (b)  $2.0478 \times 10^{24}$   
 (c)  $3.0115 \times 10^{24}$  (d)  $4.0956 \times 10^{24}$
11. The number of molecules in 16 g of methane is  
 (a)  $3.0 \times 10^{23}$  (b)  $6.02 \times 10^{23}$   
 (c)  $\frac{16}{6.02} \times 10^{23}$  (d)  $\frac{16}{3.0} \times 10^{23}$
12. Number of molecules in 100 ml of each of  $O_2$ ,  $NH_3$  and  $CO_2$  at STP are [Bihar MADT 1985]  
 (a) In the order  $CO_2 < O_2 < NH_3$   
 (b) In the order  $NH_3 < O_2 < CO_2$   
 (c) The same  
 (d)  $NH_3 = CO_2 < O_2$
13. The molecular weight of hydrogen peroxide is 34. What is the unit of molecular weight [MP PMT 1986]  
 (a) g (b) mol  
 (c)  $g \text{ mol}^{-1}$  (d)  $\text{mol g}^{-1}$
14. The number of water molecules in 1 litre of water is [EAMCET 1990]  
 (a) 18 (b)  $18 \times 1000$   
 (c)  $N_A$  (d)  $55.55 N_A$
15. The number of electrons in a mole of hydrogen molecule is [CPMT 1987]  
 (a)  $6.02 \times 10^{23}$  (b)  $12.046 \times 10^{23}$   
 (c)  $3.0115 \times 10^{23}$  (d) Indefinite
16. The numbers of moles of  $BaCO_3$  which contain 1.5 moles of oxygen atoms is [EAMCET 1991]  
 (a) 0.5 (b) 1  
 (c) 3 (d)  $6.02 \times 10^{23}$

### The mole concept

1. Which one of the following pairs of gases contains the same number of molecules [EAMCET 1987]  
 (a) 16 g of  $O_2$  and 14 g of  $N_2$   
 (b) 8 g of  $O_2$  and 22 g of  $CO_2$   
 (c) 28 g of  $N_2$  and 22 g of  $CO_2$   
 (d) 32 g of  $O_2$  and 32 g of  $N_2$
2. Number of gm of oxygen in 32.2 g  $Na_2SO_4 \cdot 10H_2O$  is [Haryana PMT 2000]  
 (a) 20.8 (b) 22.4  
 (c) 2.24 (d) 2.08
3. 250 ml of a sodium carbonate solution contains 2.65 grams of  $Na_2CO_3$ . If 10 ml of this solution is diluted to one litre, what is the concentration of the resultant solution (mol. wt. of  $Na_2CO_3 = 106$ ) [EAMCET 2001]  
 (a) 0.1 M (b) 0.001 M  
 (c) 0.01 M (d)  $10^{-4}$  M
4. A molar solution is one that contains one mole of a solute in [IIT 1986]  
 (a) 1000 g of the solvent (b) One litre of the solvent  
 (c) One litre of the solution (d) 22.4 litres of the solution
5. The number of oxygen atoms in 4.4 g of  $CO_2$  is approx. [CBSE PMT 1990]  
 (a)  $1.2 \times 10^{23}$  (b)  $6 \times 10^{22}$

17. Which of the following is Loschmidt number  
(a)  $6 \times 10^{23}$  (b)  $2.69 \times 10^{19}$   
(c)  $3 \times 10^{23}$  (d) None of these
18. How many molecules are present in one gram of hydrogen  
[AIIMS 1982]  
(a)  $6.02 \times 10^{23}$  (b)  $3.01 \times 10^{23}$   
(c)  $2.5 \times 10^{23}$  (d)  $1.5 \times 10^{23}$
19. The total number of gm-molecules of  $SO_2Cl_2$  in 13.5 g of sulphuryl chloride is  
[CPMT 1992]  
(a) 0.1 (b) 0.2  
(c) 0.3 (d) 0.4
20. The largest number of molecules is in  
[BHU 1997]  
(a) 34 g of water (b) 28 g of  $CO_2$   
(c) 46 g of  $CH_3OH$  (d) 54 g of  $N_2O_5$
21. The number of moles of sodium oxide in 620 g of it is  
[BHU 1992]  
(a) 1 mol (b) 10 moles  
(c) 18 moles (d) 100 moles
22. 2 g of oxygen contains number of atoms equal to that in  
[BHU 1992]  
(a) 0.5 g of hydrogen (b) 4 g of sulphur  
(c) 7 g of nitrogen (d) 2.3 g of sodium
23. Molarity of liquid  $HCl$  with density equal to  $1.17 \text{ g/cc}$  is  
[CBSE PMT 2001]  
(a) 36.5 (b) 18.25  
(c) 32.05 (d) 4.65
24. How many atoms are contained in one mole of sucrose ( $C_{12}H_{22}O_{11}$ )  
[Pb. PMT 2002]  
(a)  $45 \times 6.02 \times 10^{23}$  atoms/mole  
(b)  $5 \times 6.62 \times 10^{23}$  atoms/mole  
(c)  $5 \times 6.02 \times 10^{23}$  atoms/mole  
(d) None of these
25. The number of molecules of  $CO_2$  present in 44 g of  $CO_2$  is  
[BCECE 2005]  
(a)  $6.0 \times 10^{23}$  (b)  $3 \times 10^{23}$   
(c)  $12 \times 10^{23}$  (d)  $3 \times 10^{10}$
26. A sample of phosphorus trichloride ( $PCl_3$ ) contains 1.4 moles of the substance. How many atoms are there in the sample [Kerala PMT 2004]  
(a) 4 (b) 5.6  
(c)  $8.431 \times 10^{23}$  (d)  $3.372 \times 10^{24}$   
(e)  $2.409 \times 10^{24}$
27. The number of sodium atoms in 2 moles of sodium ferrocyanide is  
[BHU 2004]  
(a)  $12 \times 10^{23}$  (b)  $26 \times 10^{23}$   
(c)  $34 \times 10^{23}$  (d)  $48 \times 10^{23}$
- (a) 40 (b) 60  
(c) 8 (d) 10
2. The percentage of nitrogen in urea is about [KCET 2001]  
(a) 46 (b) 85  
(c) 18 (d) 28
3. If two compounds have the same empirical formula but different molecular formula, they must have [MP PMT 1986]  
(a) Different percentage composition  
(b) Different molecular weights  
(c) Same viscosity  
(d) Same vapour density
4. A compound (80 g) on analysis gave  $C = 24 \text{ g}$ ,  $H = 4 \text{ g}$ ,  $O = 32 \text{ g}$ . Its empirical formula is [CPMT 1981]  
(a)  $C_2H_2O_2$  (b)  $C_2H_2O$   
(c)  $CH_2O_2$  (d)  $CH_2O$
5. The empirical formula of a compound is  $CH_2O$ . 0.0835 moles of the compound contains 1.0 g of hydrogen. Molecular formula of the compound is  
(a)  $C_2H_{12}O_6$  (b)  $C_5H_{10}O_5$   
(c)  $C_4H_8O_8$  (d)  $C_3H_6O_3$
6. The empirical formula of an acid is  $CH_2O_2$ , the probable molecular formula of acid may be [AFMC 2000]  
(a)  $CH_2O$  (b)  $CH_2O_2$   
(c)  $C_2H_4O_2$  (d)  $C_3H_6O_4$
7. In which of the following pairs of compounds the ratio of C, H and O is same  
(a) Acetic acid and methyl alcohol  
(b) Glucose and acetic acid  
(c) Fructose and sucrose  
(d) All of these

### Chemical stoichiometry

1. How much of  $NaOH$  is required to neutralise  $1500 \text{ cm}^3$  of 0.1 N  $HCl$  ( $Na = 23$ ) [KCET 2001]  
(a) 40 g (b) 4 g  
(c) 6 g (d) 60 g
2. How much water should be added to 200 c.c of semi normal solution of  $NaOH$  to make it exactly deci normal [AFMC 1983]  
(a) 200 cc (b) 400 cc  
(c) 800 cc (d) 600 cc
3. 2.76 g of silver carbonate on being strongly heated yield a residue weighing [Pb. CET 2003]  
(a) 2.16 g (b) 2.48 g  
(c) 2.64 g (d) 2.32 g

### Percentage composition & Molecular formula

1. The percentage of oxygen in  $NaOH$  is [CPMT 1979]

4. In the reaction,  $4\text{NH}_3(\text{g}) + 5\text{O}_2(\text{g}) \rightarrow 4\text{NO}(\text{g}) + 6\text{H}_2\text{O}(\text{g})$ , When 1 mole of ammonia and 1 mole of  $\text{O}_2$  are made to react to completion
- 1.0 mole of  $\text{H}_2\text{O}$  is produced
  - 1.0 mole of  $\text{NO}$  will be produced
  - All the oxygen will be consumed
  - All the ammonia will be consumed
5. Haemoglobin contains 0.33% of iron by weight. The molecular weight of haemoglobin is approximately 67200. The number of iron atoms (At. wt. of  $\text{Fe} = 56$ ) present in one molecule of haemoglobin is [CBSE PMT 1998]
- 6
  - 1
  - 4
  - 2
6. What quantity of ammonium sulphate is necessary for the production of  $\text{NH}_3$  gas sufficient to neutralize a solution containing 292 g of  $\text{HCl}$ ? [ $\text{HCl} = 36.5$ ;  $(\text{NH}_4)_2\text{SO}_4 = 132$ ;  $\text{NH}_3 = 17$ ] [CPMT 1992]
- 272 g
  - 403 g
  - 528 g
  - 1056 g
7. The percentage of  $\text{P}_2\text{O}_5$  in diammonium hydrogen phosphate  $(\text{NH}_4)_2\text{HPO}_4$  is [CPMT 1992]
- 23.48
  - 46.96
  - 53.78
  - 71.00
8. If  $1\frac{1}{2}$  moles of oxygen combine with  $\text{Al}$  to form  $\text{Al}_2\text{O}_3$  the weight of  $\text{Al}$  used in the reaction is ( $\text{Al} = 27$ ) [EAMCET 1980]
- 27 g
  - 54 g
  - 49.5 g
  - 31 g
9. The percentage of  $\text{Se}$  in peroxidase anhydrous enzyme is 0.5% by weight (atomic weight = 78.4). Then minimum molecular weight of peroxidase anhydrous enzyme is [CBSE PMT 2001]
- $1.568 \times 10^4$
  - $1.568 \times 10^3$
  - 15.68
  - $3.136 \times 10^4$
10.  $\text{H}_2$  evolved at STP on complete reaction of 27 g of Aluminium with excess of aqueous  $\text{NaOH}$  would be [CPMT 1991]
- 22.4
  - 44.8
  - 67.2
  - 33.6 litres
11. What is the % of  $\text{H}_2\text{O}$  in  $\text{Fe}(\text{CNS})_3 \cdot 3\text{H}_2\text{O}$
- 45
  - 30
  - 19
  - 25
12. What weight of  $\text{SO}_2$  can be made by burning sulphur in 5.0 moles of oxygen
- 640 grams
  - 160 grams
  - 80 grams
  - 320 grams
13. What is the normality of a 1 M solution of  $\text{H}_3\text{PO}_4$  [AIIMS 1982]
- 0.5 N
  - 1.0 N
  - 2.0 N
  - 3.0 N
14. Normality of 2M sulphuric acid is [AIIMS 1992]
- 2N
  - 4N
  - $\frac{N}{2}$
  - $\frac{N}{4}$
15. How many g of a dibasic acid (Mol. wt. = 200) should be present in 100 ml of its aqueous solution to give decinormal strength [AIIMS 1992]
- 1 g
  - 2 g
  - 10 g
  - 20 g
16. The solution of sulphuric acid contains 80% by weight  $\text{H}_2\text{SO}_4$ . Specific gravity of this solution is 1.71. Its normality is about [CBSE 1991]
- 18.0
  - 27.9
  - 1.0
  - 10.0
17. Mohr's salt is dissolved in dil.  $\text{H}_2\text{SO}_4$  instead of distilled water to
- Enhance the rate of dissolution
  - Prevent cationic hydrolysis
  - Increase the rate of ionisation
  - Increase its reducing strength
18. Acidified potassium permanganate solution is decolourised by
- Bleaching powder
  - White vitriol
  - Mohr's salt
  - Microcosmic salt
19. Approximate atomic weight of an element is 26.89. If its equivalent weight is 8.9, the exact atomic weight of element would be [DPMT 1984]
- 26.89
  - 8.9
  - 17.8
  - 26.7
20. Vapour density of a gas is 22. What is its molecular mass [AFMC 2000]
- 33
  - 22
  - 44
  - 11
21. Equivalent weight of  $\text{KMnO}_4$  acting as an oxidant in acidic medium is [CPMT 1990; MP PET 1999]
- The same as its molecular weight
  - Half of its molecular weight
  - One-third of its molecular weight
  - One-fifth of its molecular weight
22. 0.16 g of dibasic acid required 25 ml of decinormal  $\text{NaOH}$  solution for complete neutralisation. The molecular weight of the acid will be [CPMT 1989]
- 32
  - 64
  - 128
  - 256



23. To neutralise 20 ml of  $M/10$  sodium hydroxide, the volume of  $M/20$  hydrochloric acid required is [CPMT 1992]  
[Andhra MBBS 1980]  
(a) 10 ml (b) 15 ml  
(c) 20 ml (d) 40 ml
24. Hydrochloric acid solutions A and B have concentration of 0.5 N and 0.1 N respectively. The volume of solutions A and B required to make 2 litres of 0.2 N hydrochloric acid [KCET 1993]  
(a) 0.5 l of A + 1.5 l of B  
(b) 1.5 l of A + 0.5 l of B  
(c) 1.0 l of A + 1.0 l of B  
(d) 0.75 l of A + 1.25 l of B
25. 5 ml of N HCl, 20 ml of  $N/2$   $H_2SO_4$  and 30 ml of  $N/3$   $HNO_3$  are mixed together and volume made to one litre. The normality of the resulting solution is [MNR 1991]  
(a)  $N/5$  (b)  $N/10$   
(c)  $N/20$  (d)  $N/40$
26. Under similar conditions of pressure and temperature, 40 ml of slightly moist hydrogen chloride gas is mixed with 20 ml of ammonia gas, the final volume of gas at the same temperature and pressure will be [CBSE PMT 1993]  
(a) 100 ml (b) 20 ml  
(c) 40 ml (d) 60 ml
27.  $KMnO_4$  reacts with oxalic acid according to the equation,  $2MnO_4^- + 5C_2O_4^{2-} + 16H^+ \rightarrow 2Mn^{2+} + 10CO_2 + 8H_2O$ , here 20 ml of 0.1 M  $KMnO_4$  is equivalent to [CBSE PMT 1996]  
(a) 20 ml of 0.5 M  $H_2C_2O_4$  (b) 50 ml of 0.1 M  $H_2C_2O_4$   
(c) 50 ml of 0.5 M  $H_2C_2O_4$  (d) 20 ml of 0.1 M  $H_2C_2O_4$
28. In order to prepare one litre normal solution of  $KMnO_4$ , how many grams of  $KMnO_4$  are required if the solution is used in acidic medium for oxidation [MP PET 2002]  
(a) 158 g (b) 31.6 g  
(c) 790 g (d) 62 g
29. What is the concentration of nitrate ions if equal volumes of 0.1 M  $AgNO_3$  and 0.1 M  $NaCl$  are mixed together [NCERT 1981; CPMT 1983]  
(a) 0.1 N (b) 0.2 M  
(c) 0.05 M (d) 0.25 M
30. 30 ml of acid solution is neutralized by 15 ml of a 0.2 N base. The strength of acid solution is [CPMT 1986]  
(a) 0.1 N (b) 0.15 N  
(c) 0.3 N (d) 0.4 N
31. A solution containing  $Na_2CO_3$  and  $NaOH$  requires 300 ml of 0.1 N HCl using phenolphthalein as an indicator. Methyl orange is then added to the above titrated solution when a further 25 ml of 0.2 N HCl is required. The amount of  $NaOH$  present in solution is ( $NaOH = 40$ ,  $Na_2CO_3 = 106$ )  
(a) 0.6 g (b) 1.0 g  
(c) 1.5 g (d) 2.0 g
32. In the preceding question, the amount of  $Na_2CO_3$  present in the solution is [CPMT 1992]  
(a) 2.650 g (b) 1.060 g  
(c) 0.530 g (d) 0.265 g
33. How many ml of 1 (M)  $H_2SO_4$  is required to neutralise 10 ml of 1 (M)  $NaOH$  solution [MP PET 1998; MNR 1982; MP PMT 1987]  
(a) 2.5 (b) 5.0  
(c) 10.0 (d) 20.0
34. Which of the following cannot give iodometric titrations [AIIMS 1997]  
(a)  $Fe^{3+}$  (b)  $Cu^{2+}$   
(c)  $Pb^{2+}$  (d)  $Ag^+$
35.  $KMnO_4$  reacts with ferrous ammonium sulphate according to the equation  $MnO_4^- + 5Fe^{2+} + 8H^+ \rightarrow Mn^{2+} + 5Fe^{3+} + 4H_2O$ , here 10 ml of 0.1 M  $KMnO_4$  is equivalent to [CPMT 1999]  
(a) 20 ml of 0.1 M  $FeSO_4$   
(b) 30 ml of 0.1 M  $FeSO_4$   
(c) 40 ml of 0.1 M  $FeSO_4$   
(d) 50 ml of 0.1 M  $FeSO_4$
36.  $Ca(OH)_2 + H_3PO_4 \rightarrow CaHPO_4 + 2H_2O$  the equivalent weight of  $H_3PO_4$  in the above reaction is [Pb. PMT 2004]  
(a) 21 (b) 27  
(c) 38 (d) 49
37. The mass of  $BaCO_3$  produced when excess  $CO_2$  is bubbled through a solution of 0.205 mol  $Ba(OH)_2$  is [UPSEAT 2004]  
(a) 81 g (b) 40.5 g  
(c) 20.25 g (d) 162 g
38. The amount of water that should be added to 500 ml of 0.5 N solution of  $NaOH$  to give a concentration of 10 mg per ml is  
(a) 100 (b) 200  
(c) 250 (d) 500
39. Number of moles of  $KMnO_4$  required to oxidize one mole of  $Fe(C_2O_4)$  in acidic medium is [Haryana CEE 1996]  
(a) 0.6 (b) 0.167  
(c) 0.2 (d) 0.4
40. A hydrocarbon contains 86% carbon, 488ml of the hydrocarbon weight 1.68 g at STP. Then the hydrocarbon is an

- (a) Alkane (b) Alkene (c) 18 g (d) 19 g  
(c) Alkyne (d) Arene
41. The ratio of amounts of  $H_2S$  needed to precipitate all the metal ions from 100 ml of 1 M  $AgNO_3$  and 100 ml of 1 M  $CuSO_4$  will be  
(a) 1:1 (b) 1:2  
(c) 2:1 (d) None of these
42. An electric discharge is passed through a mixture containing 50 c.c. of  $O_2$  and 50 c.c. of  $H_2$ . The volume of the gases formed (i) at room temperature and (ii) at  $110^\circ C$  will be  
(a) (i) 25 c.c. (ii) 50 c.c. (b) (i) 50 c.c. (ii) 75 c.c.  
(c) (i) 25 c.c. (ii) 75 c.c. (d) (i) 75 c.c. (ii) 75 c.c.
43. 100 ml of 0.1 N hypo decolourised iodine by the addition of  $x$  g of crystalline copper sulphate to excess of KI. The value of 'x' is (molecular wt. of  $CuSO_4 \cdot 5H_2O$  is 250)  
(a) 5.0 g (b) 1.25 g  
(c) 2.5 g (d) 4 g
44. How many grams of caustic potash required to completely neutralise 12.6 gm  $HNO_3$   
(a) 22.4 KOH (b) 1.01 KOH  
(c) 6.02 KOH (d) 11.2 KOH
45. If isobutane and n-butane are present in a gas, then how much oxygen should be required for complete combustion of 5 kg of this gas  
(a) 17.9 kg (b) 9 kg  
(c) 27 kg (d) 1.8 kg
46. 16.8 litre gas containing  $H_2$  and  $O_2$  is formed at NTP on electrolysis of water. What should be the weight of electrolysed water  
(a) 5 g (b) 9 g  
(c) 10 g (d) 12 g
47. On electrical decomposition of 150 ml dry and pure  $O_2$ , 10% of  $O_2$  gets changed to O, then the volume of gaseous mixture after reaction and volume of remaining gas left after passing in turpentine oil will be  
(a) 145 ml (b) 149 ml  
(c) 128 ml (d) 125 ml
48. What should be the weight of 50% HCl which reacts with 100 g of limestone  
(a) 50% pure (b) 25% pure  
(c) 10% pure (d) 8% pure
49. What should be the weight and moles of AgCl precipitate obtained on adding 500ml of 0.20 M HCl in 30 g of  $AgNO_3$  solution? ( $AgNO_3 = 170$ )  
(a) 14.35 g (b) 15 g
50. A solution of 10 ml  $\frac{M}{10} FeSO_4$  was titrated with  $KMnO_4$  solution in acidic medium. The amount of  $KMnO_4$  used will be [CPMT 1984]  
(a) 5 ml of 0.1 M (b) 10 ml of 1.1 M  
(c) 10 ml of 0.5 M (d) 10 ml of 0.02 M
51. 1.12 ml of a gas is produced at STP by the action of 4.12 mg of alcohol, with methyl magnesium iodide. The molecular mass of alcohol is [Roorkee 1992; IIT 1993]  
(a) 16.0 (b) 41.2  
(c) 82.4 (d) 156.0
52. The simplest formula of a compound containing 50% of element X (atomic mass 10) and 50% of element Y (atomic mass 20) is [Roorkee 1994]  
(a) XY (b)  $X_2Y$   
(c)  $XY_3$  (d)  $X_2Y_3$
53. A compound contains atoms of three elements in A, B and C. If the oxidation number of A is +2, B is +5 and that of C is -2, the possible formula of the compound is [CBSE PMT 2000]  
(a)  $A_3(BC_4)_2$   
(b)  $A_3(B_4C)_2$   
(c)  $ABC_2$   
(d)  $A_2(BC_3)_2$
54. What will be the volume of  $CO_2$  at NTP obtained on heating 10 grams of (90% pure) limestone [Pb. CET 2001]  
(a) 22.4 litre  
(b) 2.016 litre  
(c) 2.24 litre  
(d) 20.16 litre
55. The ratio of the molar amounts of  $H_2S$  needed to precipitate the metal ions from 20mL each of 1M  $Cd(NO_3)_2$  and 0.5M  $CuSO_4$  is [CPMT 1997]  
(a) 1 : 1  
(b) 2 : 1  
(c) 1 : 2  
(d) Indefinite
56. 12g of Mg (at. mass 24) will react completely with acid to give [MNR 1985]  
(a) One mole of  $H_2$   
(b)  $1/2$  mole of  $H_2$   
(c)  $2/3$  mole of  $O_2$   
(d) Both  $1/2$  mol of  $H_2$  and  $1/2$  mol of  $O_2$

57. 1.5 mol of  $O_2$  combine with  $Mg$  to form oxide  $MgO$ . The mass of  $Mg$  (at. mass 24) that has combined is

[KCET 2001]

- (a) 72 g (b) 36 g  
(c) 48 g (d) 24 g

58. 100 g  $CaCO_3$  reacts with 1 litre 1 N  $HCl$ . On completion of reaction how much weight of  $CO_2$  will be obtain

[Kerala CET 2005]

- (a) 5.5 g (b) 11 g  
(c) 22 g (d) 33 g  
(e) 44 g

## Critical Thinking

### Objective Questions

1. Mixture of sand and sulphur may best be separated by

[Kerala CET 2001]

- (a) Fractional crystallisation from aqueous solution  
(b) Magnetic method  
(c) Fractional distillation  
(d) Dissolving in  $CS_2$  and filtering

2. Irrespective of the source, pure sample of water always yields 88.89% mass of oxygen and 11.11% mass of hydrogen. This is explained by the law of

[Kerala CEE 2002]

- (a) Conservation of mass (b) Constant composition  
(c) Multiple proportions (d) Constant volume

3. Zinc sulphate contains 22.65% of zinc and 43.9% of water of crystallization. If the law of constant proportions is true, then the weight of zinc required to produce 20 g of the crystals will be

- (a) 45.3 g (b) 4.53 g  
(c) 0.453 g (d) 453 g

4. 10  $dm^3$  of  $N_2$  gas and 10  $dm^3$  of gas X at the same temperature contain the same number of molecules. The gas X is

- (a)  $CO$  (b)  $CO_2$   
(c)  $H_2$  (d)  $NO$

5. The molar heat capacity of water at constant pressure is 75  $JK^{-1} mol^{-1}$ . When 1.0 kJ of heat is supplied to 100 g of water which is free to expand, the increases in temperature of water is

- (a) 6.6 K (b) 1.2 K  
(c) 2.4 K (d) 4.8 K

6. A compound possesses 8% sulphur by mass. The least molecular mass is

[AIIMS 2002]

- (a) 200 (b) 400  
(c) 155 (d) 355

7. Which of the following contains maximum number of atoms

[JIPMER 2000]

- (a)  $6.023 \times 10^{21}$  molecules of  $CO_2$

- (b) 22.4 L of  $CO_2$  at STP

- (c) 0.44 g of  $CO_2$

- (d) None of these

8. In a mole of water vapour at STP, the volume actually occupied or taken by the molecules (i.e., Avogadro's No.  $\times$  Volume of one molecule) is

[Kerala EEE 2000]

- (a) Zero  
(b) Less than 1% of 22.4 litres  
(c) About 10% of the volume of container  
(d) 1% to 2% of 22.4 litres  
(e) Between 2% to 5% of 22.4 litres

9. If  $10^{21}$  molecules are removed from 200mg of  $CO_2$ , then the number of moles of  $CO_2$  left are

[IIT 1983]

- (a)  $2.85 \times 10^{-3}$  (b)  $28.8 \times 10^{-3}$   
(c)  $0.288 \times 10^{-3}$  (d)  $1.68 \times 10^{-2}$

10. The set of numerical coefficient that balances the equation  $K_2CrO_4 + HCl \rightarrow K_2Cr_2O_7 + KCl + H_2O$  is

[Kerala CEE 2001]

- (a) 1, 1, 2, 2, 1 (b) 2, 2, 1, 1, 1  
(c) 2, 1, 1, 2, 1 (d) 2, 2, 1, 2, 1

11. One litre hard water contains 12.00 mg  $Mg^{2+}$  milli equivalent of washing soda required to remove its hardness is

- (a) 1 (b) 12.15  
(c)  $1 \times 10^{-3}$  (d)  $12.15 \times 10^{-3}$

12. In standardization of  $Na_2S_2O_3$  using  $K_2Cr_2O_7$  by iodometry, the equivalent weight of  $K_2Cr_2O_7$  is

[IIT 2000]

- (a)  $MW/2$  (b)  $MW/3$   
(c)  $MW/6$  (d)  $MW/1$

13. 3.92 g of ferrous ammonium sulphate crystals are dissolved in 100 ml of water, 20 ml of this solution requires 18 ml of  $KMnO_4$  during titration for complete oxidation. The weight of  $KMnO_4$  present in one litre of the solution is

[Tamilnadu CET 2002]

- (a) 3.476 g (b) 12.38 g  
(c) 34.76 g (d) 1.238 g

14. A 100 ml solution of 0.1 N  $HCl$  was titrated with 0.2 N  $NaOH$  solution. The titration was discontinued after adding 30 ml of  $NaOH$  solution. The remaining titration was completed by adding 0.25 N  $KOH$  solution. The volume of  $KOH$  required for completing the titration is

[DCE 1999]

- (a) 70 ml (b) 32 ml  
(c) 35 ml (d) 16 ml

15. What volume of Hydrogen gas, at 273 K and 1 atm pressure will be consumed in obtaining 21.6 g of elemental boron (atomic mass = 10.8) from the reduction of boron trichloride by Hydrogen

[AIEEE 2003]

- (a) 22.4 L (b) 89.6 L  
(c) 67.2 L (d) 44.8 L

16. The mass of 112  $cm^3$  of  $CH_4$  gas at STP is

[Karnataka CET 2001]

- (a) 0.16 g (b) 0.8 g  
(c) 0.08 g (d) 1.6 g
17. Complete combustion of 0.858 g of compound X gives 2.63 g of  $CO_2$  and 1.28 g of  $H_2O$ . The lowest molecular mass X can have [Kerala MEE 2000]  
(a) 43 g (b) 86 g  
(c) 129 g (d) 172 g
18. In the following reaction, which choice has value twice that of the equivalent mass of the oxidising agent  
 $SO_2 + H_2O \longrightarrow 3S + 2H_2O$  [DPMT 2000]  
(a) 64 (b) 32  
(c) 16 (d) 48

## Assertion & Reason

For AIIMS Aspirants

Read the assertion and reason carefully to mark the correct option out of the options given below :

- (a) If both assertion and reason are true and the reason is the correct explanation of the assertion.  
(b) If both assertion and reason are true but reason is not the correct explanation of the assertion.  
(c) If assertion is true but reason is false.  
(d) If the assertion and reason both are false.  
(e) If assertion is false but reason is true.

1. Assertion : Volume of a gas is inversely proportional to the number of moles of a gas.  
Reason : The ratio by volume of gaseous reactants and products is in agreement with their mole ratio. [AIIMS 1995]
2. Assertion : Molecular weight of oxygen is 16.  
Reason : Atomic weight of oxygen is 16. [AIIMS 1996]
3. Assertion : Atoms can neither be created nor destroyed.  
Reason : Under similar condition of temperature and pressure, equal volume of gases does not contain equal number of atoms. [AIIMS 1994,2002]
4. Assertion : One mole of  $SO_2$  contains double the number of molecules present in one mole of  $O_2$ .  
Reason : Molecular weight of  $SO_2$  is double to that of  $O_2$ .
5. Assertion : 1.231 has three significant figures.  
Reason : All numbers right to the decimal point are significant.
6. Assertion : 22.4 L of  $N_2$  at NTP and 5.6 L  $O_2$  at NTP contain equal number of molecules.

- Reason : Under similar conditions of temperature and pressure all gases contain equal number of molecules.
7. Assertion : One atomic mass unit (amu) is mass of an atom equal to exactly one-twelfth the mass of a carbon-12 atom.  
Reason : Carbon-12 isotope was selected as standard.
8. Assertion : Molecular mass of A is  $\frac{M}{4}$  if the molecular mass of B is M.  
Reason : Vapour density of A four times that of B.
9. Assertion : Pure water obtained from different sources such as, river, well, spring, sea etc. always contains hydrogen and oxygen combined in the ratio 1 : 8 by mass.  
Reason : A chemical compound always contains elements combined together in same proportion by mass, it was discovered by French chemist, Joseph Proust (1799).
10. Assertion : As mole is the basic chemical unit, the concentration of the dissolved solute is usually specified in terms of number of moles of solute.  
Reason : The total number of molecules of reactants involved in a balanced chemical equation is known as molecularity of the reaction.
11. Assertion : A certain element X, forms three binary compounds with chlorine containing 59.68%, 68.95% and 74.75% chlorine respectively. These data illustrate the law of multiple proportions.  
Reason : According to law of multiple proportions, the relative amounts of an element combining with some fixed amount of a second element in a series of compounds are the ratios of small whole numbers.
12. Assertion : Equivalent weight of Cu in  $CuO$  is 63.6 and in  $Cu_2O$  31.8.  
Reason : Equivalent weight of an element =  $\frac{\text{Atomic weight of the element}}{\text{Valency of the element}}$
13. Assertion : Mass spectrometer is used for the determination of isotopes.  
Reason : Isotopes are the atoms of same element differing in mass numbers.
14. Assertion : Gases combine in simple ratio of their volume but, not always.  
Reason : Gases deviate from ideal behaviour.
15. Assertion : Isomorphous substances form crystals of same shape and can grow in saturated solution of each other.  
Reason : They have similar constitution and chemical formulae.
16. Assertion : Atomicity of oxygen is 2.

Reason : 1 mole of an element contains  $6.023 \times 10^{23}$  atoms.

17. Assertion : 1 amu equals to  $1.66 \times 10^{-24}$  g .

Reason :  $1.66 \times 10^{-24}$  g equals to  $\frac{1}{12}$ th of mass of a  $C^{12}$  atom.

# Answers

## Significant figures, Units for measurement, Matter and Separation of mixture

1	a	2	d	3	a	4	c	5	d
6	b	7	c	8	d	9	c	10	c
11	c	12	b	13	a	14	c	15	b
16	b	17	b	18	a	19	a	20	c
21	b	22	d	23	a	24	a	25	b
26	b	27	d						

## Laws of chemical combination

1	a	2	d	3	c	4	a	5	c
6	c	7	c	8	b	9	b	10	a
11	c	12	b	13	a	14	d	15	b
16	a	17	c	18	d	19	c	20	a
21	c	22	d						

## Atomic, Molecular and Equivalent masses

1	c	2	b	3	a	4	a	5	b
6	c	7	d	8	b	9	a	10	b
11	a	12	b	13	a	14	c	15	b
16	c	17	a	18	d	19	a	20	a
21	b	22	b	23	d	24	c	25	a
26	a	27	c	28	d	29	a	30	c
31	a	32	d	33	b	34	a	35	c
36	b	37	b	38	c	39	a	40	b
41	c	42	d	43	a	44	d	45	d
46	b	47	c	48	c	49	b	50	b
51	a	52	b	53	b	54	c	55	b
56	a	57	d	58	d	59	a	60	d
61	c	62	d	63	b				

## The mole concept

1	a	2	b	3	b	4	c	5	a
6	b	7	a	8	c	9	d	10	c
11	b	12	c	13	c	14	d	15	a
16	a	17	b	18	b	19	a	20	a
21	b	22	b	23	c	24	a	25	a
26	c	27	d						

## Percentage composition & Molecular formula

1	a	2	a	3	b	4	d	5	a
6	b	7	b						

## Chemical stoichiometry

1	c	2	c	3	a	4	c	5	c
6	c	7	c	8	b	9	a	10	d
11	c	12	d	13	d	14	b	15	a
16	b	17	b	18	c	19	d	20	c
21	d	22	c	23	d	24	a	25	d
26	b	27	b	28	b	29	c	30	a
31	b	32	c	33	b	34	c	35	d
36	d	37	b	38	d	39	a	40	b
41	b	42	c	43	c	44	d	45	a
46	b	47	a	48	a	49	a	50	d
51	c	52	b	53	a	54	b	55	b
56	b	57	a	58	c				

## Critical Thinking Questions

1	d	2	b	3	b	4	a	5	c
6	b	7	b	8	b	9	a	10	d
11	a	12	c	13	a	14	d	15	c
16	c	17	a	18	b				

## Assertion & Reason

1	e	2	e	3	c	4	e	5	d
6	d	7	a	8	c	9	a	10	b
11	a	12	e	13	e	14	a	15	a
16	b	17	a						

# AS Answers and Solutions

**Significant figures, Units of measurement, Matter and Separation of mixture**

4. (c)  $\text{Pressure} = \frac{\text{Force}}{\text{Area}} = \frac{[MLT^{-2}]}{[L^2]} = [ML^{-1}T^{-2}]$
- $\text{Energy per unit volume} = \frac{[ML^2T^{-2}]}{[L^3]} = [ML^{-1}T^{-2}]$
17. (b)  $\frac{(29.2 - 20.2)(1.79 \times 10^5)}{1.37} = \frac{9.0 \times 1.79 \times 10^5}{1.37}$
- Least precise terms *i.e.*, 9.0 has only two significant figures.  
Hence, final answer will have two significant figures.
18. (a) Pure ethyl alcohol =  $81.4 - 0.002 = 81.398$ .
19. (a)  $JPa^{-1}$ ; Unit of work is *Joule* and unit of pressure is *Pascal*.  
Dimension of *Joule i.e.* work =  $F \times L = MLT^{-2} \times L$   
 $= [ML^2T^{-2}]$
- $\frac{1}{Pa} = \frac{1}{\text{Pressure}} = \frac{1}{\frac{F}{A}} = \frac{1 \times A}{F} = [MLT^{-1}]$
- So,  $JPa^{-1} = [ML^2T^{-2}] = [L^2 \times L] = [L^3]$ .
22. (d) 1 zepto =  $10^{-21}$
23. (a) As we know that all non zero unit are significant number.  
Therefore significant figure is 2.
24. (a) Number of significant figures in 6.0023 are 5 because all the zeroes stand between two non zero digit are counted towards significant figures.
25. (b) Given  $P = 0.0030m$ ,  $Q = 2.40m$  &  $R = 3000m$  In  $P(0.0030)$  initial zeros after the decimal point are not significant. Therefore, significant figures in  $P(0.0030)$  are 2. Similarly in  $Q(2.40)$  significant figures are 3 as in this case final zero is significant. In  $R = (3000)$  all the zeroes are significant hence, in  $R$  significant figures are 4.
26. (b) All the zeroes between two non zero digit are significant. Hence in 60.0001 significant figures is 6.
27. (d) Round off the digit at 2<sup>nd</sup> position of decimal 3.929 = 3.93.

**Laws of chemical combination**

12. (b)  $\begin{matrix} X & + & Y & \rightleftharpoons & R & + & S \\ ng & mg & pg & & qg \end{matrix}$
- $n + m = p + q$  by law of conservation of mass.

**Atomic, Molecular and Equivalent masses**

6. (c) As the given sulphate is isomorphous with  $ZnSO_4 \cdot 7H_2O$  its formula would be  $MSO_4 \cdot 7H_2O$ .  $m$  is the atomic weight of  $M$ , molecular weight of  $MSO_4 \cdot 7H_2O$  =  $m + 32 + 64 + 126 = m + 222$
- Hence % of  $M = \frac{m}{m + 222} \times 100 = 9.87$  (given) or
- $100m = 9.87m + 222 \times 9.87$  or  $90.13m = 222 \times 9.87$
- or  $m = \frac{222 \times 9.87}{90.13} = 24.3$ .
5. (b) The atomic weight of sulphur = 32  
In  $SCl_2$  valency of sulphur = 2
- So equivalent mass of sulphur =  $\frac{32}{2} = 16$ .