

9. In a catalytic conversion of  $N_2$  to  $NH_3$  by Haber's process, the rate of reaction was expressed as change in the concentration of ammonia per time is  $40 \times 10^{-3} \text{ mol litre}^{-1} \text{ s}^{-1}$ . If there are no side reaction, the rate of the reaction as expressed in terms of hydrogen is (in  $\text{mol litre}^{-1} \text{ s}^{-1}$ )
- (a)  $60 \times 10^{-3}$  (b)  $20 \times 10^{-3}$   
(c) 1.200 (d)  $10.3 \times 10^{-3}$
10. If the concentration of the reactants is increased, the rate of reaction [MP PMT 1989]
- (a) Remains unaffected (b) Increases  
(c) Decreases (d) May increase or decrease
11. Time required for completion of ionic reactions in comparison to molecular reactions is
- (a) Maximum (b) Minimum  
(c) Equal (d) None
12. For reaction  $2A + B \rightarrow$  products, the active mass of  $B$  is kept constant and that of  $A$  is doubled. The rate of reaction will then
- (a) Increase 2 times (b) Increase 4 times  
(c) Decrease 2 times (d) Decrease 4 times
13. In a reaction  $2A + B \rightarrow A_2B$ , the reactant  $A$  will disappear at [MP PET 1993]
- (a) Half the rate that  $B$  will decrease  
(b) The same rate that  $B$  will decrease  
(c) Twice the rate that  $B$  will decrease  
(d) The same rate that  $A_2B$  will form
14. The rate of a gaseous reaction is given by the expression  $K[A][B]$ . If the volume of the reaction vessel is suddenly reduced to  $1/4$ th of the initial volume, the reaction rate relating to original rate will be [Roorkee 1992]
- (a)  $1/10$  (b)  $1/8$   
(c) 8 (d) 16
15. A catalyst increases the rate of reaction because it [EAMCET 1992]
- (a) Increases the activation energy  
(b) Decreases the energy barrier for reaction  
(c) Decreases the collision diameter  
(d) Increases the temperature coefficient
16. For the reaction  $N_2(g) + 3H_2(g) \rightarrow 2NH_3(g)$  under certain conditions of temperature and partial pressure of the reactants, the rate of formation of  $NH_3$  is  $0.001 \text{ kg h}^{-1}$ . The rate of conversion of  $H_2$  under the same conditions is [IIT 1994]
- (a)  $1.82 \times 10^{-4} \text{ kg/hr}$  (b)  $0.0015 \text{ kg/hr}$   
(c)  $1.52 \times 10^{-4} \text{ kg/hr}$  (d)  $1.82 \times 10^{-14} \text{ kg/hr}$
17. In the reaction  $2A + B \rightarrow A_2B$ , if the concentration of  $A$  is doubled and of  $B$  is halved, then the rate of the reaction will
- (a) Increase by four times (b) Decrease by two times  
(c) Increase by two times (d) Remain the same
18. The term  $\left(-\frac{dc}{dt}\right)$  in a rate equation refers to the [MP PMT 1996]
- (a) Concentration of the reactant  
(b) Decrease in concentration of the reactant with time  
(c) Increase in concentration of the reactant with time  
(d) Velocity constant of the reaction
19. The rate of a reaction depends upon the [Pb. PMT 1999]
- (a) Volume  
(b) Force  
(c) Pressure  
(d) Concentration of reactant
20. For a given reaction  $3A + B \rightarrow C + D$  the rate of reaction can be represented by [DCE 2000]
- (a)  $-\frac{1}{3} \frac{d[A]}{dt} = -\frac{d[B]}{dt} = \frac{d[C]}{dt} = \frac{d[D]}{dt}$   
(b)  $-\frac{1}{3} \frac{d[A]}{dt} = \frac{d[C]}{dt} = K[A]^m[B]^n$   
(c)  $+\frac{1}{3} \frac{d[A]}{dt} = -\frac{d[C]}{dt} = K[A]^n[B]^m$   
(d) None of these
21. For the reaction  $N_2 + 3H_2 \rightarrow 2NH_3$  if  $\frac{\Delta[NH_3]}{\Delta t} = 2 \times 10^{-4} \text{ mol l}^{-1} \text{ s}^{-1}$ , the value of  $\frac{-\Delta[H_2]}{\Delta t}$  would be [MP PMT 2000]
- (a)  $1 \times 10^{-4} \text{ mol l}^{-1} \text{ s}^{-1}$  (b)  $3 \times 10^{-4} \text{ mol l}^{-1} \text{ s}^{-1}$   
(c)  $4 \times 10^{-4} \text{ mol l}^{-1} \text{ s}^{-1}$  (d)  $6 \times 10^{-4} \text{ mol l}^{-1} \text{ s}^{-1}$
22. A gaseous hypothetical chemical equation  $2A \rightleftharpoons 4B + C$  is carried out in a closed vessel. The concentration of  $B$  is found to increase by  $5 \times 10^{-3} \text{ mol l}^{-1}$  in 10 second. The rate of appearance of  $B$  is [AFMC 2001]
- (a)  $5 \times 10^{-4} \text{ mol l}^{-1} \text{ sec}^{-1}$  (b)  $5 \times 10^{-5} \text{ mol l}^{-1} \text{ sec}^{-1}$   
(c)  $6 \times 10^{-5} \text{ mol l}^{-1} \text{ sec}^{-1}$  (d)  $4 \times 10^{-4} \text{ mol l}^{-1} \text{ sec}^{-1}$
23. The rate of a chemical reaction depends upon [AFMC 2002]
- (a) Time (b) Pressure  
(c) Concentration (d) All of these
24. The rate of disappearance of  $SO_2$  in the reaction  $2SO_2 + O_2 \rightarrow 2SO_3$  is  $1.28 \times 10^{-3} \text{ g/sec}$  then the rate of formation of  $SO_3$  is
- (a)  $0.64 \times 10^{-3} \text{ g/sec}$  (b)  $0.80 \times 10^{-3} \text{ g/sec}$   
(c)  $1.28 \times 10^{-3} \text{ g/sec}$  (d)  $1.60 \times 10^{-3} \text{ g/sec}$
25. When the concentration of  $A$  in the reaction  $A + B \rightleftharpoons AB$  is doubled, the rate of reaction will be [MP PMT 2002]
- (a) Doubled (b) Decreased by half  
(c) Unchanged (d) Increased by four times
26. The velocity of the chemical reaction doubles every  $10^\circ \text{C}$  rise of temperature. If the temperature is raised by  $50^\circ \text{C}$ , the velocity of the reaction increases to about
- (a) 32 times (b) 16 times  
(c) 20 times (d) 50 times
27. An increase in temperature by  $10^\circ \text{C}$ , generally increases the rate of a reaction by [MP PET 1994; CBSE PMT 2000]
- (a) 2 times (b) 10 times  
(c) 9 times (d) 100 times
28. The temperature coefficient for reaction in which food deteriorates is 2. Then food deteriorates ..... times as rapidly at  $25^\circ \text{C}$  as it does at  $5^\circ \text{C}$
- (a) Two (b) Four  
(c) Six (d) Twenty
29. The rate of a reaction is doubled for every  $10^\circ$  rise in temperature. The increase in reaction rate as a result of temperature rise from  $10^\circ$  to  $100^\circ$  is [KCET 1993; Kerala PET 2002; MP PET 2003]

- (a) 112 (b) 512  
(c) 400 (d) 614
30. A catalyst increases the rate of a chemical reaction by [MNR 1988; CPMT 1999; Pb. PMT 2000]  
(a) Increasing the activation energy  
(b) Decreasing the activation energy  
(c) Reacting with reactants  
(d) Reacting with products
31. Velocity constant of a reaction at 290 K was found to be  $3.2 \times 10^{-3}$ . At 310 K it will be about [KCET 1989, 91]  
(a)  $1.28 \times 10^{-2}$  (b)  $9.6 \times 10^{-3}$   
(c)  $6.4 \times 10^{-3}$  (d)  $3.2 \times 10^{-4}$
32. The temperature coefficient of a reaction is  
(a) Specific reaction rate at  $25^\circ\text{C}$   
(b) Rate of the reaction at  $100^\circ\text{C}$   
(c) Ratio of the rate constants at temperatures  $35^\circ\text{C}$  and  $25^\circ\text{C}$   
(d) Ratio of the rate constants at two temperatures differing by  $1^\circ\text{C}$
33. The main function of a catalyst in speeding up a reaction is  
(a) To increase the rate of the forward reaction  
(b) To change the reaction path so as to decrease the energy of activation for the reaction  
(c) To reduce the temperature at which the reaction can occur  
(d) To increase the energy of the molecules of the reactants
34. The rate of a reaction [CPMT 1973]  
(a) Increases with increase in temperature  
(b) Decreases with increase in temperature  
(c) Does not depend on temperature  
(d) Does not depend on concentration
35. Which of the following statements is false in relation to enzyme  
(a) pH affects their functioning  
(b) Temperature affects their functioning  
(c) They always increase activation energy  
(d) Their reactions are specific
36. A reaction is catalysed by 'X'. Here 'X' [MP PMT 2003]  
(a) Decreases the rate constant of reaction  
(b) Does not affect the equilibrium constant of reaction  
(c) Decreases the enthalpy of reaction  
(d) Decreases the activation energy
37. Which reaction characteristics are changing by the addition of a catalyst to a reaction at constant temperature  
(i) Activation energy (ii) Equilibrium constant  
(iii) Reaction entropy (iv) Reaction enthalpy [DCE 2003]  
(a) (i) Only (b) (iii) only  
(c) (i) and (ii) only (d) All of these
38. The velocity constant of a reaction at 290 K was found to be  $3.2 \times 10^{-3}$ . At 300 K it will be [MP PMT 2004]  
(a)  $1.28 \times 10^{-2}$  (b)  $6.4 \times 10^{-3}$   
(c)  $9.6 \times 10^{-3}$  (d)  $3.2 \times 10^{-4}$
39. In which of the following cases, does the reaction go farthest to completion [UPSEAT 2001]  
(a)  $K = 10^3$  (b)  $K = 10^{-2}$   
(c)  $K = 10$  (d)  $K = 1$
40. Rate of reaction [Pb. CET 2004]  
(a) Decreases with increase in temperature  
(b) Increases with increase in temperature  
(c) May increase or decrease with increase in temperature  
(d) Does not depend on temperature
41. For the reaction  $2\text{N}_2\text{O}_{5(g)} \rightarrow 4\text{NO}_{2(g)} + \text{O}_{2(g)}$ , if concentration of  $\text{NO}_2$  in 100 seconds is increased by  $5.2 \times 10^{-3} \text{ m}$ . Then rate of reaction will be [Kerala CET 2005]  
(a)  $1.3 \times 10^{-5} \text{ ms}^{-1}$  (b)  $5 \times 10^{-4} \text{ ms}^{-1}$   
(c)  $7.6 \times 10^{-4} \text{ ms}^{-1}$  (d)  $2 \times 10^{-3} \text{ ms}^{-1}$   
(e)  $2.5 \times 10^{-5} \text{ ms}^{-1}$
42. A first order reaction complete its 10% in 20 minutes then time required to complete its 19% is [Kerala CET 2005]  
(a) 30 minutes (b) 40 minutes  
(c) 50 minutes (d) 38 minutes  
(e) 45 minutes

### Rate law and Rate constant

1. Which of these does not influence the rate of reaction [KCET 2005]  
(a) Nature of the reactants  
(b) Concentration of the reactants  
(c) Temperature of the reaction  
(d) Molecularity of the reaction
2. The rate law for reaction  $A + 2B \rightarrow C + 2D$  will be  
(a)  $\text{Rate} = K[A][B]$  (b)  $\text{Rate} = K[A][2B]$   
(c)  $\text{Rate} = K[A][B]^2$  (d)  $\text{Rate} = K \frac{[C][D]^2}{[A][B]^2}$
3. In the reaction  $2\text{N}_2\text{O}_5 \rightarrow 4\text{NO}_2 + \text{O}_2$ , initial pressure is 500 atm and rate constant  $K$  is  $3.38 \times 10^{-5} \text{ sec}^{-1}$ . After 10 minutes the final pressure of  $\text{N}_2\text{O}_5$  is [Orissa JEE 2005]  
(a) 490 atm (b) 250 atm  
(c) 480 atm (d) 420 atm
4. The rate law for the reaction  
 $\text{Sucrose} + \text{Water} \xrightarrow{H^+} \text{Glucose} + \text{Fructose}$  is given by  
(a)  $\text{Rate} = K [\text{sucrose}] [\text{water}]$   
(b)  $\text{Rate} = K [\text{sucrose}] [\text{water}]^0$   
(c)  $\text{Rate} = K [\text{sucrose}]^0 [\text{water}]$   
(d)  $\text{Rate} = K [\text{sucrose}]^{1/2} [\text{water}]^{1/2}$
5.  $A + 2B \rightarrow C + D$ . If  $-\frac{d[A]}{dt} = 5 \times 10^{-4} \text{ mol l}^{-1} \text{ s}^{-1}$ , then  $-\frac{d[B]}{dt}$  is [DPMT 2005]  
(a)  $2.5 \times 10^{-4} \text{ mol l}^{-1} \text{ s}^{-1}$  (b)  $5.0 \times 10^{-4} \text{ mol l}^{-1} \text{ s}^{-1}$   
(c)  $2.5 \times 10^{-3} \text{ mol l}^{-1} \text{ s}^{-1}$  (d)  $1.0 \times 10^{-3} \text{ mol l}^{-1} \text{ s}^{-1}$
6. The data for the reaction  $A + B \rightarrow C$  is [CBSE PMT 1994]

Exp.	$[A]_0$	$[B]_0$	Initial rate
(1)	0.012	0.035	0.10
(2)	0.024	0.070	0.80
(3)	0.024	0.035	0.10

(4)	0.012	0.070	0.80
-----	-------	-------	------

The rate law corresponds to the above data is

- (a)  $\text{Rate} = k[B]^3$  (b)  $\text{Rate} = k[B]^4$   
 (c)  $\text{Rate} = k[A][B]^3$  (d)  $\text{Rate} = k[A]^2[B]^2$

7. The experimental data for the reaction  $2A + B_2 \rightarrow 2AB$  is

[CBSE PMT 1997]

Exp.	$[A]_0$	$[B]_0$	Rate ( $\text{mole s}^{-1}$ )
(1)	0.50	0.50	$1.6 \times 10^{-4}$
(2)	0.50	1.00	$3.2 \times 10^{-4}$
(3)	1.00	1.00	$3.2 \times 10^{-4}$

The rate equation for the above data is

- (a)  $\text{Rate} = k[B_2]$  (b)  $\text{Rate} = k[B_2]^2$   
 (c)  $\text{Rate} = k[A]^2[B]^2$  (d)  $\text{Rate} = k[A]^2[B]$

8. The reaction  $2NO(g) + O_2(g) \rightleftharpoons 2NO_2(g)$  is of first order. If volume of reaction vessel is reduced to  $1/3$ , the rate of reaction would be [MP PMT 2001]

- (a)  $1/3$  times (b)  $2/3$  times  
 (c) 3 times (d) 6 times

9. For a reaction  $2A + B \rightarrow \text{Products}$ , doubling the initial concentration of both the reactants increases the rate by a factor of 8, and doubling the concentration of  $B$  alone doubles the rate. The rate law for the reaction is

[MP PET 2001]

- (a)  $\gamma = k[A][B]^2$  (b)  $\gamma = k[A]^2[B]$   
 (c)  $\gamma = k[A][B]$  (d)  $\gamma = k[A]^2[B]^2$

10. For a reactions  $A + B \rightarrow \text{product}$ , it was found that rate of reaction increases four times if concentration of 'A' is doubled, but the rate of reaction remains unaffected. If concentration of 'B' is doubled. Hence, the rate law for the reaction is [MP PET/PMT 1998; MP PMT 2003]

- (a)  $\text{rate} = k[A][B]$  (b)  $\text{rate} = k[A]^2$   
 (c)  $\text{rate} = k[A]^2[B]^1$  (d)  $\text{rate} = k[A]^2[B]^2$

11. Velocity constant  $K$  of a reaction is affected by

- (a) Change in the concentration of the reactant  
 (b) Change of temperature  
 (c) Change in the concentration of the product  
 (d) None of the above

12. Point out the wrong statement :

For a first order reaction

- (a) Time for half-change ( $t_{1/2}$ ) is independent of initial concentration  
 (b) Change in the concentration unit does not change the rate constant ( $K$ )  
 (c) Time for half-change  $\times$  rate constant = 0.693  
 (d) The unit of  $K$  is  $\text{mole}^{-1} \text{min}^{-1}$

13. The rate constant of a reaction depends on

[CPMT 1989; DPMT 2001]

- (a) Temperature (b) Mass  
 (c) Weight (d) Time

14. In a first order reaction the concentration of reactant decreases from  $800 \text{ mol/dm}^3$  to  $50 \text{ mol/dm}^3$  is  $2 \times 10^2 \text{ sec}$ . The rate constant of reaction in  $\text{sec}^{-1}$  is

[IIT-JEE (Screening) 2003]

- (a)  $2 \times 10^4$  (b)  $3.45 \times 10^{-5}$   
 (c)  $1.386 \times 10^{-2}$  (d)  $2 \times 10^{-4}$

15. For a reaction  $A \rightarrow B$ , the rate of reaction quadrupled when the concentration of A is doubled. The rate expression of the reaction is  $r = K(A)^n$ . when the value of  $n$  is

- (a) 1 (b) 0  
 (c) 3 (d) 2

16. The velocity constant of a reaction is  $K$ . Which of the following statements is not true regarding  $K$

- (a)  $K$  is a constant for a reaction at a given temperature  
 (b) The value of  $K$  changes when the temperature changes  
 (c)  $K$  is the velocity of the reaction at unit concentrations of the reactant  
 (d)  $K$  is a constant for all reactions

17. For the following reaction scheme (homogeneous), the rate constant has units :  $A + B \xrightarrow{K} C$  [MP PET 1999]

- (a)  $\text{sec}^{-1} \text{ mole}$  (b)  $\text{sec}^{-1}$   
 (c)  $\text{sec}^{-1} \text{ litremole}^{-1}$  (d)  $\text{sec}$

18. Which of the following oxides of nitrogen will be the most stable one [NCERT 1978]

- (a)  $2NO_2(g) \rightleftharpoons N_2(g) + 2O_2(g); K = 6.7 \times 10^{16} \text{ mol l}^{-1}$   
 (b)  $2NO(g) \rightleftharpoons N_2(g) + O_2(g); K = 2.2 \times 10^{30} \text{ mol l}^{-1}$   
 (c)  $2N_2O_5(g) \rightleftharpoons 2N_2(g) + 5O_2(g); K = 1.2 \times 10^{34} \text{ mol l}^{-5}$   
 (d)  $2N_2O(g) \rightleftharpoons 2N_2(g) + O_2(g); K = 3.5 \times 10^{33} \text{ mol litre}^{-1}$

19. The rate of reaction is determined by slow step reaction. The step is called

- (a) Reaction rate (b) Activation step  
 (c) Rate determining step (d) None of the above

20. The rate of the reaction


$CCl_3CHO + NO \rightarrow CHCl_3 + NO + CO$  is given by Rate =  $K[CCl_3CHO][NO]$ . If concentration is expressed in moles/litre, the units of  $K$  are [MP PET 1993]

- (a)  $\text{litre}^2 \text{ mole}^{-2} \text{ sec}^{-1}$  (b)  $\text{mole litre}^{-1} \text{ sec}^{-1}$   
 (c)  $\text{litre mole}^{-1} \text{ sec}^{-1}$  (d)  $\text{sec}^{-1}$

21. Rate constant for a reaction  $H_2 + I_2 \rightarrow 2HI$  is 49, then rate constant for reaction  $2HI \rightarrow H_2 + I_2$  is [Bihar MEE 1997]

- (a) 7 (b)  $1/49$   
 (c) 49 (d) 21  
 (e) 63

22. The reaction



- is of first order in  $N_2O_5$  with rate constant  $6.2 \times 10^{-1} s^{-1}$ . What is the value of rate of reaction when  $[N_2O_5] = 1.25 \text{ mole l}^{-1}$  [AFMC 1998]
- (a)  $7.75 \times 10^{-1} \text{ mole l}^{-1} s^{-1}$  (b)  $6.35 \times 10^{-3} \text{ mole l}^{-1} s^{-1}$   
(c)  $5.15 \times 10^{-5} \text{ mole l}^{-1} s^{-1}$  (d)  $3.85 \times 10^{-1} \text{ mole l}^{-1} s^{-1}$
23. A reaction that is of the first order with respect to reactant A has a rate constant  $6 \text{ min}^{-1}$ . If we start with  $[A] = 0.5 \text{ mol l}^{-1}$ , when would  $[A]$  reach the value  $0.05 \text{ mol l}^{-1}$  [KCET 2000]
- (a) 0.384 min (b) 0.15 min  
(c) 3 min (d) 3.84 min
24. The rate constant for the reaction,  $2N_2O_5 \rightarrow 4NO_2 + O_2$  is  $3 \times 10^{-5} \text{ sec}^{-1}$ . If the rate is  $2.40 \times 10^{-5} \text{ mol litre}^{-1} \text{ sec}^{-1}$ . Then the concentration of  $N_2O_5$  (in  $\text{mol litre}$ ) is [IIT Screening 2000]
- (a) 1.4 (b) 1.2  
(c) 0.04 (d) 0.8
25. For the reaction  $2N_2O_5 \rightarrow 4NO_2 + O_2$  rate of reaction and rate constant are  $1.02 \times 10^{-4}$  and  $3.4 \times 10^{-5} \text{ sec}^{-1}$  respectively. The concentration of  $N_2O_5$  at that time will be [BHU 2001]
- (a) 1.732 (b) 3  
(c)  $1.02 \times 10^{-4}$  (d)  $3.4 \times 10^5$
26. The rate law of the reaction  $2N_2O_5 \rightarrow 4NO_2 + O_2$  is [MP PET 2002]
- (a)  $r = K[N_2O_5]$  (b)  $r = K[N_2O_5]^2$   
(c)  $r = K[N_2O_5]^0$  (d)  $r = K[NO_2]^4 [O_2]$
27. If  $R = K[NO]^2 [O_2]$ , rate constant may be increased by [BHU 2003]
- (a) Increasing temperature  
(b) Decreasing temperature  
(c) Increasing concentration of  $O_2$   
(d) Increasing concentration of  $NO$
28. The value of rate constant  $A + B \rightarrow \text{products}$  depends on [BHU 2003]
- (a) Concentration of A and B  
(b) Pressure  
(c) Temperature  
(d) All of these
29. The rate constant of a reaction depends upon [BHU 2004]
- (a) Extent of reaction  
(b) Time of reaction  
(c) Temperature of the system  
(d) Concentration of the system
30. The rate equation for the reaction  $2A + B \rightarrow C$  is found to be: rate  $= k[A][B]$ . The correct statement in relation to this reaction is that the [AIEEE 2004]
- (a) Rate of formation of C is twice the rate of disappearance of A  
(b)  $t_{1/2}$  is a constant  
(c) Unit of  $k$  must be  $s^{-1}$   
(d) Value of  $k$  is independent of the initial concentrations of A and B
31. The specific rate constant of a first order reaction depends on the [IIT 1981, 83; DPMT 1991; Bihar MEE 1995; KCET 1998]
- (a) Concentration of the reactants  
(b) Concentration of the products  
(c) Time of reaction  
(d) Temperature of reaction
32. If the concentration is expressed in moles per litre, the unit of the rate constant for a first order reaction is [MNR 1986; MP PET 1994, 2000, 01; Bihar MEE 1996; CPMT 1997; MP PMT 1995, 96, 99; AFMC 2002]
- (a)  $\text{mole litre}^{-1} \text{ sec}^{-1}$  (b)  $\text{mole litre}^{-1}$   
(c)  $\text{sec}^{-1}$  (d)  $\text{mole}^{-1} \text{ litre}^{-1} \text{ sec}^{-1}$
33. The dimension of rate constant of a second order reaction involves
- (a) Neither time nor concentration  
(b) Only time  
(c) Time and concentration  
(d) Time and square of concentration
34. The unit of rate constant of second order reaction is usually expressed as [NCERT 1983, 84; MNR 1983; MP PMT 1994, 99]
- (a)  $\text{mole litre sec}^{-1}$  (b)  $\text{mole}^{-1} \text{ litre}^{-1} \text{ sec}^{-1}$   
(c)  $\text{mole litre}^{-1} \text{ sec}^{-1}$  (d)  $\text{mole}^{-1} \text{ litre sec}^{-1}$
35. A zero order reaction is one whose rate is independent of [NCERT 1981]
- (a) Temperature of the reaction  
(b) The concentrations of the reactants  
(c) The concentration of the products  
(d) The material of the vessel in which the reaction is carried out
36. The unit of rate constant for a zero order reaction is [NCERT 1981; MP PMT 2000; RPET 2000]
- (a)  $\text{litre sec}^{-1}$  (b)  $\text{litre mole}^{-1} \text{ sec}^{-1}$   
(c)  $\text{mole litre}^{-1} \text{ sec}^{-1}$  (d)  $\text{mole sec}^{-1}$
37. Which of the following rate laws has an overall order of 0.5 for reaction involving substances x, y and z [AIIMS 1983]
- (a) Rate  $= K(C_x)(C_y)(C_z)$   
(b) Rate  $= K(C_x)^{0.5} (C_y)^{0.5} (C_z)^{0.5}$   
(c) Rate  $= K(C_x)^{1.5} (C_y)^{-1} (C_z)^0$   
(d) Rate  $= K(C_x)(C_z)^n / (C_y)^2$
38. The rates of a certain reaction ( $dc/dt$ ) at different times are as follows
- | Time | Rate ( $\text{mole litre}^{-1} \text{ sec}^{-1}$ ) |
|------|--|
| 0    | $2.8 \times 10^{-2}$                               |
| 10   | $2.78 \times 10^{-2}$                              |
| 20   | $2.81 \times 10^{-2}$                              |
| 30   | $2.79 \times 10^{-2}$                              |
- The reaction is [NCERT 1978]
- (a) Zero order (b) First order  
(c) Second order (d) Third order
39. For a chemical reaction  $A \rightarrow B$  it is found that the rate of reaction doubles, when the concentration of A is increased four times. The order in A for this reaction is [NCERT 1979; AIIMS 1997; J & K CET 2005]
- (a) Two (b) One  
(c) Half (d) Zero
40. The following data are for the decomposition of ammonium nitrate in aqueous solution

- |                       |                |  |
|-----------------------|----------------|--|
| Volume of $N_2$ in cc | Time (minutes) |  |
| 6.25                  | 10             |  |
| 9.50                  | 15             |  |
| 11.42                 | 20             |  |
| 13.65                 | 25             |  |
| 35.05                 | Finally        |  |
- The order of the reaction is [NCERT 1980]
- (a) Zero (b) One  
(c) Two (d) Three
41. The hydrolysis of ethyl acetate is a reaction of [MP PMT 1987]
- $$CH_3COOEt + H_2O \xrightarrow{H^+} CH_3COOH + EtOH$$
- (a) First order (b) Second order  
(c) Third order (d) Zero order
42. The rate of reaction between  $A$  and  $B$  increases by a factor of 100, when the concentration of  $A$  is increased 10 folds. The order of reaction with respect to  $A$  is [CPMT 1985]
- (a) 10 (b) 1  
(c) 4 (d) 2
43. Which of the following is a first order reaction [MP PMT 1987]
- (a)  $NH_4NO_2 \rightarrow N_2 + 2H_2O$   
(b)  $2HI \rightarrow H_2 + I_2$   
(c)  $2NO_2 \rightarrow 2NO + O_2$   
(d)  $2NO + O_2 \rightarrow 2NO_2$
44. The inversion of cane sugar is represented by  
 $C_{12}H_{22}O_{11} + H_2O \rightarrow C_6H_{12}O_6 + C_6H_{12}O_6$   
 It is a reaction of [AFMC 1982; MP PMT 1993; RPET 2000]
- (a) Second order (b) Unimolecular  
(c) Pseudo unimolecular (d) None of the three
45. Which one of the following formula represents a first order reaction
- (a)  $K = \frac{x}{t}$  (b)  $K = \frac{1}{2t} \left[ \frac{1}{(a-x)^2} - \frac{1}{a^2} \right]$   
 (c)  $K = \frac{2.303}{t} \log_{10} \frac{a}{(a-x)}$  (d)  $K = \frac{1}{t} \frac{x}{a(a-x)}$
46. The first order rate constant for the decomposition of  $N_2O_5$  is  $6.2 \times 10^{-4} \text{ sec}^{-1}$ . The half life period for this decomposition in seconds is [MNR 1991; MP PET 1997; UPSEAT 2000]
- (a) 1117.7 (b) 111.7  
(c) 223.4 (d) 160.9
47. A first order reaction which is 30% complete in 30 minutes has a half-life period of [AIIMS 1998]
- (a) 24.2 min (b) 58.2 min  
(c) 102.2 min (d) 120.2 min
48. The order of a reaction which has the rate expression  
 $\frac{dc}{dt} = K[E]^{3/2}[D]^{3/2}$  is
- (a) 3/2 (b) 3  
(c) 2 (d) 0
49. The reaction  $2N_2O_5 \rightleftharpoons 2NO_2 + O_2$  follows first order kinetics. Hence, the molecularity of the reaction is
- (a) Unimolecular  
(b) Pseudo-unimolecular  
(c) Bimolecular  
(d) None of the above
50. A reaction involving two different reactants [KCET 1989; AIEEE 2005]
- (a) Can never be a second order reaction  
(b) Can never be a unimolecular reaction  
(c) Can never be a bimolecular reaction  
(d) Can never be a first order reaction
51. By "the overall order of a reaction", we mean
- (a) The number of concentration terms in the equation for the reaction  
(b) The sum of powers to which the concentration terms are raised in the velocity equation  
(c) The least number of molecules of the reactants needed for the reaction  
(d) The number of reactants which take part in the reaction
52. Catalyst decomposition of hydrogen peroxide is a ..... order reaction
- (a) First (b) Second  
(c) Third (d) Zero
53. The half life of a first order reaction is
- (a) Independent of the initial concentration of the reactant  
(b) Directly proportional to the initial concentration of the reactants  
(c) Inversely proportional to the initial concentration of the reactant  
(d) Directly proportional to the square of the initial concentration of the reactant
54. The decomposition of  $N_2O_5$  is a first order reaction represented by  $N_2O_5 \rightarrow N_2O_4 + \frac{1}{2}O_2$ . After 15 minutes the volume of  $O_2$  produced is 9 ml and at the end of the reaction 35 ml. The rate constant is equal to [MP PET 1995]
- [MP PMT 1999]
- (a)  $\frac{1}{15} \ln \frac{35}{44}$  (b)  $\frac{1}{15} \ln \frac{44}{26}$   
(c)  $\frac{1}{15} \ln \frac{44}{35}$  (d)  $\frac{1}{15} \ln \frac{35}{26}$
55. The unit of specific reaction rate constant for a first order (if the concentration expressed in molarity) would be [MNR 1988; UPSEAT 2000, 01]
- (a)  $\text{mole litre}^{-1} \text{ s}^{-1}$  (b)  $\text{mole litre}^{-1}$   
(c)  $\text{mole s}^{-1}$  (d)  $\text{s}^{-1}$
56. A first order reaction requires 30 minutes for 50% completion. The time required to complete the reaction by 75% will be
- (a) 45 minutes (b) 15 minutes  
(c) 60 minutes (d) None of these
57. Inversion of canesugar in dilute acid (conversion into glucose and fructose) is a
- (a) Unimolecular reaction  
(b) Bimolecular reaction  
(c) Trimolecular reaction  
(d) Pseudo-unimolecular reaction
58. The half life period of a first order reaction [AMU 2000]

- (a)  $\frac{0.693}{t}$  (b)  $\frac{0.693}{K}$   
(c)  $\frac{2.303}{t}$  (d)  $\frac{0.303}{K_1}$
59. The order of a reaction is said to be 2 with respect to a reactant  $X$ , when [DPMT 2000]  
(a) The rate of the reaction is proportional to  $[X]$   
(b) The rate of the reaction is proportional  $[X]^2$   
(c) Two molecules of  $X$  are present in the stoichiometric equation  
(d) The reaction occurs in two steps
60. Decay constant of a reaction is  $1.1 \times 10^{-9}$  /sec, then the half life of the reaction is  
(a)  $1.2 \times 10^8$  (b)  $6.3 \times 10^8$   
(c)  $3.3 \times 10^8$  (d)  $2.1 \times 10^8$
61. If the half life period of a reaction is inversely proportional to the initial concentration, the order of the reaction is  
(a) Zero (b) One  
(c) Two (d) Three
62. Which one of the following statements is wrong  
(a) Molecularity of a reaction is always a whole number  
(b) Order and molecularity of a reaction need not be same  
(c) Order of a reaction may be zero  
(d) Order of a reaction depends upon the mechanism of the reaction
63. The velocity constant of first order reaction is expressed in the units  
(a) Concentration per unit time  
(b) Time per unit concentration  
(c) Per unit time  
(d) Unit time per unit concentration
64. For reaction,  $A + B \rightarrow$  products, it is found that the rate of the reaction is proportional to the concentration of  $A$ , but it is independent of the concentration of  $B$ , then  
(a) The order of the reaction 2 and molecularity 1  
(b) Molecularity of the reaction is 2 but order is 1  
(c) Order is 2 and molecularity is 2  
(d) Order of the reaction is 2 but molecularity is 0
65. For a zero order reaction  
(a) The concentration of the reactant does not change during the reaction  
(b) The concentration change only when the temperature changes  
(c) The rate remains constant throughout  
(d) The rate of the reaction is proportional to the concentration
66. If ' $a$ ' is the initial concentration and ' $n$ ' is the order of the reaction and the half life period is ' $T$ ', then [MH CET 2000]  
(a)  $T \propto a^{n-1}$  (b)  $T \propto a^n$   
(c)  $T \propto \frac{1}{a^n}$  (d)  $T \propto \frac{1}{a^{n-1}}$
67. In presence of  $HCl$ , sucrose gets hydrolysed into glucose and fructose. The concentration of sucrose was found to reduce from 0.4  $M$  to 0.2  $M$  in 1 hour and 0.1  $M$  in 2 hours. The order of the reaction is  
(a) Zero (b) One  
(c) Two (d) None of these
68. The time of half change of a first order reaction in ..... initial concentration  
(a) Proportional to  
(b) Inversely proportional to  
(c) Independent of  
(d) Equal to
69. Half life period of a first order reaction is 138.6 minutes. The velocity constant of the reaction is  
(a)  $0.05 \text{ min}^{-1}$  (b)  $0.00005 \text{ min}^{-1}$   
(c)  $0.005 \text{ min}^{-1}$  (d)  $200 \text{ min}^{-1}$
70. An example of a pseudo –unimolecular reaction is  
(a) Dissociation of hydrogen iodide  
(b) Hydrolysis of methyl acetate in dilute solution  
(c) Dissociation of phosphorus pentachloride  
(d) Decomposition of hydrogen peroxide
71. About half life period of a first order reaction, which one of the following statements is generally false  
(a) It is independent of initial concentration  
(b) It is independent of temperature  
(c) It decreases with the introduction of a catalyst  
(d) It increases with increase of temperature
72. Decomposition of nitrogen pentoxide is known to be a first order reaction 75 percent of the oxide had decomposed in the first 24 minutes. At the end of an hour, after the start of the reaction, the amount of oxide left will be  
(a) Nil (b) About 1%  
(c) About 2% (d) About 3%
73. A reaction  $2A \rightarrow$  products is found to follow zero order kinetics, then  
(a)  $\frac{dx}{dt} = k[A]^2$  (b)  $\frac{dx}{dt} = k[A]^0$   
(c)  $\frac{dx}{dt} = k[A]$  (d)  $\frac{dx}{dt} = k[2A]$
74. The alkaline hydrolysis of ethyl acetate is represented by the equation  
 $CH_3COOC_2H_5 + NaOH \rightarrow CH_3COONa + C_2H_5OH$   
Experimentally it is found that for this reaction  
 $\frac{dx}{dt} = k[CH_3COOC_2H_5][NaOH]$   
Then the reaction is [JIPMER 1999]  
(a) Bimolecular and of first order  
(b) Bimolecular and of second order  
(c) Pseudo-bimolecular  
(d) Pseudo-unimolecular
75. For the reaction  $2HI \rightleftharpoons H_2 + I_2$ , the rate of the reaction is proportional to  $[HI]^2$ . This means that the reaction is [AMU 1985; MP PET 2000]  
(a) Unimolecular (b) Bimolecular  
(c) Of first order (d) Of second order
76. Inversion of sucrose is [AMU 1988; MP PET 2000]  
(a) Zero order reaction (b) First order reaction  
(c) Second order reaction (d) Third order reaction
77. The one which is unimolecular reaction is [MP PMT 1999; UPSEAT 2001]  
(a)  $2HI \rightarrow H_2 + I_2$  (b)  $N_2O_5 \rightarrow N_2O_4 + \frac{1}{2}O_2$

- (c)  $H_2 + Cl_2 \rightarrow 2HCl$  (d)  $PCl_3 + Cl_2 \rightarrow PCl_5$
78. Integrated velocity equation for first order reaction is  
 (a)  $[A]_o = [A]e^{-Kt}$  (b)  $K = [A]_o e^{-A/t}$   
 (c)  $Kt = 2.303 \log \frac{[A]_o}{[A]}$  (d)  $\log \frac{[A]_o}{[A]} = -2.303 Kt$
79. If the surface area of the reactants increases, then order of the reaction  
 (a) Increases  
 (b) Decreases  
 (c) Remain constant  
 (d) Sometimes increases and sometimes decreases
80. Half life period  $t_{1/2}$  for first order reaction is  
 (a)  $K$  (b)  $\frac{1.303 \log 2}{K}$   
 (c)  $\frac{2.303 \log 2}{K}$  (d)  $\frac{9}{K}$
81. Molecularity of reaction of inversion of sugar is  
 (a) 3 (b) 2  
 (c) 1 (d) 0
82. For any reaction, if we plot a graph between time ' $t$ ' and  $\log(a-x)$ , a simple line is obtained. The order of reaction is  
 (a) Zero (b) One  
 (c) Two (d) Three
83. Value of velocity constant for first order reaction is  $3.46 \times 10^{-3} \text{ min}^{-1}$ , the time for half change is  
 (a) 100 minutes (b) 400 minutes  
 (c) 200 minutes (d) 346 minutes
84. The unit of the velocity constant in case of zero order reaction is [CPMT 1994]  
 (a)  $\text{Conc.} \times \text{time}^{-1}$  (b)  $\text{Conc.}^{-1} \times \text{time}$   
 (c)  $\text{Conc.}^{-1} \times \text{time}^{-1}$  (d)  $\text{Conc.} \times (\text{time})^2$
85. For the reaction  $H_2(g) + Br_2(g) \rightarrow 2HBr(g)$ , the experimental data suggest, rate  $= K[H_2][Br_2]^{1/2}$ . The molecularity and order of the reaction are respectively [CPMT 1988; MP PET 1993]  
 (a)  $2, \frac{3}{2}$  (b)  $\frac{3}{2}, \frac{3}{2}$   
 (c) 1, 1 (d)  $1, \frac{1}{2}$
86. The incorrect order indicated against the rate of reaction  $A + B \xrightarrow{K} C$  is [BHU 1990]  

Rate	Order
(a) $\frac{d[C]}{dt} = K[A]$	1
(b) $\frac{d[C]}{dt} = K[A][B]$	2
(c) $\frac{-d[A]}{dt} = K[A][B]^0$	2
- (d)  $\frac{-d[A]}{dt} K[A]$  1
87. Which of the following statements regarding the molecularity of a reaction is wrong [CBSE PMT 1989]  
 (a) It is the number of molecules of the reactants taking part in a single step chemical reaction  
 (b) It is calculated from the reaction mechanism  
 (c) It may be either a whole number or fractional  
 (d) It depends on the rate determining step in the reaction
88. Diazonium salt decomposes as  
 $C_6H_5N_2^+Cl^- \rightarrow C_6H_5Cl + N_2$   
 At  $0^\circ C$ , the evolution of  $N_2$  becomes two times faster when the initial concentration of the salt is doubled. Therefore, it is [MNR 1994; UPSEAT 2]  
 (a) A first order reaction  
 (b) A second order reaction  
 (c) Independent of the initial concentration of the salt  
 (d) A zero order reaction
89. In the reaction  $A + B \rightarrow \text{Products}$ , if  $B$  is taken in excess, then it is an example of [EAMCET 1992]  
 (a) Second order reaction  
 (b) Zero order reaction  
 (c) Pseudounimolecular reaction  
 (d) First order reaction
90. The half life of a first order reaction is 69.35 sec. The value of the rate constant of the reaction is [CBSE PMT 1990]  
 (a)  $1.0 s^{-1}$  (b)  $0.1 s^{-1}$   
 (c)  $0.01 s^{-1}$  (d)  $0.001 s^{-1}$
91. The half life for the reaction  $N_2O_5 \rightleftharpoons 2NO_2 + \frac{1}{2}O_2$  in 24 hrs at  $30^\circ C$ . Starting with 10 g of  $N_2O_5$  how many grams of  $N_2O_5$  will remain after a period of 96 hours [KCET 1992]  
 (a) 1.25 g (b) 0.63 g  
 (c) 1.77 g (d) 0.5 g
92. The half life of a first order reaction is 10 minutes. If initial amount is  $0.08 \text{ mol/litre}$  and concentration at some instant is  $0.01 \text{ mol/litre}$ , then  $t =$  [Roorkee 1990]  
 (a) 10 minutes (b) 30 minutes  
 (c) 20 minutes (d) 40 minutes
93. Half life period of second order reaction is [MP PMT 1994]  
 (a) Proportional to the initial concentration of reactants  
 (b) Independent of the initial concentration of reactants  
 (c) Inversely proportional to initial concentration of reactants  
 (d) Inversely proportional to square of initial concentration of reactants
94. The reaction  $2H_2O_2 \rightarrow 2H_2O + O_2$  is a [Manipal MEE 1995]  
 (a) Zero order reaction (b) First order reaction  
 (c) Second order reaction (d) Third order reaction
95. In a reaction involving hydrolysis of an organic chloride in presence of large excess of water  
 $RCI + H_2O \rightarrow ROH + HCl$  [MP PET 1995]  
 (a) Molecularity is 2, order of reaction is also 2

- (b) Molecularity is 2, order of reaction is 1  
(c) Molecularity is 1, order of reaction is 2  
(d) Molecularity is 1, order of reaction is also 1
96. The thermal decomposition of a compound is of first order. If a sample of the compound decomposes 50% in 120 minutes, in what time will it undergo 90% decomposition  
[MP PET 1996]  
(a) Nearly 240 minutes (b) Nearly 480 minutes  
(c) Nearly 450 minutes (d) Nearly 400 minutes
97. The order of a reaction with rate equals  $kC_A^{3/2}C_B^{-1/2}$  is  
[MP PET 1996, 2001]  
(a) 2 (b) 1  
(c)  $-\frac{1}{2}$  (d)  $\frac{3}{2}$
98. If the rate expression for a chemical reaction is given by Rate  $= k[A]^m[B]^n$   
[MP PMT 1996]  
(a) The order of the reaction is  $m$   
(b) The order of the reaction is  $n$   
(c) The order of the reaction is  $m+n$   
(d) The order of the reaction is  $m-n$
99. The half-life period of a first order reaction is 100 sec. The rate constant of the reaction is  
[MP PMT 1997; MP PET 2001]  
(a)  $6.93 \times 10^{-3} \text{ sec}^{-1}$  (b)  $6.93 \times 10^{-4} \text{ sec}^{-1}$   
(c)  $0.693 \text{ sec}^{-1}$  (d)  $69.3 \text{ sec}^{-1}$
100. For the first order reaction with rate constant  $k$ , which expression gives the half-life period? (Initial concentration  $= a$ )  
[MP PET 1998]  
(a)  $\frac{1^2}{k}$  (b)  $\frac{1}{ka}$   
(c)  $\frac{0.693}{k}$  (d)  $\frac{3}{2ka^2}$
101. The rate constant of a first order reaction whose half-life is 480 seconds, is  
[MP PET 1999]  
(a)  $2.88 \times 10^{-3} \text{ sec}^{-1}$  (b)  $1.44 \times 10^{-3} \text{ sec}^{-1}$   
(c)  $1.44 \text{ sec}^{-1}$  (d)  $0.72 \times 10^{-3} \text{ sec}^{-1}$
102. The conversion of  $A \rightarrow B$  follows second order kinetics. Doubling the concentration of  $A$  will increase the rate of formation of  $B$  by a factor  
[MP PET 1999; DCE 1999; KCET 2001; BCECE 2005]  
(a) 1/4 (b) 2  
(c) 1/2 (d) 4
103. The reaction  $2FeCl_3 + SnCl_2 \rightarrow 2FeCl_2 + SnCl_4$  is an example of  
[CBSE PMT 1996; MP PET 1999]  
(a) First order reaction  
(b) Second order reaction  
(c) Third order reaction  
(d) None of these
104. If reaction between  $A$  and  $B$  to give  $C$  shows first order kinetics in  $A$  and second order in  $B$ , the rate equation can be written as  
[MP PET 1999]  
(a) Rate  $= k[A][B]^{1/2}$  (b) Rate  $= k[A]^{1/2}[B]$   
(c) Rate  $= k[A][B]^2$  (d) Rate  $= k[A]^2[B]$
105. For a first order reaction, the half-life period is independent of  
[CBSE PMT 1999]  
(a) Initial concentration  
(b) Cube root of initial concentration  
(c) First power of final concentration  
(d) Square root of final concentration
106. Order of a reaction can have  
[DPMT 1996]  
(a) +ve values (b) Whole number values  
(c) Fractional values (d) All of the above
107. The order of the reaction occurring by following mechanism should be  
[JIPMER 1997]  
(i)  $A_2 \rightarrow A + A$  (fast)  
(ii)  $A + B_2 \rightarrow AB + B$  (slow)  
(iii)  $A + B \rightarrow$  (fast)  
(a)  $1\frac{1}{2}$  (b)  $3\frac{1}{2}$   
(c) 2 (d) None of these
108. For the reaction  $A \rightarrow B$ , the rate law expression is: Rate  $= k[A]$   
Which of the following statements is incorrect  
[Pb. PMT 1998]  
(a) The reaction is said to follow first order kinetics  
(b) The half life of the reaction will depend on the initial concentration of the reactant  
(c)  $k$  is constant for the reaction at a constant temperature  
(d) The rate law provides a simple way of predicting the concentration of reactants and products at any time after the start of the reaction
109. If initial concentration is reduced to its 1/4th in a zero order reaction, the time taken for half of the reaction to complete  
[BHU 1998]  
(a) Remains same (b) Becomes 4 times  
(c) Becomes one-fourth (d) Doubles
110. For a reaction whose rate expression is:  
Rate  $= k[A]^{1/2}[B]^{3/2}$ , the order would be  
[Pune CET 1998]  
(a) 1.5 (b) 2  
(c) 3 (d) 1
111. For the reaction  $A \rightarrow B$ , the rate increases by a factor of 2.25 when the concentration of  $A$  is increased by 1.5. What is the order of the reaction  
[KCET 1998]  
(a) 3 (b) 0  
(c) 2 (d) 1
112. For the reaction  $H_2 + Cl_2 \xrightarrow{\text{Sunlight}} 2HCl$  taking place on water, the order of reaction is  
[KCET 1998; AIIMS 2002; Pb. PMT 2002]  
(a) 1 (b) 2  
(c) 3 (d) 0
113. For a first order reaction, rate constant is  $0.6932 \text{ hr}^{-1}$ , then half-life for the reaction is  
[Bihar MEE 1997]  
(a) 0.01 hr (b) 1 hr  
(c) 2 hr (d) 10 hr  
(e) 0.1 hr
114. The rate constant of a reaction is  $0.69 \times 10^{-1} \text{ min}^{-1}$  and the initial concentration is  $0.2 \text{ mol l}^{-1}$ . The half-life period is  
[AIIMS 1998]



- (a) 400 sec (b) 600 sec  
(c) 800 sec (d) 1200 sec
115. The rate constant of a first order reaction is  $3 \times 10^{-6}$  per second. If the initial concentration is 0.10 M, the initial rate of reaction is  
[AFMC 1999 Pb. PMT 1999, 2000; BHU 1999; AIIMS 1999; KCET 2000; DCE 2004]
- (a)  $3 \times 10^{-5} \text{ ms}^{-1}$  (b)  $3 \times 10^{-6} \text{ ms}^{-1}$   
(c)  $3 \times 10^{-8} \text{ ms}^{-1}$  (d)  $3 \times 10^{-7} \text{ ms}^{-1}$
116. Certain bimolecular reactions which follow the first order kinetics are called [KCET (Med.) 1999]
- (a) First order reactions  
(b) Unimolecular reactions  
(c) Bimolecular reactions  
(d) Pseudounimolecular reactions
117. The rate law of the reaction  $A + 2B \rightarrow \text{Product}$  is given by  $\frac{d[B]}{dt} = k[B]^2$ . If A is taken in excess, the order of the reaction will be [AMU (Engg.) 1999]
- (a) 1 (b) 2  
(c) 3 (d) 0
118. For a first order reaction  $A \rightarrow \text{product}$ , the rate of reaction at  $[A] = 0.2 \text{ mol l}^{-1}$  is  $1.0 \times 10^{-2} \text{ mol l}^{-1} \text{ min}^{-1}$ . The half life period for the reaction is [Roorkee 1999]
- (a) 832 s (b) 440 s  
(c) 416 s (d) 13.86 s
119. For the reaction  $A + B \rightarrow \text{products}$ , doubling the concentration of A the rate of the reaction is doubled, but on doubling the concentration of B rate remains unaltered. The over all order of the reaction is [JIPMER 1999]
- (a) 1 (b) 0  
(c) 2 (d) 3
120. Which among the following is a false statement [KCET 1999]
- (a) Half life of a third order reaction is inversely proportional to the square of initial concentration of the reactant.  
(b) Molecularity of a reaction may be zero or fractional  
(c) For a first order reaction  $t_{1/2} = \frac{0.693}{K}$   
(d) Rate of zero order reaction is independent of initial concentration of reactant
121. After how many seconds will the concentration of the reactants in a first order reaction be halved, if the decay constant is  $1.155 \times 10^{-3} \text{ sec}^{-1}$  [CBSE PMT 2000]
- (a) 100 sec (b) 200 sec  
(c) 400 sec (d) 600 sec
122. What is the order of a reaction which has a rate expression rate  $= K[A]^{3/2}[B]^{-1}$  [DCE 2000]
- (a) 3/2 (b) 1/2  
(c) 0 (d) None of these
123. Which of the following expression is correct for first order reaction? (CO) refers to initial concentration of reactant [DCE 2000]
- (a)  $t_{1/2} \propto CO$  (b)  $t_{1/2} \propto CO^{-1}$   
(c)  $t_{1/2} \propto CO^{-2}$  (d)  $t_{1/2} \propto CO^0$
124. For a reaction  $2NO(g) + Cl_2(g) \rightleftharpoons 2NOCl(g)$ . When concentration of  $Cl_2$  is doubled, the rate of reaction becomes two times of the original. When the concentration of NO is doubled the rate becomes four times. What is the order of the reaction
- (a) 1 (b) 2  
(c) 3 (d) 4
125. The rate constant for a second order reaction is  $8 \times 10^{-5} \text{ M}^{-1} \text{ min}^{-1}$ . How long will it take a 1M solution to be reduced to 0.5 M [MH CET 2001]
- (a)  $8 \times 10^{-5} \text{ min}$  (b)  $8.665 \times 10^3 \text{ min}$   
(c)  $4 \times 10^{-5} \text{ min}$  (d)  $1.25 \times 10^4 \text{ min}$
126. The rate for a first order reaction is  $0.6932 \times 10^{-2} \text{ mol l}^{-1} \text{ min}^{-1}$  and the initial concentration of the reactants is 1M,  $T_{1/2}$  is equal to [JIPMER (Med.) 2001]
- (a) 6.932 min (b) 100 min  
(c)  $0.6932 \times 10^{-3} \text{ min}$  (d)  $0.6932 \times 10^{-2} \text{ min}$
127. For a given reaction  $t_{1/2} = \frac{1}{Ka}$ . The order of the reaction is [KCET 2001]
- (a) 1 (b) 0  
(c) 3 (d) 2
128. 75% of a first order reaction is completed in 30 minutes. What is the time required for 93.75% of the reaction (in minutes)
- (a) 45 (b) 120  
(c) 90 (d) 60
129. A First order reaction is half completed in 45 minutes. How long does it need 99.9% of the reaction to be completed [AIIMS 2001]
- (a) 5 hours (b) 7.5 hours  
(c) 10 hours (d) 20 hours
130. A substance 'A' decomposes by a first order reaction starting initially with  $[A] = 2.00 \text{ M}$  and after 200 min  $[A] = 0.15 \text{ M}$ . For this reaction what is the value of  $k$  [AIIMS 2001]
- (a)  $1.29 \times 10^{-2} \text{ min}^{-1}$  (b)  $2.29 \times 10^{-2} \text{ min}^{-1}$   
(c)  $3.29 \times 10^{-2} \text{ min}^{-1}$  (d)  $4.40 \times 10^{-2} \text{ min}^{-1}$
131. Which of the following statements about zero order reaction is not true [DCE 2001]
- (a) Its unit is  $\text{sec}^{-1}$   
(b) The graph between log (reactant) versus rate of reaction is a straight line  
(c) The rate of reaction increases with the decrease in concentration of reactants  
(d) Rate of reaction is independent of concentration of reactants
132. The given reaction  $2NO + O_2 \rightarrow 2NO_2$  is an example of [CBSE PMT 2001; JIPMER 2002]
- (a) First order reaction (b) Second order reaction  
(c) Third order reaction (d) None of these
133. Order of a reaction is decided by [KCET 2002]
- (a) Pressure  
(b) Temperature

- (c) Molecularity  
(d) Relative concentration of reactants
134. From the following which is a second order reaction [AMU 2002]
- (a)  $K = 5.47 \times 10^{-4} \text{ sec}^{-1}$   
(b)  $K = 3.9 \times 10^{-3} \text{ mole lit sec}^{-1}$   
(c)  $K = 3.94 \times 10^{-4} \text{ lit mole}^{-1} \text{ sec}^{-1}$   
(d)  $K = 3.98 \times 10^{-5} \text{ lit mole}^{-2} \text{ sec}^{-1}$
135. For the reaction  $A + 2B \rightarrow C$ , rate is given by  $R = [A][B]^2$  then the order of the reaction is [AIEEE 2002]
- (a) 3 (b) 6  
(c) 5 (d) 7
136. Units of rate constant of first and zero order reactions in terms of molarity  $M$  unit are respectively [AIEEE 2002]
- (a)  $\text{sec}^{-1}$ ,  $M \text{ sec}^{-1}$  (b)  $\text{sec}^{-1}$ ,  $M$   
(c)  $M \text{ sec}^{-1}$ ,  $\text{sec}^{-1}$  (d)  $M, \text{sec}^{-1}$
137. The reaction  $2N_2O_5 \rightleftharpoons 2N_2O_4 + O_2$  is [MP PMT 2002]
- (a) Bimolecular and second order  
(b) Unimolecular and first order  
(c) Bimolecular and first order  
(d) Bimolecular and zero order
138. The half-life period for a first order reaction is 693 seconds. The rate constants for this reaction would be [MP PET 2002]
- (a)  $0.1 \text{ sec}^{-1}$  (b)  $0.01 \text{ sec}^{-1}$   
(c)  $0.001 \text{ sec}^{-1}$  (d)  $0.0001 \text{ sec}^{-1}$
139. For an elementary reaction,  $2A + B \rightarrow C + D$  the molecularity is
- (a) Zero (b) One  
(c) Two (d) Three
140. If the order of the reaction  $x + y \xrightarrow{h\nu} x + y$  is zero, it means that the rate of [Kurukshetra CEE 2002]
- (a) Reaction is independent of temperature  
(b) Formation of activated complex is zero  
(c) Reaction is independent of the concentration of reacting species  
(d) Decomposition of activated complex is zero
141. For a first order reaction velocity constant,  $K = 10^{-3} \text{ s}^{-1}$ . Two third life for it would be [MP PET 2001; UPSEAT 2003]
- (a) 1100 s (b) 2200 s  
(c) 3300 s (d) 4400 s
142. In a reaction, the concentration of reactant is increased two times and three times then the increases in rate of reaction were four times and nine times respectively, order of reaction is
- (a) Zero (b) 1  
(c) 2 (d) 3
143. For a chemical reaction....can never be a fraction [EAMCET 2003]
- (a) Order (b) Half-life  
(c) Molecularity (d) Rate constant
144. 75% of a first order reaction was completed in 32 minutes when was 50% of the reaction completed [AMU 1999; Kerala (Med.) 2003]
- (a) 16 min. (b) 24 min.  
(c) 8 min. (d) 4 min.
145. The decomposition of  $N_2O_5$  occurs as,  $2N_2O_5 \rightarrow 4NO_2 + O_2$ , and follows 1<sup>st</sup> order kinetics, hence [BVP 2003]
- (a) The reaction is unimolecular  
(b) The reaction is bimolecular  
(c)  $T_{1/2} \propto a^0$   
(d) None of these
146. Which equation is correct for first order reactions [MP PMT 2003]
- (a)  $t_{1/2} \propto C^{-1}$  (b)  $t_{1/2} \propto C$   
(c)  $t_{1/2} \propto C^0$  (d)  $t_{1/2} \propto C^{1/2}$
147. For the reaction system  $2NO(g) + O_2(g) \rightarrow 2NO_2(g)$  volume is suddenly produced to half its value by increasing the pressure on it. If the reaction is of first order with respect to  $O_2$  and second order with respect to  $NO$ , the rate of reaction will [AIEEE 2003]
- (a) Diminish to one fourth of its initial value  
(b) Diminish to one eighth of its initial value  
(c) Increase to eight times of its initial value  
(d) Increase to four times of its initial value
148. If the rate of the reaction is equal to the rate constant, the order of the reaction is [CBSE PMT 2003]
- (a) 3 (b) 0  
(c) 1 (d) 2
149. The reaction  $A \rightarrow B$  follows first order kinetics. The time taken for 0.8 mole of  $A$  to produce 0.6 mole of  $B$  is 1 hour. What is the time taken for conversion of 0.9 mole of  $A$  to produce 0.675 mole of  $B$  [Kurukshetra CEE 2002]
- (a) 2 hours (b) 1 hour  
(c) 0.5 hour (d) 0.25 hour
150. The unit of velocity constant in case of zero order reaction is [MP PMT 2004]
- (a)  $\text{Concentration} \times \text{Time}^{-1}$  (b)  $\text{Concentration}^{-1} \times \text{Time}^{-1}$   
(c)  $\text{Concentration} \times \text{Time}^2$  (d)  $\text{Concentration}^{-1} \times \text{Time}$
151. Which one of the following is wrongly matched [KCET 2004]
- (a) Saponification of  $CH_3COOC_2H_5$  – Second order reaction  
(b) Hydrolysis of  $CH_3COOCH_3$  – Pseudo uni-molecular reaction  
(c) Decomposition of  $H_2O_2$  – First order reaction  
(d) Combination of  $H_2$  and  $Br_2$  to give  $HBr$  – Zero order reaction
- [UPSEAT 2003]
152. Which of the following is an example of pseudo unimolecular reaction [Pb. CET 2001]
- (a)  $CH_3COOCH_3 + H_2O \xrightarrow{H^+} CH_3COOH + CH_3OH$   
(b)  $CH_3COOCH_3 + H_2O \xrightarrow{OH^-} CH_3COOH + CH_3OH$   
(c)  $2FeCl_3 + SnCl_2 \rightarrow SnCl_4 + 2FeCl_2$   
(d)  $NaOH + HCl \rightarrow NaCl + H_2O$
153. Hydrolysis of DDT is a first order reaction, its half life is 10 years. Time to hydrolyse 10 g DDT to half is [BVP 2004]

- (a) 100 years (b) 50 years  
(c) 5 years (d) 10 years
154. In a first order reaction, the concentration of the reactant, decreases from 0.8 M to 0.4 M in 15 minutes. The time taken for the concentration to change from 0.1 M to 0.025 M is [AIEEE 2004]  
(a) 7.5 minutes (b) 15 minutes  
(c) 30 minutes (d) 60 minutes
155. In the first order reaction, the concentration of the reactant is reduced to 25% in one hour. The half life period of the reaction is [DCE 2004]  
(a) 2 hr (b) 4 hr  
(c) 1/2 hr (d) 1/4 hr
156. For a reaction,  $X(g) \rightarrow Y(g) + Z(g)$  the half life period is 10 min. In What period of time would the concentration of X be reduced to 10% of original concentration [DCE 2004]  
(a) 20 min (b) 33 min  
(c) 15 min (d) 25 min
157. A first order reaction with respect to the reactant A has a rate constant of  $6 \text{ sec}^{-1}$ . If we start with  $[A] = 0.5 \text{ mol/litre}$ , then in what time the concentration of A becomes 0.05 mol/litre [DCE 2004]  
(a) 0.384 sec (b) 0.214 sec  
(c) 3.84 sec (d) 0.402 sec
158. Order of radioactive disintegration reaction is [JEE Orissa 2004]  
(a) Zero (b) First  
(c) Second (d) Third
159. The rate of a first order reaction is  $1.5 \times 10^{-2} \text{ mol L}^{-1} \text{ min}^{-1}$  at 0.5 M concentration of the reactant. The half life of the reaction is [CBSE PMT 2004]  
(a) 8.73 min (b) 7.53 min  
(c) 0.383 min (d) 23.1 min
160. A first order reaction was started with a decimolar solution of the reactant, 8 minutes and 20 seconds later its concentration was found to be  $M/100$ . So the rate of the reaction is [Kerala PMT 2004]  
(a)  $2.303 \times 10^{-5} \text{ sec}^{-1}$  (b)  $2.303 \times 10^{-4} \text{ sec}^{-1}$   
(c)  $4.606 \times 10^{-3} \text{ sec}^{-1}$  (d)  $2.606 \times 10^{-5} \text{ sec}^{-1}$   
(e)  $2.603 \times 10^{-4} \text{ sec}^{-1}$
161. Which is correct about zero order reaction [JEE Orissa 2004]  
(a) Rate of reaction depends on decay constant  
(b) Rate of reaction is independent of concentration  
(c) Unit of rate constant is concentration  
(d) Unit of rate constant is concentration  $\cdot$  time
162. Decay of  ${}_{92}\text{U}^{235}$  is .....order reaction [JEE Orissa 2004]  
(a) Zero (b) First  
(c) Second (d) Third
163. The half-life of 2 sample are 0.1 and 0.4 seconds. Their respective concentration are 200 and 50 respectively. What is the order of the reaction [JEE Orissa 2004]  
(a) 0 (b) 2  
(c) 1 (d) 4
164. The following statements(s) is(are) correct [IIT 1999]  
(a) A plot of  $\log K_p$  versus  $1/T$  is linear  
(b) A plot of  $\log[X]$  versus time is linear for a first order reaction  $X \rightarrow P$   
(c) A plot of  $\log P$  versus  $1/T$  is linear at constant volume  
(d) A plot of  $P$  versus  $1/V$  is linear at constant temperature
165. For a first order reaction [IIT 1998]  
(a) The degree of dissociation is equal to  $(1 - e^{-kt})$   
(b) A plot of reciprocal concentration of the reactant vs time gives a straight line  
(c) The time taken for the completion of 75% reaction is thrice the  $t_{1/2}$  of the reaction  
(d) The pre-exponential factor in the Arrhenius equation has the dimension of time  $T^{-1}$
166. For reaction  $a A \rightarrow x P$ , when  $[A] = 2.2 \text{ mM}$ , the rate was found to be  $2.4 \text{ mM s}^{-1}$ . On reducing concentration of A to half, the rate changes to  $0.6 \text{ mM s}^{-1}$ . The order of reaction with respect to A is [AIIMS 2005]  
(a) 1.5 (b) 2.0  
(c) 2.5 (d) 3.0
167. Which one of the following statement for order of reaction is not correct [IIT 2005]  
(a) Order can be determined experimentally.  
(b) Order of reaction is equal to sum of the powers of concentration terms in differential rate law.  
(c) It is not affected with the stoichiometric coefficient of the reactants.  
(d) Order cannot be fractional.
168.  $t_{1/4}$  can be taken as the time taken for the concentration of a reactant to drop to  $\frac{3}{4}$  of its initial value. If the rate constant for a first order reaction is  $K$ , the  $t_{1/4}$  can be written as [AIEEE 2005]  
(a)  $0.10 / K$  (b)  $0.29 / K$   
(c)  $0.69 / K$  (d)  $0.75 / K$
169. For a first order reaction  $A \rightarrow B$  the reaction rate at reactant concentration of 0.01M is found to be  $2.0 \times 10^{-5} \text{ mol L}^{-1} \text{ s}^{-1}$ . The half life period of the reaction is [CBSE PMT 2005]  
(a) 220 s (b) 30 s  
(c) 300 s (d) 347 s
170. The rate of reaction between two reactants A and B decreases by a factor of 4 if the concentration of reactant B is doubled. The order of this reaction with respect to reactant B is  
(a) -1 (b) -2  
(c) 1 (d) 2
171. If a substance with half life 3 days is taken at other place in 12 days. What amount of substance is left now [AFMC 2005]  
(a) 1/4 (b) 1/8  
(c) 1/16 (d) 1/32
172. The half-life of a first order reaction having rate constant  $K = 1.7 \times 10^3 \text{ s}$  is [BHU 2005]  
(a) 12.1 h (b) 9.7 h  
(c) 11.3 h (d) 1.8 h
173. For the reaction  $A + B \rightarrow C$ , it is found that doubling the concentration of A increases the rate by 4 times, and doubling the concentration of B doubles the reaction rate. What is the overall order of the reaction. [KCET 2005]

- (a) 4 (b)  $3/2$   
(c) 3 (d) 1

174. Which of the following reactions end in finite time

[DPMT 2005]

- (a) 0 order (b) 1st order  
(c) 2nd order (d) 3rd order

### Collision theory, Energy of activation and Arrhenius equation

1. A large increase in the rate of a reaction for a rise in temperature is due to [EAMCET 1980; MP PET 1995]

- (a) The decrease in the number of collisions  
(b) The increase in the number of activated molecules  
(c) The shortening of the mean free path  
(d) The lowering of the activation energy

2. Which of the following statements is not true according to collision theory of reaction rates

- (a) Collision of molecules is a precondition for any reaction to occur  
(b) All collisions result in the formation of the products  
(c) Only activated collisions result in the formation of the products  
(d) Molecules which have acquired the energy of activation can collide effectively

3. According to the collision theory of chemical reactions

- (a) A chemical reaction occurs with every molecular collision  
(b) Rate is directly proportional to the number of collisions per second  
(c) Reactions in the gas phase are always of zero order  
(d) Reaction rates are of the order of molecular speeds

4. According to the collision theory of reaction rates, rate of reaction increases with temperature due to

- (a) Greater number of collisions  
(b) Greater velocity of the reacting molecules  
(c) Greater number of molecules have activation energy  
(d) None of the above

5. The reaction rate at a given temperature becomes slower, then

[MP PMT 1993; DPMT 2000]

- (a) The free energy of activation is higher  
(b) The free energy of activation is lower  
(c) The entropy changes  
(d) The initial concentration of the reactants remains constant

6. A rise in temperature increases the velocity of a reaction. It is because it results in

- (a) An increased number of molecular collisions  
(b) An increased momentum of colliding molecules  
(c) An increase in the activation energy  
(d) A decrease in the activation energy

7. The number of collisions depend upon

- (a) Pressure (b) Concentration  
(c) Temperature (d) All the above

8. If  $E_f$  and  $E_r$  are the activation energies of forward and reverse reactions and the reaction is known to be exothermic, then

(a)  $E_f > E_r$

(b)  $E_f < E_r$

(c)  $E_f = E_r$

(d) No relation can be given between  $E_f$  and  $E_r$  as data are not sufficient

9. According to Arrhenius theory, the activation energy is

- (a) The energy it should possess so that it can enter into an effective collision  
(b) The energy which the molecule should possess in order to undergo reaction  
(c) The energy it has to acquire further so that it can enter into a effective collision  
(d) The energy gained by the molecules on colliding with another molecule

10. The energy of activation is

- (a) The energy associated with the activated molecules  
(b) Threshold energy – energy of normal molecules  
(c) Threshold energy + energy of normal molecules  
(d) Energy of products – energy of reactants

11. Which one of the following does not represent Arrhenius equation

(a)  $k = Ae^{-E/RT}$

(b)  $\log_e k = \log_e A - \frac{E}{RT}$

(c)  $\log_{10} k = \log_{10} A - \frac{E}{2.303 RT}$

(d)  $k = AE^{-RT}$

12. On increasing the temperature, the rate of the reaction increases because of [MP PMT 1997]

- (a) Decrease in the number of collisions  
(b) Decrease in the energy of activation  
(c) Decrease in the number of activated molecules  
(d) Increase in the number of effective collisions

13. Energy of activation of a reactant is reduced by

- (a) Increased temperature (b) Reduced temperature  
(c) Reduced pressure (d) Increased pressure

14. The minimum energy a molecule should possess in order to enter into a fruitful collision is known as

[Kurukshetra CEE 2002]

- (a) Reaction energy (b) Collision energy  
(c) Activation energy (d) Threshold energy

15. Activation energy is


- (a) The amount of energy to be added to the actual energy of a molecule so that the threshold energy is reached  
(b) The amount of energy the molecule must contain so that it reacts  
(c) The energy which a molecule should have in order to enter into an effective collision  
(d) The average kinetic energy of the molecule

16. The reason for almost doubling the rate of reaction on increasing the temperature of the reaction system by  $10^\circ\text{C}$  is

[J & K 2005]

- (a) The value of threshold energy increases  
(b) Collision frequency increases

- (c) The fraction of the molecule having energy equal to threshold energy or more increases  
(d) Activation energy decreases
17. The activation energy for a simple chemical reaction  $A \rightarrow B$  is  $E_a$  in forward direction. The activation energy for reverse reaction
- (a) Is always double of  $E_a$   
(b) Is negative of  $E_a$   
(c) Is always less than  $E_a$   
(d) Can be less than or more than  $E_a$
18. Arrhenius equation is
- (a)  $\frac{d \ln K}{dT} = \Delta E^* / RT$  (b)  $\frac{d \ln K}{dT} = \Delta E^* / RT^2$   
(c)  $\frac{d \ln K}{dT} = -\Delta E^* / RT^2$  (d)  $\frac{d \ln K}{dT} = -\Delta E^* / RT$
19. Activation energy of any reaction depends on
- (a) Temperature  
(b) Nature of reactants  
(c) Number of collisions per unit time  
(d) Concentration of reactants
20. Relation between rate constant and temperature by Arrhenius equation is
- (a)  $\log_e A = \log_e K + \frac{E_a}{RT}$  (b)  $\log K = A \frac{E_a}{RT}$   
(c)  $\log_e K = \log_e A - \frac{E_a}{RT^2}$  (d)  $\log A = RT \ln E_a - \ln K$
21. An endothermic reaction  $A \rightarrow B$  has an activation energy  $15 \text{ kcal/mole}$  and energy of reaction  $5 \text{ kcal/mole}$ . The activation energy of the reaction  $B \rightarrow A$  is [Pb. CET 1985]
- (a)  $20 \text{ kcal/mole}$  (b)  $15 \text{ kcal/mole}$   
(c)  $10 \text{ kcal/mole}$  (d) None of these
22. Which of the following plots is in accordance with the Arrhenius equation




23. The Arrhenius Equation expressing the effect of temperature on the rate constant of a reaction is [MP PET 1997]
- (a)  $k = e^{-E_a/RT}$  (b)  $k = E_a / RT$   
(c)  $k = \log_e \frac{E_a}{RT}$  (d)  $k = A e^{-E_a/RT}$
24. For a reaction, activation energy ( $E_a$ ) = 0 and rate constant ( $K$ ) =  $3.2 \times 10^6 \text{ s}^{-1}$  at 300 K. What is the value of the rate constant at 310 K [KCET (Med.) 1999]
- (a)  $3.2 \times 10^{-12} \text{ s}^{-1}$  (b)  $3.2 \times 10^6 \text{ s}^{-1}$   
(c)  $6.4 \times 10^{12} \text{ s}^{-1}$  (d)  $6.4 \times 10^6 \text{ s}^{-1}$
25. Activation energy is given by the formula [DCE 1999]
- (a)  $\log \frac{K_2}{K_1} = \frac{E_a}{2.303R} \left[ \frac{T_2 - T_1}{T_1 T_2} \right]$

$$(b) \log \frac{K_1}{K_2} = -\frac{E_a}{2.303R} \left[ \frac{T_2 - T_1}{T_1 T_2} \right]$$

$$(c) \log \frac{K_1}{K_2} = -\frac{E_a}{2.303R} \left[ \frac{T_1 - T_2}{T_1 T_2} \right]$$

(d) None of these

26. A reaction having equal activation energies for forward and reverse reaction has [MP PMT 2002]
- (a)  $\Delta H = 0$  (b)  $\Delta S = 0$   
(c) Zero order (d) None of these
27. Collision theory is applicable to [MP PMT 2002]
- (a) First order reactions (b) Zero order reactions  
(c) Bimolecular reactions (d) Intra molecular reactions
28. A graph plotted between  $\log K$  vs  $1/T$  for calculating activation energy is shown by [MP PET 2002]



29. The rate constant of a reaction at temperature 200 K is 10 times less than the rate constant at 400 K. What is the activation energy ( $E_a$ ) of the reaction ( $R$  = gas constant) [EAMCET 2003]
- (a)  $1842.4 R$  (b)  $921.2 R$   
(c)  $460.6 R$  (d)  $230.3 R$
30. In respect of the equation  $k = A e^{-E_a/RT}$  in chemical kinetics, which one of the following statement is correct [AIEEE 2003]
- (a)  $k$  is equilibrium constant  
(b)  $A$  is adsorption factor  
(c)  $E_a$  is energy of activation  
(d)  $R$  is Rydberg's constant

31. The rate constant is doubled when temperature increases from  $27^\circ \text{C}$  to  $37^\circ \text{C}$ . Activation energy in  $\text{kJ}$  is [JEE Orissa 2004]


- (a) 34 (b) 54  
(c) 100 (d) 50

32. The activation energy of a reaction is zero. The rate constant of this reaction
- (a) Increases with increase of temperature  
(b) Decreases with an increase of temperature  
(c) Decreases with decrease of temperature  
(d) Is independent of temperature

33. The rate constant is given by the equation  $k = p z e^{-E/RT}$ . Which factor should register a decrease for the reaction to proceed more rapidly [MP PET/PMT 1998]

- (a)  $T$  (b)  $Z$   
(c)  $E$  (d)  $p$

34. An endothermic reaction with high activation energy for the forward reaction is given by the diagram: [AIIMS 2005]



35. Consider an endothermic reaction  $X \rightarrow Y$  with the activation energies  $E_b$  and  $E_f$  for the backward and forward reactions, respectively, in general [AIEEE 2005]

- (a)  $E_b < E_f$   
 (b)  $E_b > E_f$   
 (c)  $E_b = E_f$   
 (d) There is no definite relation between  $E_b$  and  $E_f$

36. Temperature dependent equation can be written as

[Orissa JEE 2005]

- (a)  $\ln k = \ln A - e^{E_a/RT}$  (b)  $\ln k = \ln A + e^{E_a/RT}$   
 (c)  $\ln k = \ln A - e^{RT/E_a}$  (d) All of these

### Photochemical reactions

- The formation of starch in plants results from
  - Photosynthesis
  - Photolysis
  - Flash photolysis
  - None of the above
- The photolysis of water gives the substance
  - $OH^- + H^+$
  - $H_2 + OH^-$
  - $H_2 + O_2$
  - $H_2O + H_2O_2$
- The law of photochemical equivalence was given by
  - Drapper
  - Grauths
  - Einstein
  - Labbert
- If  $I'$  is the intensity of absorbed light and  $C$  is the concentration of  $AB$  for the photochemical process  $AB + h\nu \rightarrow AB^*$ , the rate of formation of  $AB^*$  is directly proportional to
  - $C$
  - $I$
  - $I^2$
  - $C/I$

## Critical Thinking

### Objective Questions

1. For the reaction  $H_2(g) + I_2(g) \rightleftharpoons 2HI(g)$ , the rate of reaction is expressed as [CBSE PMT 1997; AIEEE 2002]

(a)  $-\frac{\Delta[I_2]}{\Delta t} = -\frac{\Delta[H_2]}{\Delta t} = \frac{1}{2} \frac{\Delta[HI]}{\Delta t}$

(b)  $\frac{\Delta[I_2]}{\Delta t} = \frac{\Delta[H_2]}{\Delta t} = \frac{\Delta[HI]}{2\Delta t}$

(c)  $\frac{\Delta[H_2]}{\Delta t} = \frac{1}{2} \frac{\Delta[I_2]}{\Delta t} = -\frac{\Delta[HI]}{\Delta t}$

(d) None of these

2. If  $3A \rightarrow 2B$ , then the rate of reaction of  $+\frac{d(B)}{dt}$  is equal to

[CBSE PMT 2002]

(a)  $+2 \frac{d(A)}{dt}$  (b)  $-\frac{1}{3} \frac{d(A)}{dt}$

(c)  $-\frac{2}{3} \frac{d(A)}{dt}$  (d)  $-\frac{3}{2} \frac{d(A)}{dt}$

3. The differential rate law for the reaction  $H_2 + I_2 \rightarrow 2HI$  is

[AIEEE 2002]

(a)  $-\frac{d[H_2]}{dt} = -\frac{d[I_2]}{dt} = +\frac{1}{2} \frac{d[HI]}{dt}$

(b)  $\frac{d[H_2]}{dt} = \frac{d[HI]}{dt} = \frac{1}{2} \frac{d[HI]}{dt}$

(c)  $\frac{1}{2} \frac{d[H_2]}{dt} = \frac{1}{2} \frac{d[I_2]}{dt} = -\frac{d[HI]}{dt}$

(d)  $-2 \frac{d[H_2]}{dt} = -2 \frac{d[I_2]}{dt} = +\frac{d[HI]}{dt}$

4. The rate law for a reaction between the substances  $A$  and  $B$  is given by, rate  $= k[A]^n[B]^m$ . On doubling the concentration of  $A$  and halving the concentration of  $B$ , the ratio of the new rate to the earlier rate of the reaction will be as

(a)  $\frac{1}{2^{(m+n)}}$  (b)  $(m+n)$

(c)  $(n-m)$  (d)  $2^{(n-m)}$

5. If we plot a graph between  $\log K$  and  $\frac{1}{T}$  by Arrhenius equation, the slope is

[UPSEAT 2001]

(a)  $-\frac{E_a}{R}$  (b)  $+\frac{E_a}{R}$

(c)  $-\frac{E_a}{2.303 R}$  (d)  $+\frac{E_a}{2.303 R}$

6. For an endothermic reaction, where  $\Delta H$  represents the enthalpy of the reaction in  $kJ/mol$ , the minimum value for the energy of activation will be

[IIT 1992]

- (a) Less than  $\Delta H$  (b) Zero  
 (c) More than  $\Delta H$  (d) Equal to  $\Delta H$

7. The rate constant ( $K'$ ) of one reaction is double of the rate constant ( $K''$ ) of another reaction. Then the relationship between the corresponding activation energies of the two reactions ( $E_a'$  and  $E_a''$ ) will be

[MP PET 1994; UPSEAT 2001]

(a)  $E_a' > E_a''$  (b)  $E_a' = E_a''$

(c)  $E_a' < E_a''$  (d)  $E_a' = 4E_a''$

8. The rate constant, the activation energy and the arrhenius parameter of a chemical reaction at  $25^\circ C$  are  $3.0 \times 10^{-4} s^{-1}$ ,

$104.4 \text{ kJ mol}^{-1}$  and  $6.0 \times 10^{14} \text{ s}^{-1}$  respectively. The value of the rate constant as  $T \rightarrow \infty$  is

[IIT 1996]

- (a)  $2.0 \times 10^{18} \text{ s}^{-1}$  (b)  $6.0 \times 10^{14} \text{ s}^{-1}$   
(c) Infinity (d)  $3.6 \times 10^{30} \text{ s}^{-1}$

9. The  $\Delta H$  value of the reaction  $\text{H}_2 + \text{Cl}_2 \rightleftharpoons 2\text{HCl}$  is  $-44.12 \text{ kcal}$ . If  $E_1$  is the activation energy of the products, then for the above reaction

[EAMCET 1997]

- (a)  $E_1 > E_2$   
(b)  $E_1 < E_2$   
(c)  $E_1 = E_2$   
(d)  $\Delta H$  is not related to  $E_1$  and  $E_2$   
(e) None is correct

10. The temperature dependence of rate constant ( $k$ ) of a chemical reaction is written in terms of Arrhenius equation,  $K = A.e^{-E^*/RT}$ . Activation energy ( $E^*$ ) of the reaction can be calculated by plotting

[CBSE PMT 2003]

- (a)  $\log k \text{ vs } \frac{1}{\log T}$  (b)  $k \text{ vs } T$   
(c)  $k \text{ vs } \frac{1}{\log T}$  (d)  $\log k \text{ vs } \frac{1}{T}$

11. Activation energy of a chemical reaction can be determined by

[CBSE PMT 1998; AFMC 1999; BHU 2000]

- (a) Changing concentration of reactants  
(b) Evaluating rate constant at standard temperature  
(c) Evaluating rate constants at two different temperatures  
(d) Evaluating velocities of reaction at two different temperatures

12. The activation energy for a reaction is  $9.0 \text{ K cal/mol}$ . The increase in the rate constant when its temperature is increased from  $298 \text{ K}$  to  $308 \text{ K}$  is

[JIPMER 2000]

- (a) 63% (b) 50%  
(c) 100% (d) 10%

13. Which of the following is the fastest reaction

[Pb. CET 2002]

- (a)  $\text{C} + \frac{1}{2} \text{O}_2 \xrightarrow{250^\circ\text{C}} \text{CO}$  (b)  $\text{C} + \frac{1}{2} \text{O}_2 \xrightarrow{500^\circ\text{C}} \text{CO}$   
(c)  $\text{C} + \frac{1}{2} \text{O}_2 \xrightarrow{750^\circ\text{C}} \text{CO}$  (d)  $\text{C} + \frac{1}{2} \text{O}_2 \xrightarrow{1000^\circ\text{C}} \text{CO}$

14. The rate constant  $k$ , for the reaction  $\text{N}_2\text{O}_5(\text{g}) \rightarrow 2\text{NO}_2(\text{g}) + \frac{1}{2} \text{O}_2(\text{g})$  is  $2.3 \times 10^{-2} \text{ s}^{-1}$ . Which equation given below describes the change of  $[\text{N}_2\text{O}_5]$  with time?  $[\text{N}_2\text{O}_5]_0$  and  $[\text{N}_2\text{O}_5]_t$  correspond to concentration of  $\text{N}_2\text{O}_5$  initially and at time,  $t$

[AIIMS 2004]

- (a)  $[\text{N}_2\text{O}_5]_t = [\text{N}_2\text{O}_5]_0 + kt$   
(b)  $[\text{N}_2\text{O}_5]_0 = [\text{N}_2\text{O}_5]_t e^{kt}$   
(c)  $\log_{10} [\text{N}_2\text{O}_5]_t = \log_{10} [\text{N}_2\text{O}_5]_0 - kt$   
(d)  $\ln \frac{[\text{N}_2\text{O}_5]_0}{[\text{N}_2\text{O}_5]_t} = kt$

15. For the reaction  $\text{CH}_3\text{COOCH}_3 + \text{H}_2\text{O} \xrightarrow{\text{H}^+} \text{CH}_3\text{COOH} + \text{CH}_3\text{OH}$  The progress of the process of reaction is followed by

- (a) Finding the amount of methanol formed at different intervals

- (b) Finding the amount of acetic acid formed at different intervals  
(c) Using a voltmeter  
(d) Using a polarimeter

16. Half life of a reaction is found to be inversely proportional to the cube of its initial concentration. The order of reaction is

[KCET 2002]

- (a) 2 (b) 5  
(c) 3 (d) 4

17. The integrated rate equation is  $Rt = \log C_0 - \log C_t$ . The straight line graph is obtained by plotting

[AIEEE 2002]

- (a) time v/s  $\log C_t$  (b)  $\frac{1}{\text{time}} \text{ v/s } C_t$   
(c) time v/s  $C_t$  (d)  $\frac{1}{\text{time}} \text{ v/s } \frac{1}{C_t}$

18. For which order reaction a straight line is obtained along with  $x$ -axis by plotting a graph between half life ( $t_{1/2}$ ) and initial concentration 'a'


[RPET 2003]

- (a) 1 (b) 2  
(c) 3 (d) 0

19. The reaction,  $X \rightarrow \text{product}$  follows first order kinetics. In 40 minutes the concentration of  $X$  changes from  $0.1 \text{ M}$  to  $0.025 \text{ M}$ . Then the rate of reaction when concentration of  $X$  is  $0.01 \text{ M}$

- (a)  $1.73 \times 10^{-4} \text{ M min}^{-1}$  (b)  $3.47 \times 10^{-5} \text{ M min}^{-1}$   
(c)  $3.47 \times 10^{-4} \text{ M min}^{-1}$  (d)  $1.73 \times 10^{-5} \text{ M min}^{-1}$

20. A substance undergoes first order decomposition. The decomposition follows two parallel first order reactions as



The percentage distribution of  $B$  and  $C$  are

[Kerala PMT 2004]

- (a) 75%  $B$  and 25%  $C$  (b) 80%  $B$  and 20%  $C$   
(c) 60%  $B$  and 40%  $C$  (d) 90%  $B$  and 10%  $C$   
(e) 76.83%  $B$  and 23.17%  $C$

## Assertion & Reason

For AIIMS Aspirants

Read the assertion and reason carefully to mark the correct option out of the options given below :

- (a) If both assertion and reason are true and the reason is the correct explanation of the assertion.  
(b) If both assertion and reason are true but reason is not the correct explanation of the assertion.  
(c) If assertion is true but reason is false.  
(d) If the assertion and reason both are false.  
(e) If assertion is false but reason is true.

1. Assertion : Instantaneous rate of reaction is equal to  $dx/dt$ .  
Reason : It is the rate of reaction at any particular instant of time.  
2. Assertion : Molecularity has no meaning for a complex reaction.

- Reason : The overall molecularity of a complex reaction is equal to the molecularity of the slowest step.
3. Assertion : If in a zero order reaction, the concentration of the reactant is doubled, the half-life period is also doubled.
- Reason : For a zero order reaction, the rate of reaction is independent of initial concentration.
4. Assertion : If the activation energy of a reaction is zero, temperature will have no effect on the rate constant.
- Reason : Lower the activation energy, faster is the reaction.
5. Assertion : According to steady state hypothesis, in a multistep reaction, the change in concentration with time for reactive intermediates is zero.
- Reason : The intermediates are so reactive that after a brief initial period their concentrations rise from zero to a small value and remains constant for most of the duration of the reaction.
6. Assertion : Half-life period of a reaction of first order is independent of initial concentration.
- Reason : Half-life period for a first order reaction  

$$t_{1/2} = \frac{2.303}{K} \log 2.$$
7. Assertion : The photochemical reactions  
 $H_2 + Cl_2 \rightarrow 2HCl$  and  $H_2 + Br_2 \rightarrow 2HBr$   
 have equal quantum efficiencies.
- Reason : Both the reactions proceed by similar mechanism.
8. Assertion : Vision is not a photochemical reaction.
- Reason : Halogenation of alkenes is a photochemical reaction.
9. Assertion : Glow worm shows chemiluminescence.
- Reason : Glow worm emits light due to oxidation of protein, luciferin present in it.
10. Assertion : The rate of reaction is always negative.
- Reason : Minus sign used in expressing the rate shows that concentration of product is decreasing.
11. Assertion : The kinetic of the reaction  
 $mA + nB + pC \rightarrow m'X + n'Y + p'Z$  obeys  
 the rate expression as  $\frac{dx}{dt} = k[A]^m[B]^n$ .
- Reason : The rate of the reaction does not depend upon the concentration of C.