

Tips & Tricks

Badecker reaction: This reaction involves the following chemical change.

 $Na_2[Fe(CN)_5NO] + Na_2SO_3 \rightarrow Na_4[Fe(CN)_5(NO.SO_3)]$

- **Exeritt's salt :** It is $K_2[Fe(CN)_6]$ obtained by reduction of prussian blue.
- **Masking:** Masking is the process in which a substance without physical separation of it is so transformed that is does not enter into a particular reaction e.g., masking of Cu^{2+} by CN^- ion.
- **Macrocyclic effect:** This term refers to the greater thermodynamic stability of a complex with a cyclic polydentate ligand when compared to the complex formed with a non-cyclic ligand. e.g., Zn(II) complex with ligand;



- \not **Prussian blue** and **Turnbull's blue** is pot. ferric ferrocyanide. However colour of Turnbull's blue is less intense than prussian blue. Decrease in colour is due to the presence in it of a white compound of the formula $K_2\{Fe[Fe(CN)_6]\}$ named as potassium ferrous ferrocyanide.
- \varnothing The colour of complexes are explained in terms of electronic transitions between the various d orbitals of different energies.
- In octahedral complexes, the energy of the d_{z^2} and $d_{x^2-y^2}$ orbitals increases much more than the energy of the d_{xy}, d_{yz}, d_{xz} orbitals. The potential energy increases as a result of repulsive interactions. The greater the repulsion, greater is the increases in energy.
- ∠ Ligands with larger groups from unstable rings than the ligands with smaller groups. This is ascribable to steric hinderance produced by a larger group.
- \varnothing When Ca^{2+} or Mg^{2+} forms complexes with EDTA, the pH of the solution decreases.
- \mathcal{E} The copper sulphate solution e.g., turns deep blue when excess of ammonia is added. This is also due to the formation of $[Cu(NH_3)_4]SO_4$
- \varnothing Co-ordination compound with a general formula MA₄, MA₃B or MABCD do not show any geometric isomerism.
- E Flexidentate character: polydentate ligand are said to

have flexidentate character if they do not use all its donar atoms to get coordinated to the metal ion e.g., EDTA generally act as a hexadentate ligand but it can also act as a pentadentate and tetradentate ligand.

 \mathcal{E} Octahedral complex of the type [M(ABCDEF)] have 15 different geometrical isomers with a pair of enantiomers. Although a few geometrical isomer have been prepared but none has been resolved e.g., [$Pt(Py)(NH_3)(NO_2)(Cl)(Br)I$].



Basic Terms

- 1. In $K_4 Fe(CN)_6$
 - (a) (CN) are linked with primary valency
 - (b) (CN) are linked with secondary valency
 - (c) *K* are linked with secondary valency
 - (d) K are linked with non-ionic valency
- 2. The co-ordination number of copper in cuprammonium sulphate is [KCET 1991, 92]
 - (a) 2

(b) 6

(c) 4

- (d) -4
- **3.** Which of the following acts as a bidentate ligand in complex formation
 - (a) Acetate
- (b) Oxalate
- (c) Thiocyanate
- (d) EDTA
- **4.** The co-ordination number of cobalt in the complex $[Co(en)_2Br_2]Cl_2$ is
 - (a) 2

(b) 6

(c) 5

- (d) 4
- **5.** Which of the following ligands forms a chelate

[MP PET/PMT 1998]

- (a) Acetate
- (b) Oxalate
- (c) Cyanide
- (d) Ammonia
- **6.** According to Werner's theory

[MP PMT 2000, 02]

- (a) Primary valency can be ionized
- (b) Secondary valency can be ionized
- (c) Primary and secondary valencies both cannot be ionized
- (d) Only primary valency cannot be ionized
- Which of the following is not true for ligand-metal complex

[MP PET 1993]

- (a) Larger the ligand, the more stable is the metal-ligand bond
- (b) Highly charged ligand forms strong bond
- (c) Larger the permanent dipole moment of ligand, the more stable is the bond
- (d) Greater the ionization potential of central metal, the stronger is the bond
- **8.** What is the co-ordination number of the metal in $[Co(en)_2 Cl_2]^+$

	908 Co-ordination Chemistry		
9.	(a) 4 (b) 5 (c) 6 (d) 3 Bidentate ligand is	20.	Which one of the following forms with an excess of <i>CN</i> (Cyanide) a complex having coordination number two
9.	(a) CN^- (b) Ethylene diammine (en)		[AIIMS 2004] (a) Cu^+ (b) Ag^+
	(c) SCN - (d) EDTA		(c) Ni^{2+} (d) Fe^{2+}
10.	The coordination number of Pt in $[Pt(NH_3)_4 Cl_2]^{++}$ ion	21.	According to Lewis the ligands are [MP PMT 2002]
	is		(a) Acidic in nature
	[MP PET 1995]		(b) Basic in nature
	(a) 2 (b) 4		(c) Neither acidic nor basic
11	(c) 6 (d) 8 Which is the example of hexadentate ligand		(d) Some are acidic and others are basic
11.	(a) 2, 2—dipyridyl	22.	The coordination number of a central metal atom in a complex is determined by [AIEEE 2004]
	(b) Dimethyl glyoxime		(a) The number of ligands around a metal ion bonded by
	(c) Aminodiacetate ion		sigma and pi-bonds both.
	(d) Ethylene diammine tetra acetate ion [EDTA]		(b) The number around a metal ion bonded by pi-bonds
12.	The coordination number of a metal in coordination compounds is [MP PET 1996; KCET (Engg./Med.) 1999]		(c) The number of ligands around a metal ion bonded by sigma bonds
	(a) Same as primary valency		(d) The number of only anionic ligands bonded to the
	(b) Sum of primary and secondary valencies		metal ion
	(c) Same as secondary valency	23.	In the extraction of which of the following, complex ion
	(d) None of these		forms [MP PET 1989] (a) Cu (b) Ag
13.	Ligand in a complex salt are [KCET 1992] (a) Anions linked by coordinate bonds to a central metal		(c) Fe (d) Na
	atom or ion	24.	Potassium ferrocyanide is a [AFMC 2000]
	(b) Cations linked by coordinate bonds to a central metal	•	(a) Normal salt (b) Mixed salt
	atom or ion		(c) Double salt (d) Complex salt
	(c) Molecules linked by coordinate bonds to a central metal atom or ion	25 .	A monodentate ligand has
	(d) Ions or molecules linked by coordinate bonds to a		(a) One co-ordinate site(b) Two co-ordinate sites
	central metal atom or ion		(c) Any number of co-ordinate sites
14.	A group of atoms can function as a ligand only when [KCET 1989; DCE 1999; MP PMT 2000]		(d) No capacity to co-ordinate
	(a) It is a small molecule	26.	EDTA has coordination number [AFMC 2004]
	(b) It has an unshared electron pair		(a) 3 (b) 4
	(c) It is a negatively charged ion	0=	(c) 5 (d) 6
	(d) It is a positively charged ion	27.	Among the properties (a) reducing (b) oxidising (c) complexing, the set of properties shown by CN^- ion
15.	Which of the following complexes show six coordination number [RPET 2000]		towards metal species is [AIEEE 2004]
	(a) $[Zn(CN)_4]^{2-}$ (b) $[Cr(H_2O)_6]^{3+}$		(a) c, a (b) b, c (c) a, b (d) a, b, c
	(c) $[Cu(CN)_4]^{2-}$ (d) $[Ni(NH_3)_4]^{2+}$	28.	That ion or molecule which forms a complex compound
16.	The number of ions formed when cuprammonium		with transitional metal ion is called
10.	sulphate is dissolved in water is [KCET 1993]		(a) Recipient (b) Ligand
	(a) 1 (b) 2		(c) Coordinate ion (d) No special name
	(c) 4 (d) Zero	29.	Coordination number of Zn in ZnS (zinc blende) is [Orissa JEE 2004]
17.	The coordination number of Cu in complex		
	$[Cu(H_2O)_4]^{++}$ is	[]	(a) 6 MP PET 1995] (c) 8 (d) 12
	(a) 4 (b) 3	30.	Wilkinson's catalyst used as a homogeneous catalyst in
	(c) 2 (d) 1		the hydrogenation of alkenes contains
18.	The primary valence of the metal ion in the co-ordination compound $K_2[Ni(CN)_4]$ is		(a) Iron (b) Aluminium
		01	(c) Rhodium (d) Cobalt Given the molecular formula of the hexa coordinated
	(a) Four(b) Zero(c) Two(d) Six	31.	complexes (A) $CoCl_3.6NH_3$ (B) $CoCl_3.5NH_3$ (C)
19.	The metal which does not form a polynuclear carbonyl is		CoCl ₃ .4 NH_3 . If the number of co-ordinated NH_3
,.	(a) Mn (b) Co		molecules in A, B and C respectively are 6, 5 and 4, the
	(c) Cr (d) Fe		primary valency in (A) , (B) and (C) are: [DCE 2003]
	• •		(a) 6, 5, 4 (b) 3, 2, 1

Co-ordination Chemistry 909 UNIVERSAL SELF SCORER



	(c) 0, 1, 2	(d) 3, 3, 3			Ferrous cyanide	(d)	Ferriferrocyanide
32.	Generally, a group of atom	s can function as a ligand if [MP PET 1996]	45.		n's reagent is		[KCET 1990]
	(a) They are positively cha			(a)	$\left[Ag(NH_3)_2\right]^+$	(b)	Ag_2O
	(b) They are free radicals			(c)	$[Cu(OH)_4]^{2-}$	(d)	Cu_2O
	charged ions	itral molecules or negatively	46.	Finel	y divided iron combin	es with	
	(d) None of these			()	F. (CO)	<i>a</i>)	[UPSEAT 2002]
33.	The ligand in potassium fer	rricyanide is			$Fe(CO)_5$		$Fe_2(CO)_9$
	(a) K^+	(b) <i>CN</i> ⁻			$Fe_2(CO)_{12}$		$Fe(CO)_6$
	(c) Fe^{3+}	(d) $(CN)_6$	47•				coordination number is
34 ·	Co-ordination number of a			(a) 6 (c) 4		(b) :	
	(a) 8 (c) 12	(b) 6 (d) 4	48.		•		les or negative groups
35∙	In $K_4 Fe(CN)_6$, Fe is in the		•	attac	hed to the central me		om in a complex ion is
33.	(a) An atom	(b) An ion		called	d Atomic number	(P)	Effective atomic number
	(c) Cationic complex	(d) Anionic complex			Coordination number		
36.	Which of the following liga	nds is expected to bidentate	49.		A combines with cation		
		[CBSE PMT 1994]			on-exchange resins		Chelates
	(a) Br	(b) $C_2 O_4^{2-}$			Clathrates		Polymers
	(c) CH_3NH_2	(d) $CH_3C \equiv N$	50.		xample of a double salt		[MP PET 2001]
37 •	In the compound lithium t is	etrahydroaluminate, the ligand			Bleaching powder $K_4[Fe(CN)_6]$	(b) 1 (d) 1	nypo Potash alum
		[AIIMS 1997]	51.	In co	mplex compounds, me	etal liga	and bond is
	(a) H^+	(b) H ⁻			Coordinate bond		Hydrogen bond
38.	(c) H Which of the following is the	(d) None of these ne odd one out [MP PET 1996]			Ionic bond		Covalent bond
30.	(a) Potassium ferrocyanid		52.				ion $[Cu(NH_3)_4]^{2+}$ with
	(b) Ferrous ammonium su	lphate			er 10ns 1n alkaline ion. What is the reasoi		ons but not in acidic [AIEEE 2003]
	(c) Potassium ferricyanide				In acidic solutions hyd		
39.	(d) Tetrammine copper (II The basic ligand is	1) sulphate					oordinate with ammonia
39.	(a) NH ₃	(b) <i>CN</i> ⁻				H_4^+ ion	ns and NH_3 molecules
	-				are not available	_	
40.	(c) F^- The negative ligand is	(d) All					isoluble $Cu(OH)_2$ is
40.	(a) Aqua	(b) Sulphato			precipitated which is so Copper hydroxide is an a		
	(c) Carboxyl	(d) Nitro sodium	53.				which type of reaction
41.	Which has yellow colour		00	_	Hydrogenation		Polymerization
	(a) Potassium cobaltinitrit(b) Potassium hexanitro co				Oxidation		Reduction
	(c) Fischer's salt	obaltate (III)	54 ·		ch of the following nometallic compound.		not considered as an
	(d) All the above			_	Cis-platia		[AIIMS 2004] Ferrocene
42.	Ligands, in complex compo	ounds [MP PMT 2003]			Zeise's salt		Gringard reagent
	(a) Accept e^- -pair		55.		-	_	pound [MP PMT 2004]
	(b) Donate e^- -pair				Lithium methoxide		Lithium dimethyl amide
	(c) Neither accept e^- -pair	r nor donate	-6		Lithium acetate		Methyl lithium
49	(d) All of these happen	is a common donor atom in	56.		queous solution of pot Гwo types of ions		m gives [UPSEAT 2004] Only one type of ion
43.	ligands	is a common donor atom in			Four types of ions		Three types of ions
		[BHU 2001]	5 7•		allite in solution in H_2		
	(a) Arsenic	(b) Nitrogen					[DCE 2003]
11	(c) Oxygen Trunbull's blue is a compo	(d) Both (b) and (c)		(a)	K^+, Mg^{2+}, Cl^-	(b)	$K^+, Cl^-, SO_4^{2-}, Br^-$
44.	Transan s sinc is a compo	[KCET 1993]		(c)	K^+, Mg^{2+}, CO_3^{2-}	(d)	$K^{+}, Mg^{2+}, Cl^{-}, Br^{-}$
	(a) Ferricyanide	(b) Ferrous ferricyanide			, , 3	(-)	, , ,



- What is the co-ordination number of cobalt in [MP PET 1994] $Co(NH_3)_3Cl_3$ (a) 3 (b) 4 6. (d) 6 (c) 5 The formula of alum is [Pb. CET 2002] (a) $K_2SO_4.Al_2(SO_4)_3.24H_2O$ (b) $K_4[Fe(CN)_6]$ 7. (c) $K_2SO_4.Al_2(SO_4)_3.6H_2O$
- (d) $Na_2CO_3.10H_2O$
- Number of ions present in $K_4[Fe(CN)_6]$ [Pb. CET 2000] (a) 2 (b) 10
- CH_3MgI is an organometallic compound due to 61.
- [DCE 2002]
 - (a) Mg-I bond
- (b) C-I bond
- (c) C Mg bond
- (d) C H bond
- What is the EAN of nickel in $Ni(CO)_A$

[BVP 2003]

(a) 34

(b) 35

(c) 32

(d) 36

Nomenclature, Oxidation state and EAN number

- How many ions are produced in aqueous solution of 1. [RPMT 2002] $[Co(H_2O)_6]Cl_2$
 - (a) 2

(b) 3

(c) 4

- (d) 6
- IUPAC name of $[Pt(NH_3)_3(Br)(NO_2)Cl]Cl$ is 2.

[CBSE PMT 1998]

- (a) Triamminechlorobromonitroplatinum (IV) chloride
- (b) Triamminebromonitrochloroplatinum (IV) chloride
- (c) Triamminebromochloronitroplatinum (IV) chloride
- (d) Triamminenitrochlorobromoplatinum (IV) chloride
- 3. Oxidation state of nitrogen is incorrectly given for

[UPSEAT 2000, 01]

Compound Oxidation state (a) $[Co(NH_3)_5 Cl]Cl_2$ 0 (b) NH_2OH -1(c) $(N_2H_5)_2SO_4$ +2(d) Mg_3N_2 -3

The formula of dichloro bis (urea) copper (II) is 4.

[CBSE PMT 1997]

- (a) $\left[Cu\{O = C(NH_2)_2\} Cl_2 \right]$
- (b) $\left[CuCl_2 \{ O = C(NH_2)_2 \}_2 \right]$
- (c) $\left[Cu\{O = C(NH_2)_2\}Cl\right]Cl$
- (d) $[CuCl_2]{O = C(NH_2)_2 H_2}$
- The IUPAC name of the complex $[Pt(NH_3), Cl_2]$ is 5.
 - (a) Platinum (II) diammino dichloride

- (b) Diammino dichloro platinate (IV)
- (c) Bis (ammino) dichloro platinum (IV)
- (d) Dichloro diammine platinum (II)
- Correct formula of diammine silver (I) chloride is
 - (a) $Ag(NH_3)Cl$
- (b) $Ag(NH_2)Cl$
- (c) $\left[Ag(NH_3)_2\right]Cl$
- (d) $\left[Ag(NH_2), Cl\right]$
- The formula of sodium nitroprusside is [AIIMS 1992]
 - (a) $Na_4[Fe(CN)_5 NOS]$
- (b) $Na_2[Fe(CN)_5NO]$
- (c) $NaFe[Fe(CN)_{\epsilon}]$

- (d) $Na_2[Fe(CN)_6NO_2]$
- The correct name of $[Pt(NH_3)_4 Cl_2][PtCl_4]$ is 8.

[MP PET 2003]

- (a) Tetraammine dichloro platinum (iv) tetrachloro platinate (ii)
- (b) Dichloro tetra ammine platinium (iv) tetrachloro platinate (ii)
- (c) Tetrachloro platinum (ii) tetraammine platinate (iv)
- (d) Tetrachloro platinum (ii) dichloro tetraammine platinate (iv)
- Correct formula of potassium ferrocyanide is 9.

[CBSE PMT 1988]

- (a) $K_4[Fe(CN)_6]$
- (b) $K_2[Fe(CN)_6]H_2O$
- (c) $K_3[Fe(CN)_6]$
- (d) None of these
- The IUPAC name of $\left[Co(NH_3)_3(NO_2)_3\right]$ is 10.
 - (a) Trinitrotriammine cobalt (III)
 - (b) Trinitrotriammine cobalt (II)
 - (c) Trinitrotriammine cobalt (III) ion
 - (d) Trinitritriammine cobaltate (III)
- 11. In $K_4[Fe(CN)_6]$, the E.A.N. of Fe is [DCE 2000]
 - (a) 33
- (b) 35
- (c) 36
- (d) 26
- Which of the following pairs is not correctly matched 12.

- (a) Effective atomic number of Pt in $[PtCl_6]^{2-} = 84$
- (b) Absorption peak for $\left[Cr^{III}(NH_3)_6\right]^{+3} = 21680cm^{-1}$
- (c) Crystal field stabilization energy of d^2 in weak ligand field = $(-)0.8\Delta_0$
- (d) Example of weak ligand field for d^5 configuration $= \left[M n^{II} F_{\epsilon} \right]^{-4}$
- The oxidation number of chromium in sodium tetrafluoro 13. oxochromate complex is
 - (a) II

- (b) IV
- (c) VI
- (d) III
- The IUPAC name of $K_4[Fe(CN)_6]$ is 14.

[CBSE PMT 1990; MP PET 1992;

MP PMT 1995, 97; Kurukshetra CET 2002]

(a) Potassium hexacyanoferrate (II)

- (b) Potassium ferrocyanide
- (c) Tetrapotassium hexacyanoferrate (II)
- (d) Tetrapotassium ferroushexacyanide (II)



15.	The IUPAC name of [Ni((CO) ₄] is [RPET 1999]				; MP PET 1997]
	(a) Tetra carbonyl nicke	el (II)		(a) Potassium ferrocya		
	(b) Tetra carbonyl nicke	el (o)		(b) Potassium hexaferr(c) Potassium ferrohex	-	
	(c) Tetra carbonyl nicke	elate (II)		(d) Potassium hexacya		
	(d) Tetra carbonyl nicke	elate (o)	27.	The EAN of iron in pota		eis
16.	The correct nomenclatur	re for $Fe_4[Fe(CN)_6]_3$ is	,	•		[Pb. CET 2000]
		[MP PMT 1994]		(a) 18	(b) 54	
	(a) Ferroso-ferric cyani	de		(c) 35	(d) 23	
	(b) Ferric-ferrous hexac	eyanate	28.	In the coordination co	ompound, $K_4[Ni(0)]$	•
	(c) Iron (III) hexacyano	oferrate (II)		state of nickel is	a >	[AIEEE 2003]
	(d) Hexacynoferrate (II	I-II)		(a) -1	(b) o	
17.	The IUPAC name of com	npound $Na_3[Co(ONO)_6]$ will be	00	(c) +1	(d) +2	m nitronmussiad
		[MP PMT 2000]	29.	According to IUPAC no is named is		BSE PMT 2003]
	(a) Hexanitritocobalt (I	II) sodium		(a) Sodium pentacyano		
	(b) Sodium cobalt nitrit	re		(b) Sodium nitroferricy		
	(c) Sodium hexanitroco	baltate (III)		(c) Sodium nitroferroc		
	(d) Sodium hexanitritoo			(d) Sodium pentacyano		
18.		ing complexes oxidation state of	30.	Pick out the complex metal atom obeys EAN		ch the central [KCET 2003]
	metal is zero	[MP PET 1997]		(a) $K_4[Fe(CN)_6]$	(b) $K_3[Fe(C)]$	
	(a) $[Pt(NH_3)_2 Cl_2]$			*		
	(c) $\left[Cr(NH_3)_3 Cl_3\right]$	(d) $\left[Cr(en)_2 Cl_2 \right]$		(c) $[Cr(H_2O)_6]Cl_3$		
19.	The oxidation number of	$f Cr in [Cr(NH_3)_6]Cl_3$ is	31.	Which of the following is $Ni(CO)_4$ has oxida		
		[CBSE PMT 2001]		•		
	(a) 8	(b) 6		(b) $Ni(CO)_4$ has zero (oxidation number i	or ivi
	(c) 4	(d) 3		(c) Ni is metal (d) CO is gas		
20.	In $[Ni(NH_3)_4]SO_4$, the	E.A.N. of Ni is	32.	Oxidation state of <i>Fe</i> in	$K_{\alpha}[Fe(CN)_{\alpha}]$	[RPMT 2002]
	(a) 34	(b) 35	J =•	(a) 2	(b) 3	[111111 2002]
	(c) 36	(d) 37		(c) 0	(d) None of	these
21.	IUPAC name of [Co(ON	$O((NH_3)_5 Cl_2)$ is	33.	Which complexes have		
		[AMU 2002]		(a) Carbonyl	(b) Ferrocya	nide
	(a) Pentaammine nitro			(c) Amine	(d) Cyanide	
	(b) Pentaammine nitrite		34.	The proper name for K	$_{2}[PtCl]_{6}$ is	[MH CET 2002]
	(c) Pentaammine nitros	so cobalt (III) chloride		(a) Potassium platinum		
	(d) Pentaammine oxo-n	nitro cobalt (III) chloride		(b) Potassium hexachle	_	
22.	The oxidation number of	f Pt in $[Pt(C_2H_4)Cl_3]^-$ is		(c) Potassium hexachle		
		[MNR 1993]	0.5	(d) Potassium hexachle IUPAC name of $K_3[Al($		
	(a) + 1	(b) +2	35.	TOTAC name of K ₃ [At(- ','3'	W
	(c) + 3	(d) +4		(a) Potassium alumino		IT 1993, 02, 03]
23.		cobalt in the complex compound		(b) Potassium aluminio		
	$[Co(NH_3)_6]Cl_3$ is			(c) Potassium trioxalat		
	(a) $+3$	(b) $+ 6$		(d) Potassium trioxalat		
	(c) +5	(d) +2	36.	The I.U.P.A.C. name of		
24.		e of potassium cuprochloride is				[MP PMT 2001]
	(a) Potassium copper (I(b) Potassium tetrachlo			(a) Potassium tri oxala		-
	(c) Tetrachloropotassiu			(b) Potassium tri oxala	to iridate (III)	
	(d) Tetrachlorocopper ((c) Potassium tris (oxa		
25.		umber of cobalt in the complex		(d) Potassium tris (oxa	lato) iridate (III)	
	$[Co(NH_3)_6]^{3+}$ is	[MP PET 2003]	37 •	The charge on $[Ag(CN)]$	₂] complex is	[AIIMS 2001]
	(a) 36	(b) 33		(a) -1	(b) +1	
	(c) 24	(d) 30		(c) +2	(d) $+3$	
26.	IUPAC name of $K_3 Fe(C)$	$(N)_6$ is				

912 Co-ordination Chemistry The IUPAC name of $[Co(NH_3)_6]Cl_3$ is [IIT-JEE 1994] 38. (a) Hexammine cobalt (III) chloride (b) Hexammine cobalt (II) chloride (c) Triammine cobalt (III) trichloride (d) None of these IUPAC name of $[Co(NH_3)_2(H_2O)_2Cl]$ Cl₂ is 39. [MP PET 1994] (a) Diaguachlorodiammine cobalt (III) chloride (b) Triamminediaquachloro cobalt (III) chloride (c) Chlorodiamminediaqua cobalt (III) chloride

- (d) Diamminediaquachloro cobalt (II) chloride
- Dichloro diammine platinum (II) complex has the 40. formula

[MP PMT 1997]

- (a) $Pt[Cl_2(NH_3)_2]$
- (b) $Pt[R.(NH_2)_2]Cl_2$
- (c) $[PtCl_2(NH_3)_2]$
- (d) $[Pt.R.(NH_2)_2]Cl_2$
- The formula of potassiumdicyano bis (oxalato) nickelate 41.
 - (a) $K_4[Ni(CN)_2(Ox)_2]$
- (b) $K_3[Ni_2(Ni_2(CN)_2(Ox)_2)]$
- (c) $K[Ni(CN)(Ox)_2]$
- (d) $K_2[Ni(CN)_2(Ox)_2]$
- The value of x which appears in the complex $[Ni(CN)_4]^x$ is 42.
 - (a) + 2
- (b) -2

(c) o

- (d) 4
- Pick the correct name of $[Co(NH_3)_5 Cl]Cl_2$ [AMU 2001] 43.
 - (a) Chloropentammine cobalt (III)
 - (b) Pentammine cobalt (III) chloride
 - (c) Chloropentammine cobalt (III) chloride
 - (d) Chloropentammine cobalt (II) chloride
- The valency of cuprammonium ion is 44.
 - (a) +4
- (b) +2
- (c) -2
- (d) -4
- In which of the following compounds transition metal has zero oxidation state [CBSE PMT 1999]
 - (a) CrO_5
- (b) $NH_2.NH_2$
- (c) NOClO₄
- (d) $[Fe(CO)_5]$
- The complex chlorocompound diaquatriammine cobalt (III) 46. chloride is represented as [CBSE PMT 2002]
 - (a) $[Co(NH_3)_3(H_2O)_3]Cl_2$
 - (b) $[Co(NH_2)_3 (H_2O)_2]Cl_2$
 - (c) $[CoCl(NH_3)_3 (H_2O)_2]Cl_3$
 - (d) $[CoCl(NH_3)_3 (H_2O)_2]Cl_2$
- The complex compound $[Co(NH_3)_3 NO_2 ClCN]$ is named 47. [MP PMT 1996]
 - (a) Chlorocyanonitrotriammine cobalt (III)
 - (b) Nitrochlorocyanotriammine cobalt (III)
 - (c) Cyanonitrochlorotriammine cobalt (III)
 - (d) Triamminenitrochlorocyano cobalt (III)
- The oxidation number of Pt in $[Pt(C_2H_4)Cl_3]^-$ is

[UPSEAT 1999, 01]

- (a) +1
- (b) +2
- (c) +3
- (d) +4

- What the structural formula of lithium 49. is tetrahydridoaluminate [MP PMT 2003]
 - (a) $Al[LiH_4]$
- (b) $Al_{2}[LiH_{4}]_{3}$
- (c) $Li[AlH_{\Delta}]$
- (d) $Li[AlH_{4}]_{2}$
- IUPAC name for $K[Ag(CN)_2]$ is 50.
 - (a) Potassium argentocyanide
 - (b) Potassium silver cyanide
 - (c) Potassium dicyanoargentate (I)
 - (d) Potassium dicyanosilver (II)
- The oxidation state of Co in $\left[Co(H_2O)_5 Cl\right]^{2+}$ is 51.
 - (a) + 2
- (b) +3
- (c) +1
- (d) + 4
- **52.** The chemical formula of diammine silver (I) chloride is [BHU 2004]
 - (a) $[Ag(NH_3)]Cl$
- (b) $[Ag(NH_3)_2]Cl$
- (c) $[Ag(NH_3)_2]Cl$
- (d) $[Ag(NH_4)_2]Cl$
- 53. IUPAC name of $[Co(NH_3)_5 NO_2]Cl_2$
- - (a) Pentamminenitrocobalt (III) chloride
 - (b) Pentamminenitrosocobalt (III) chloride
 - (c) Pentamminenitrocobalt (II) chloride
 - (d) None of these
- The pair of the compounds in which both the metals are 54. in the highest possible oxidation state is

[IIT-JEE (Screening) 2004]

- (a) $[Fe(CN)_6]^{3-}$, $[Co(CN)_6]^{3-}$
- (b) CrO_2Cl_2, MnO_4^-
- (c) TiO_3 , MnO_2
- (d) $[Co(CN)_6]^{3-}, MnO_3$
- The IUPAC name of $[Cr(NH_3)_6]^{3+}$ is 55. [Pb. CET 2001]
 - (a) Hexamminechromium (VI) ion
 - (b) Hexamminechromium (III) ion
 - (c) Hexamminechromium (II) ion
 - (d) Hexamminechloride
- 56. The IUPAC name of $K_2[Cr(CN)_2O_2(O)_2(NH_3)]$ is

- (a) Potassium ammine dicyano dioxoperoxochromate (VI)
- (b) Potassium ammine cyano peroxo dioxo chromium (VI)
- (c) Potassium ammine cyano peroxo dioxo chromiun (VI)
- (d) Potassium ammine cyano peroxo dioxo chromatic (IV)
- The IUPAC name of the coordination compound **5**7• $K_3[Fe(CN)_6]$ is [AIEEE 2005]
 - (a) Potassium hexacyanoferrate (II)
 - (b) Potassium hexacyanoferrate (III)
 - (c) Potassium hexacyanoiron (II)
 - (d) Tripotassium hexacyanoiron (II)
- Which compound is zero valent metal complex [KCET 2005] **58.**
 - (a) $[Cu(NH_3)_4]SO_4$
- (b) $[Pt(NH_3), Cl_2]$

- (c) $[Ni(CO)_A]$
- (d) $K_3[Fe(CN)_6]$

Isomerism and magnetic properties

- Which one of the following octahedral complexes will not show geometric isomerism (A and B are monodentate ligands) [CBSE PMT 2003]
 - (a) $[MA_5B]$
- (b) $[MA_2B_4]$
- (c) $[MA_3B_3]$
- (d) $[MA_4B_2]$
- The number of unpaired electrons in the complex ion 2. $[CoF_6]^{3-}$ is (Atomic no. of Co = 27) [CBSE PMT 2003]
 - (a) Zero
- (b) 2

(c) 3

- (d) 4
- Which would exhibit co-ordination isomerism 3.
 - (a) $\left[Cr(NH_3)_6\right]\left[Co(CN)_6\right]$ (b) $\left[Co(en)_2Cl_2\right]$
 - (c) $\left[Cr(NH_3)_6\right]Cl_3$
- (d) $\left[Cr(en), Cl_{2}\right]^{+}$
- and $[Co(NH_3)_5(ONO)]Cl_2$ $[Co(NH_3)_5 NO_2]Cl_2$ 4. are related to each other as
 - (a) Geometrical isomers
- (b) Optical isomers
- (c) Linkage isomers
- (d) Coordination isomers
- $[Co(NH_3)_5 SO_4]Br$ $[Co(NH_3)_5 Br] SO_4$ and are 5. examples of which type of isomerism

[MP PMT 1993, 94, 95; MP PET 1997; MP PET/PMT 1998]

- (a) Linkage
- (b) Geometrical
- (c) Ionization
- (d) Optical
- $[Co(NH_3)_4 Cl_2]NO_2$ and $[Co(NH_3)_4 Cl.NO_2]Cl$ are 6. [MP PMT 1993; MP PET 1995, 2001]
 - (a) Geometrical
- (b) Optical
- (c) Linkage
- (d) Ionization
- Which would exhibit ionisation isomerism [MP PET 1997] 7.
 - (a) $\left[Cr(NH_3)_6\right]Cl_3$
- (b) $\left[Co(NH_3)_5 Br\right] SO_4$
- (c) $\left[Cr(en), Cl_{\gamma}\right]$
- (d) $\left[Cr(en)_3 Cl_3\right]$
- $[Ti(H_2O)_6]^{+3}$ is paramagnetic in nature due to **[RPMT 2002]** 8.
 - (a) One unpaired e^-
- (b) Two unpaired e^{-}
- (c) Three unpaired e^-
- (d) No unpaired e^-
- Coordination isomerism is caused by the interchange of 9. ligands between the [UPSEAT 2002]
 - (a) Cis and Trans structure
 - (b) Complex cation and complex anion
 - (c) Inner sphere and outer sphere
 - (d) Low oxidation and higher oxidation states
- Which one of the following will not show geometrical [MP PMT 2002] isomerism
 - (a) $[Cr(NH_3)_4 Cl_2]Cl$
- (b) $[Co(en)_2Cl_2]Cl$
- (c) $[Co(NH_3)_5 NO_2]Cl_2$
- (d) $[Pt(NH_3), Cl_2]$
- Paramagnetic co-ordination compounds contain 11. electrons
 - (a) No
 - (b) Both paired and unpaired
 - (c) Paired
 - (d) Unpaired

- Which of the following isomeric pairs shows ionization 12. [MP PET 1993]
 - (a) $\left[Co(NH_3)_6\right]\left[Cr(CN)_6\right]$ and $\left[Cr(NH_3)_6\right]\left[Co(CN)_6\right]$
 - (b) $[Cr(H_2O)_6]Cl_3$ and $[Cr(H_2O)_5Cl]Cl_2.H_2O$
 - (c) $[Pt(NH_3)_2 Cl_2]$ and $[Pt(NH_3)_4][PtCl_4]$
 - (d) $[Co(NH_3)_5 Br] SO_4$ and $[Co(NH_3)_5 SO_4] Br$
- Among the following ions which one has the highest 13. paramagnetism [IIT 1993; UPSEAT 2002]
 - (a) $\left[Cr(H_2O)_6 \right]^{3+}$
- (b) $[Fe(H_2O)_6]^{2+}$
- (c) $[Cu(H_2O)_{\epsilon}]^{2+}$
- (d) $[Zn(H_2O)_6]^{2+}$
- Amongst $Ni(CO)_4$, $[Ni(CN)_4]^{2-}$ and $[NiCl_4]^{2-}$ [IIT 1991]
 - (a) $Ni(CO)_4$ and $[NiCl_4]^{2-}$ are diamagnetic and $[Ni(CN)_4]^{2-}$ is paramagnetic
 - (b) $[NiCl_4]^{2-}$ and $[Ni(CN)_4]^{2-}$ are diamagnetic and $Ni(CO)_4$ is paramagnetic
 - (c) $Ni(CO)_A$ and $[Ni(CN)_A]^{2-}$ are diamagnetic and $[NiCl_4]^{2-}$ is paramagnetic
 - (d) $Ni(CO)_4$ is diamagnetic and $[NiCl_4]^{2-}$ and $[Ni(CN)_4]^{2-}$ are paramagnetic
- $[Co(NH_3)_4 Cl_2]^+$ exhibits 15.
 - (a) Geometrical isomerism (b) Optical isomerism
 - (c) Bonding isomerism
- (d) Ionisation isomerism
- The compound which does not show paramagnetism is
 - [IIT 1992]

[AIEEE 2002]

- (a) $\left[Cu(NH_3)\right]_{A} \left[Cl_3\right]_{A}$
- (b) $\left[Ag(NH_3)\right]_2 Cl$
- (c) NO
- (d) NO_2
- The number of geometrical isomers for $[Pt(NH_3)_2 Cl_2]$ is 17. [CBSE PMT 1995]
 - (a) Two
- (b) One
- (c) Three
- (d) Four
- The pair of complex compounds $[Cr(H_2O)_6 Cl_3]$ and 18. $[Cr(H_2O)_5 Cl]Cl_2H_2O$ are an example of [MP PMT 1997]
 - (a) Linkage isomerism
- (b) Ionisation isomerism
- (c) Coordination isomerism (d) Hydrate isomerism
- The number of geometrical isomers of the complex 19. $\left[Co(NO_2)_2(NH_3)_2\right]$ is [CBSE PMT 1997]
 - (a) 2
- (b) 3
- (c) 4
- (d) o
- The type of isomerism present in nitropentamine 20. chromium (III) chloride is
 - (a) Optical
- (b) Linkage
- (c) Ionization
- (d) Polymerisation
- Which of the following compounds exhibits linkage 21. isomerism [MP PMT 2001]
 - (a) $[Co(en)_3]Cl_3$
- (b) $[Co(NH_3)_6[Cr(CN)_6]$
- (c) [Co(en), NO, Cl]Br
- (d) $[Co(NH_3)_5 Cl]Br_2$

914 Co-ordination Chemistry Pick out from the following complex compounds, a poor (c) 4 (d) 6 electrolytic conductor in solution [MP PMT 1994] The number of unpaired electrons in $Ni(CO)_4$ is (b) $[Co(NH_3)_3(NO_2)_3]$ (a) $K_2[PtCl_6]$ [AIIMS 1997] (d) $\left[Cu(NH_2)\right]_{A} \left[SO_A\right]$ (c) $K_4[Fe(CN)_6]$ (a) Zero (b) One (c) Three (d) Five The possible number of optical isomers in $[Co(en), Cl_2]^+$ 23. If the magnetic moment of $[Ag(CN)_2]^{-1}$ is zero, then the [MP PET 2003] are 35. (a) 2 (b) 3 number of unpaired electrons will be [MP PET 1995] (c) 4 (d) 6 (a) 1 (b) 2 (c) 3 (d) Zero Magnetic moment of $[Cu(NH_3)_4]^{2+}$ ion is 24. [RPET 2003] $[Pt(NH_3)_4 Cl_2]Br_2$ and $[Pt(NH_3)_4 Br_2]Cl_2$ are related to 36. (a) 1.414 each other as (d) 2.38 (c) 2.23 [MP PET 1996; AFMC 2000; CBSE PMT 2001] What is true for $[Fe(CN)_6]^{3-}$ and $[FeF_6]^{3-}$ [RPET 1999] 25. (a) Optical isomers (b) Coordinate isomers (a) Both are paramagnetic (c) Ionization isomers (d) Linkage isomers (b) Only $[Fe(CN)_6]^{3-}$ is paramagnetic 37. Which of the following complex will show geometrical as well as optical isomerism (en = ethylene diamine) (c) Only $[FeF_6]^{3-}$ is paramagnetic [KCET 1996] (d) Both are diamagnetic (a) $Pt(NH_3)_2 Cl_2$ (b) $[Pt(NH_3)Cl_A]$ Which of the following is paramagnetic [AFMC 1997] 26. (c) $[Pt(en)_2]^{4+}$ (d) $[Pt(en)_2]Cl_2$ (a) $[Ni(CO)_A]$ (b) $[Co(NH_3)_6]^{3+}$ 38. Which of the following complexes is diamagnetic (c) $[Ni(CN)_4]^{2-}$ (d) $[NiCl_A]^{2-}$ [RPMT 1997] The total number of possible isomers for the complex 27. (a) $Ni(CO)_A$ (b) $Ni(Cl_A)^{2-}$ compound $|Cu^{II}(NH_3)_4||Pt^{II}Cl_4|$ are (c) $Ni(Br)_{4}^{2-}$ (d) $NiCl_2.4H_2O$ [CBSE PMT 1998; DPMT 2004; J & K CET 2005] (a) 3 (b) 4 Which one has the highest paramagnetism 39. [AMU 2001] (c) 5 (d) 6 (a) $Ni(CO)_A$ (b) $[Ni(NH_3)_4]Cl_2$ Which following shows maximum one of the 28. (c) $[Ni(NH_3)_6]Cl_2$ (d) $[Cu(NH_3)_4]Cl_2$ paramagnetic character [AIIMS 1998] Which of the following coordination compounds would (a) $[Cr(H_2O)_6]^{3+}$ (b) $[Fe(CN)_6]^{4-}$ exhibit optical isomerism [CBSE PMT 2004] (c) $[Fe(CN)_6]^{3-}$ (d) $[Cu(H_2O)_6]^{2+}$ (a) trans-dicyanobis (ethylenediamine) chromium (III) chloride The $[Co(NH_3)_6][Cr(C_2O_4)_3]$ 29. complexes and (b) tris-(ethylenediamine) cobalt (III) bromide [AMU 2002] $[Cr(NH_3)_6][Co(C_2O_4)_3]$ (c) pentaamminenitrocobalt (III) iodide (a) Linkage isomerism (b) Geometrical isomerism (d) diamminedichloroplatinum (II) (c) Coordination isomerism (d) Ionisation isomerism Which of the following does not have optical isomer Which of the following exhibits highest molar [AIIMS 2004] 30. conductivity (a) $[Co(NH_3)_3 Cl_3]$ [MP PET 1994] (b) $[Co(en)_3]Cl_3$ (a) $\left[Co(NH_3)_6\right]Cl_3$ (b) $[Co(NH_3)_5 Cl]Cl_2$ (c) $[Co(en)_2Cl_2]Cl$ (d) $[Co(en)(NH_3)_2Cl_2]Cl$ (c) $[Co(NH_3)_A Cl_2]Cl$ (d) $\left[Co(NH_3)_3 Cl_3\right]$ Change in composition of co-ordination sphere yields 42. Which of the following compounds is colourless which types of isomers [DCE 2002] 31. (a) Optical (b) Geometrical [MP PET 1994] (d) None of these (c) Ionisation (a) $Cu_2(CH_3COO)_4.2H_2O$ (b) Cu_2Cl_2

Types of isomerism shown by

(b) Ionisation

(b) $\left[Co(NH_3)_4 Cl_2\right]Cl$

(d) $\left[Co(NH_3)_6\right]Cl_3$

[MP PET 2003]

(d) Linkage

Which of the following will not give a precipitate with

 $[Cr(NH_3)_5 NO_2]Cl_2$ is

(a) $[Co(NH_3)_3 Cl_3]$

(c) $[Co(NH_3)_5 Cl]Cl_7$

(a) Optical

 $AgNO_3$

(c) Geometrical

43.

(d) $[Cu(NH_3)_4]SO_4.4H_2O$

(b) Diamagnetism

(d) None of these

[IIT 1994]

[MP PET 1994]

The type of magnetism exhibited by $[Mn(H_2O)_6]^{2+}$ ion is

The number of isomers possible for square planar

(b) 3

(c) $CuSO_4.5H_2O$

(a) Paramagnetism

(c) Both (a) and (b)

complex $K_2[PdClBr_2(SCN)]$ is

32.

33.

(a) 2



45.	How many ions are produced from $[Co(NH_3)_6]Cl_3$ in		(b) Show optical Isomerism
	solution [RPET 1999]		(c) Show ionic Isomerism
	(a) 6 (b) 4		(d) A octahedral complex
	(c) 3 (d) 2		(e) A cationic complex
46.	The colour of $CoCl_3.5NH_3.H_2O$ is [Kerala (Med.) 2002]		
	(a) Orange yellow (b) Orange		Hybridisation and Geometry
	(c) Green (d) Violet		
	(e) Pink	1.	The correct structural formula of zeise's salt is
47.	Which one of the following is expected to be a paramagnetic complex [MP PMT 1991, 2000]		(a) $K^+ \left[PtCl_3 - \eta^2 - (C_2H_4) \right]^-$
	(a) $[Ni(H_2O)_6]^{2+}$ (b) $[Ni(CO)_4]$		(b) $K_2 \left[PtCl_3 - \eta^2 - C_2H_4 \right]$
	(c) $\left[Zn(NH_3)_A\right]^{2+}$ (d) $\left[Co(NH_3)_6\right]^{+3}$		(c) $K^{+} \left[PtCl_{2} - \eta^{2} - (C_{2}H_{4}) \right] Cl^{-}$
40	Which one of the following will give a white precipitate		•
48.	with $AgNO_3$ in aqueous mediun [MP PMT 1994]		(d) $K^{+}[PtCl_{3}(C_{2}H_{4}]^{-}]$
	(a) $[Co(NH_3), Cl](NO_2)$, (b) $[Pt(NH_3), Cl_2]$	2.	The correct order of hybridisations of central atom in
	- 1 212 - 12		$NH_3, [PtCl_4]^{2^-}, PCl_5 \text{ and } BCl_3 \text{ is}$ [MP PMT 2003]
	(c) $[Pt(en)Cl_2]$ (d) $[Pt(NH_3)_4]Cl_2$		(a) dsp^2 , dsp^3 , sp^2 and sp^3
49.	How many ions will be produced in solution from one molecule of chloropentammine cobalt (III) chloride		(b) sp^3, sp^3, sp^3d and sp^2
	(a) 1 (b) 2		(c) dsp^2, sp^2, sp^3 and dsp^3
50.	(c) 3 (d) 4 Which of the following complex will give white precipitate		(d) dsp^2 , sp^3 , sp^2 and dsp^3
J 0.	with $BaCl_2$ (aq.) [JIPMER 1997]	3.	One mole of the complex compound $Co(NH_3)_5 Cl_3$, gives
	(a) $[Co(NH_3)_4 SO_4]NO_2$ (b) $[Cr(NH_3)_5 SO_4]Cl$	Ü	3 moles of ions on dissolution in water. One mole of the
	(c) $[Cr(NH_3)_5Cl]SO_4$ (d) Both (b) and (c)		same complex reacts with two moles of $AgNO_3$ solution
51.	The number of precipitable halide ions in the sample		to yield two moles of $AgCl(s)$. The structure of the
J.,	$[Pt(NH_3)Cl_2Br]Cl$ will be		complex is
	(a) 2 (b) 3		[AIEEE 2003]
	(c) 4 (d) 1		(a) $[Co(NH_3), Cl]Cl_2$
52.	The colour of tetrammine copper (II) sulphate is		(b) $[Co(NH_3)_3 Cl_3].2NH_3$
	(a) Blue (b) Red		(c) $[Co(NH_3)_4 Cl_2]Cl.NH_3$
	(c) Violet (d) Green		(d) $[Co(NH_3)_4 Cl]Cl_2.NH_3$
53 ·	The number of ions per mole of a complex	4.	Cuprammonium ion $\left[Cu(NH_3)_4\right]^{2+}$ is
	$[CoCl_2.5 NH_3]Cl_2$ in aqueous solution will be	Τ.	[MP PMT 1997; KCET 2002]
	[MP PMT 2001] (a) Nine (b) Four		(a) Tetrahedral (b) Square planar
	(a) Nine (b) Four (c) Three (d) Two		(c) Triangular bipyramid (d) Octahedral
54.	How many unpaired electrons are present in the central	5.	In the complex $[SbF_5]^{2-}$, sp^3d hydridisation is present.
94.	metal ion of $[CoCl_4]^{2-}$ [Orissa JEE 2005]		Geometry of the complex is [Pb. PMT 2000]
	(a) 3 (b) 4		(a) Square pyramidal (b) Square bipyramidal
	(c) 5 (d) 2		(c) Tetrahedral (d) Square
55.	What is the magnetic moment of $K_3[FeF_6]$	6.	The type of hybridization involved in the metal ion of
	[Orissa JEE 2005]		$[Ni(H_2O)_6]^{2+}$ complex is
	(a) 5.91 BM (b) 4.89 BM		(a) $d^3 sp^2$ (b) $sp^3 d^2$
	(c) 3.87 BM (d) 6.92 BM		(c) sp^{3} (d) dsp^{2}
56.	(i) $K_4[Fe(CN)_6]$ (ii) $K_3[Cr(CN)_6]$ (iii) $K_3[Fe(CN)_6]$	7•	In the formation of $K_4 Fe(CN)_6$, the hybridisation involved
	(iv) $K_2[Ni(CN)_4]$	/ •	is $K_4 = (e^{\gamma t})_6$, the hybridisation involved
	Choose the complex which is paramagnetic		(a) sp^2 (b) d^2sp^3
	[Kerala CET 2005] (a) (i), (ii) and (iii) (b) (i), (iii) and (iv)		(c) d^3sp^2 (d) d^4p
	(c) (i), (iii) and (iv) (d) (i), (ii) and (iv)	8.	Which of the following is not true for metal carbonyls
	(e) (ii) and (iv)	٠.	[MP PET 1993]
57.	In coordination compound $[Co(en)_2 Cl_2]Cl$ which is false		(a) The oxidation state of the metal in the carbonyls is

zero

(b) The secondary carbonyls are obtained from photo-decomposition $% \left(\frac{1}{2}\right) =\frac{1}{2}\left(\frac{1}{2}\right) =\frac{1}{2}\left($

[Kerala CET 2005]

(a) Show geometrical Isomerism



- (c) Metal carbonyls are single bonded species
- (d) $d\pi p\pi$ overlap is observed in metal carbonvls
- An octahedral complex is formed, when hybrid orbitals of 9. the following type are involved [DCE 2003]
 - (a) sp^3
- (b) dsp^2
- (c) sp^3d^2
- (d) sp^2d
- Which one is an example of octahedral complex 10.

[MP PET 2000]

- (a) FeF_6^{3-}
- (b) $Zn(NH_3)_4^{2+}$
- (c) $Ni(CN)_4^{2-}$
- (d) $Cu(NH_3)_4^{2+}$
- Which of the following complexes has a square planar 11. geometry
 - (a) $Ag(NH_3)_2^+$
- (b) $Cu(en)_{2}^{2+}$
- (c) $[MnCl_4]^{2-}$
- (d) $Ni(CO)_{A}$
- The shape of $[Fe(CN)_6]^{4-}$ ion is 12.
 - (a) Hexagonal
- (b) Pyrimidal (d) Octagonal
- (c) Octahedral What is the shape of $Fe(CO)_5$
- 13.
- [CBSE PMT 2000]
- (a) Linear
- (b) Tetrahedral
- (c) Square planar (d) Trigonal bipyramidal
- What type of hybridization is involved in $[Fe(CN)_6]^{3-}$ 14.

[AMU 1999]

- (a) d^2sp^3
- (b) dsp^2
- (c) sp^3d^2
- (d) dsp^3
- The example of dsp^2 hybridisation is 15.

[MP PET 1999; AIIMS 2001]

- (a) $Fe(CN)_6^{3-}$
- (b) $Ni(CN)_4^{2-}$
- (c) $Zn(NH_3)_4^{2+}$
- (d) FeF_6^{3-}
- The shape of $\left[Cu(NH_3)_a\right]^{2+}$ is square planar, Cu^{2+} in this 16. complex is [NCERT 1989; RPET 1999]
 - (a) sp^3 hybridised
- (b) dsp^2 hybridised
- (c) sp^3d hybridised
- (d) sp^3d^2 hybridised
- The geometry of $Ni(CO)_4$ and $Ni(PPh_3)_2 Cl_2$ are 17.

[IIT-JEE 1999; DCE 2002]

- (a) Both square planar
- (b) Tetrahedral and square planar respectively
- (c) Both tetrahedral
- (d) Square planar and tetrahedral respectively
- Which complex has square planar structure [JIPMER 2002] 18.
 - (a) $Ni(CO)_{A}$
- (b) $[NiCl_4)^{2-}$
- (c) $[Ni(H_2O)_6]^{2+}$
- (d) $[Cu(NH_3)_4]^{2+}$
- $[Pt(NH_3)_4]Cl_2$ is 19.
- [DCE 2001]
- (a) Square planar
- (b) Tetrahedral
- (c) Pyramidal
- (d) Pentagonal
- A complex involving dsp^2 hybridization has
 - (a) A square planar geometry
 - (b) A tetrahedral geometry
 - (c) An octahedral geometry
 - (d) Trigonal planar geometry

- A tetrahedral complex ion is formed due to hybridization
 - (a) sp^2
- (b) sp^{3}
- (c) dsp^2
- (d) $d^2 s p^3$
- Back bonding is involved in which of the organometallic compounds
 - (a) $[(CH_3)_3 Al]_3$
- (b) $Mg^{2+}(C_5H_5^-)_2$
- (c) R Mg X
- (d) $[(C_5H_5)_7Fe]$
- d^2sp^3 hybridisation leads to 23.
 - (a) Hexagonal shape
- (b) Trigonal bipyrimidal
- (c) Octahedral shape
- (d) Tetrahedral shape
- Chromium hexacarbonyl is an octahedral compound 24. involving
 - (a) $sp^{3}d^{2}$
- (b) dsp^2
- (c) $d^2 s p^3$
- (d) d^3sp^2 orbitals
- $[CoF_6]^{-3}$ is formed by hybridization 25.
 - (a) $d^2 s p^3$
- (b) $d^3 sp^2$
- (c) $d^2 s p^3$
- (d) $sp^{3}d^{2}$
- The species havoing tetrahedral shape is **26.**

[IIT-JEE (Screening) 2004]

- (a) $[PdCl_4]^{2-}$
- (b) $[Ni(CN)_{4}]^{2-}$
- (c) $[Pd(CN)_4]^{2-}$
- (d) $[NiCl_{4}]^{2-}$
- Among $[Ni(CO)_4], [Ni(CN)_4]^{2-}, [NiCl_4]^{2-}$ 27. hybridization states at the Ni atom are, respectively
 - [CBSE PMT 2004; MP PMT 1992; BHU 1995; AFMC 1997]
 - (a) sp^{3}, sp^{3}, dsp^{2}
- (d) dsp^2, sp^3, sp^3
- (c) sp^3, dsp^2, dsp^2
- (d) sp^3, dsp^2, sp^3
- (At. no. of Ni = 28)
- The bond in $K_4[Fe(CN)_6]$ are: 28.
- [MP PET 2004]
 - (a) All ionic
 - (b) All covalent
 - (c) Ionic and covalent
 - (d) Ionic, covalent and co-ordinate covalent
- Hybridization of Fe in $K_3Fe(CN)_6$ is 29. [DCE 2002]
 - (a) sp^3
- (b) dsp^3
- (c) sp^3d^2
- (d) d^2sp^3
- The complex ion which has no 'd' electrons in the central 30. metal atom is [IIT-JEE Screening 2001]
 - (a) $[MnO_4]^-$
- (b) $[Co(NH_3)_6]^{3+}$
- (c) $[Fe(CN)_6]^{3-}$

32.

- (d) $[Cr(H_2O)_6]^{3+}$
- Which of the following statement is correct 31.
 - (a) $\left[Cu(NH_3)_6\right]^{2+}$ is a colourless ion
 - (b) $\left[Zn(H_2O)_6\right]^{2+}$ ion is blue coloured
 - (c) $[Ni(CN)_A]^{2-}$ ion has a tetrahedral shape
 - (d) Nickel dimethyl glyoxides is red in colour
 - Which of the following shall form an octahedral complex [DCE 2001]
 - (a) d^4 (low spin)
- (b) d^8 (high spin)
- (c) d^6 (low spin)
- (d) None of these



	(a) CN^{-} (b) NO_{2}^{-}		(a) $[Fe(OH)_3]^{3-}$ (b) $[Fe(Cl)_6]^{3-}$
	(c) en (d) NH_3		(c) $[Fe(CN)_6]^{3-}$ (d) $[Fe(H_2O)_6]^{3+}$
34.	The strongest ligand in the following is [MP PET 199	5l 3 .	The most stable complex among the following is
	(a) CN^- (b) Br^-		[MP PMT 2002]
	(c) HO^- (d) F^-		(a) $K_3[Al(C_2O_4)_3]$ (b) $[Pt(en)_2]Cl_2$
35.	The neutral ligand is		(c) $Ag(NH_3)_2Cl$ (d) $K_2(Ni(EDTA)]$
	(a) Chloro (b) Hydroxo	4.	Which of the following factors tends to increase the
a.e	(c) Ammine (d) Oxalato	1	stability of metal ion complexes (a) Higher ionic radius of the metal ion
36.	The ligands which can get attached to the central met ion through more than one atom are called	.aı	(b) Higher charge/size ratio of the metal ion
	(a) Ambident ligands (b) Polydentate ligands		(c) Lower ionisation potential of the metal ion
	(c) Chelate ligands (d) Neutral ligands	_	(d) Lower basicity of the ligand
37•	A strong ligand gives a complex which is generally called	5.	CuSO ₄ reacts with KCN solution and forms:[DPMT 2004]
	(a) High spin (b) High energy (c) Low spin (d) Stable		(a) $K_3[Cu(CN)_4]$ (b) $Cu(CN)$
.0			(c) $Cu(CN)_2$ (d) $K_4[Cu(CN)_6]$
38.	CN^- is a strong field ligand. This is due to the fact that [CBSE PMT 200	6.	A co-ordination complex compound of cobalt has the
	(a) It can accept electron from metal species	-1 1	molecular formulae containing five ammonia molecules, one nitro group and two chlorine atoms for one cobalt
	(b) It forms high spin complexes with metal species		atom. One mole of this compound produces three mole
	(c) It carries negative charge.		ions in an aqueous solution on reacting with excess of
	(d) It is a pseudohalide		$AgNO_3$, $AgCl$ precipitate. The ionic formula for this
39.	Considering H_2O as a weak field ligand, the number		complex would be: [DPMT 2004; Kerala PMT 2004]
	unpaired electrons in $[Mn(H_2O)_6]^{2+}$ will be (At. No. of Mathematical Equation 2)		(a) $[Co(NH_3)_5(NO_2)]Cl_2$
	= 25) asas a ajkk [CBSE PMT 200	4]	(b) $[Co(NH_3)_5 Cl][Cl(NO_2)]$
	(a) Two (b) Four (c) Three (d) Five		(c) $[Co(NH_3)_4(No_2)Cl][(NH_3)Cl]$
40.	Which of the following is a π - complex		(d) $[Co(NH_3)_5][(NO_2)_2Cl_2]$
•	(a) Trimethyl aluminium (b) Ferrocene	7•	In any ferric salt, on adding potassium ferrocyanide, a
	(c) Diethyl zinc (d) Nickel carbonyl	/ •	prussian blue colour is obtained, which is
41.	Which of the following is a π – acid ligand		[BIT 1992; BHU 2002]
	[KCET 1996; AHMS 200	3]	(a) $K_3 Fe(CN)_6$ (b) $KFe[Fe(CN)_6]$
	(a) NH_3 (b) CO		(d) $FeSO_4.Fe(CN)_6$ (d) $Fe_4[Fe(CN)_6]_3$
	(c) F^- (d) Ethylene diamine	8.	Prussian blue is formed when [CBSE PMT 1990]
42.	The value of the 'spin only' magnetic moment for one the following configurations is 2.84 BM. the correct one	of	(a) Ferrous sulphate reacts with FeCl ₃
	[AIEEE 200	5]	(b) Ferric sulphate reacts with $K_4[Fe(CN)_6]$
	(a) d^4 (in strong ligand field)		(c) Ferrous ammonium sulphate reacts with $FeCl_3$
	(b) d^4 (in weak ligand field)		(d) Ammonium sulphate reacts with FeCl ₃
	(c) d^3 (in weak as well as in strong fields)	9.	Complex salt can be made by the combination of
	(d) d^5 (in strong ligand field)		$[Co^{III}(NH_3)_5 Cl]^X$ with [RPMT 2000, AFMC 2002]
43.	The geometry of $Ni(CO)_4$ and $Ni(PPh_3)_2Cl_2$ are		(a) PO_4^{3-} (b) Cl^{-}
	[BHU 200	5]	• • • • • • • • • • • • • • • • • • • •
	(a) Both square planer	10.	(c) $2Cl^-$ (d) $2K^+$ Which reagent can be used to identify nickel ion
	(b) Tetrahedral and square planar respectively(c) Both tetrahedral	10.	(a) Resorcinol
	(d) Square planar and tetrahedral respectively		(b) Dimethyl glyoxime [DMG]
	(1) 14-11-17	_	(c) Diphenyl benzidine
	Complexes and complex stability		(d) Potassium ferrocyanide
		— 11.	Dimethyl glyoxime forms a coloured complex with
1.	Which of the following complexes has the highest stabili	ty	(a) Ag (b) Ni
	constant at 298 K	12.	(c) Cr (d) Zn Silver chloride dissolves in excess of NH_4OH . The cation
	(a) $[CdCl_4]^{2-}$ (b) $[CdBr_4]^{2-}$	14.	present in this solution is [EAMCET 1998]
	(c) $[CdI_4]^{2-}$ (d) $[Cd(CN)_4]^{2-}$		present in this solution is [EAMCEI 1996]
	- · · · · · · · · · · · · · · · · · · ·		

2.

The most stable ion is

Which one of the following is a strong field ligand

33.



- (a) Ag^+
- (b) $[Ag(NH_2)_2]^+$
- (c) $\left[Ag(NH_3)_4\right]^+$
- (d) $[Ag(NH_3)_6]^+$
- Silver sulphide dissolved in a solution of sodium 13. cynamide to form the complex [AMU 1999]
 - (a) $Na[Ag(CN)_2]$
- (b) $Na_3[Ag(CN)_4]$
- (c) $Na_5[Ag(CN)_6]$
- (d) $Na_{2}[Ag(CN)_{2}]$
- Which one will give Fe^{3+} ions in solution 14.
 - (a) $[Fe(CN)_6]^{3-}$
 - (b) $Fe_2(SO_4)_3$
 - (c) $[Fe(CN)_6]^{4-}$
 - (d) $NH_4(SO_4)_2.FeSO_4.6H_2O$
- The cation that does not form an amine complex with 15. excess of ammonia is
 - (a) Cd^{2+}
- (b) Al^{3+}
- (c) Cu²⁺
- (d) Ag^+

Application of organometallics

- Ziegler-Natta catalyst is an organometallic compound of 1. which metal [J & K CET 2005]
 - (a) Iron
- (b) Zirconium
- (c) Rhodium
- (d) Titanium
- In the case of small cuts, bleeding is stopped by applying 2. potash alum. Here alum acts as

[KCET (Med.) 2001]

- (a) Fungicide
- (b) Disinfectant
- (c) Germicide
- (d) Coagulating agent
- The formula of ferrocene is 3.
 - (a) $[Fe(CN)_6]^{4-}$
- (b) $[Fe(CN)_6]^{3-}$
- (c) $\left[Fe(CO)_5 \right]$
- (d) $[(C_5H_5), Fe]$
- Which of the following is formed when n-butyl lithium 4. reacts with tin (II) chloride [AFMC 2001]
 - (a) LiBr
- (b) Et_4Pb
- (c) $(C_4H_9)_4Sn$
- (d) $(C_2H_5)_4Pb$
- Which of the following organo-silicon compound on 5. hydrolysis will give a three dimensional silicone

[Orissa JEE 2003]

- (a) R_3SiCl
- (b) $RSiCl_3$
- (c) SiCl₄
- (d) R_2SiCl_2
- Which one is not an organometallic compound 6.

[J & K CET 2005; Pb. CET 2003]

- (a) RMgX
- (b) C_2H_5ONa
- (c) $(CH_3)_4 Sn$
- (d) KC_4H_9
- The complex used as an anticancer agent is [AIIMS 2003]
 - (a) trans $-[Co(NH_3)_3Cl_3]$ (b) cis $-[PtCl_2(NH_3)_2]$
 - (c) $cis K_2[PtCl_2Br_2]$
- (d) Na_2CO_3
- 8. The compound that is not olefinic organometallic is
 - (a) $K[C_2H_4PtCl_3]3H_2O$ (b) $Be(CH_2)_2$
 - (c) $(C_2H_4PtCl_3)_2$
- (d) $C_{\perp}H_{\perp}Fe(CO)_{3}$

- Among the following, which is not the π -bonded Q. organometallic compound [CBSE PMT 2003]
 - (a) $(CH_3)_4 Sn$
- (b) $K[PtCl_3(\eta^2 C_2H_4)]$
- (c) $Fe(\eta^5 C_5H_5)_2$
- (d) $Cr(\eta^6 C_6H_6)_2$
- Wilkinson's catalyst is used in 10.
 - (a) Polymerization
- (b) Condensation
- (c) Halogenation
- (d) Hydrogenation
- What is the use of tetraethyl lead 11.
 - (a) As a catalyst in addition reaction of alkenes
 - (b) As a catalyst in polymerization reaction of alkenes
 - (c) For reducing knocking
 - (d) For creating knocking
- Which of the following is an organo-metallic compound 12.

[MP PMT 2001]

- (a) Lithium ethoxide
- (b) Ethyl lithium
- (c) Lithium acetate
- (d) Lithium carbide
- Which of the following is an organometallic compound 13. [AIIMS 1997]
 - (a) $Ti(C_2H_5)_4$
- (b) $Ti(OC_2H_5)_4$
- (c) $Ti(OCOCH_3)_4$
- (d) $Ti(OC_6H_5)_4$
- Which of the following is not an organometallic 14. compound

[MP PET 1996; BHU 2002]

- (a) Ethyl magnesium bromide
- (b) Tetraethyl lead
- (c) Sodium ethoxide
- (d) Trimethyl aluminium
- An organometallic compound amongst the following is 15.
 - (a) Ferrocene

16.

18.

- (b) CaC_2
- (c) Tetraethyl lead (TEL) (d) All of these
- Which of the following does not have a metal carbon bond [CBSE PMT 2004]
- (a) $K[Pt(C_2H_4)Cl_3]$
 - (b) $Ni(CO)_4$
- (c) $Al(OC_2H_5)_3$
- (d) C_2H_5MgBr
- Coordination compounds have great importance in 17. biological systems. In this context which of the following statements is incorrect [AIEEE 2004]
 - (a) Cyanocobalamin is B_{12} and contains cobalt.
 - (b) Haemoglobin is the red pigment of blood and contains iron.
 - (c) Chlorophylls are green pigments in plants and contains calcium.
 - (d) Carboxypepticase-A is an enzyme and contains zinc.
 - Ziegler-Natta catalyst is (a) $(Ph_3P)_3RhCl$
- (b) $K[PtCl_3(C_2H_4)]$

[Pb. CET 2004]

- (c) $[Al_2(C_2H_6)_6] + TiCl_4$
- (d) $[Fe(C_2H_5)_2]$
- Dimethyl glyoxime gives a red precipitate with Ni^{2+} , 19. which is used for its detection. To get this precipitate readily the best *pH* range is [AIIMS 2004] (a) < 1(b) 2-3
- (d) 9-11 (c) 3-4 The π -bonded organo metallic compound which has 20. ethene as one of its component is [J & K CET 2005]
 - (a) Zeise's salt
- (b) Ferrocene
- (c) Dibenzene chromium
- (d) Tetraethyl tin.



. Critical Thinking

Objective Questions

- The coordination number and oxidation state of Cr in $K_3[Cr(C_2O_4)_3]$ are, respectively [CBSE PMT 1995]
 - (a) 4 and +2
- (b) 6 and +3
- (c) 3 and +3
- (d) 3 and o
- The complex compounds which result from the 2. coordination of carbon monoxide are known as
 - (a) Electronic
- (b) Carbonyls
- (c) Carbonates
- (d) Carbon permono
- Mixture X = 0.02 mol of $[Co(NH_3)_5 SO_4]Br$ and 0.02 3. mol of $[Co(NH_3)_5Br]SO_4$ was prepared in 2 litre of
 - 1 litre of mixture $X + \text{excess } AgNO_3 \rightarrow Y$.
 - 1 *litre* of mixture $X + \text{excess } BaCl_2 \rightarrow Z$

Number of moles of Y and Z are

[IIT JEE 2003]

- (a) 0.01, 0.01
- (b) 0.02, 0.01
- (c) 0.01, 0.02 (d) 0.02, 0.02
- Which of the following organometallic compound is σ and π bonded [MH CET 2001; PCET 2002]
 - (a) $Fe(CH_3)_3$
 - (b) $[Fe(\eta^5 C_5H_5)_2]$
 - (c) $[Co(CO)_5 NH_3]^{2+}$
 - (d) $K[PtCl_3(n^2 C_2H_4)]$
- In $[Ni(NH_3)_4]SO_4$, the valency and coordination number 5. of Ni will be respectively
 - (a) 3 and 6
- (b) 2 and 4
- (c) 4 and 2
- (d) 4 and 4
- Co-ordination number of Fe in the complexes 6. $[Fe(CN)_6]^{4-}$, $[Fe(CN)_6]^{3-}$ and $[FeCl_A]^{-}$ would be respectively [MP PET 2003]
 - (a) 2, 3, 3
- (b) 6, 6, 4
- (c) 6, 3,3
- (d) 6, 4, 6
- On hydrolysis $(Me)_2 SiCl_2$ will produce [IIT-JEE 2003] 7.
 - (a) $(Me)_2 Si(OH)_2$
 - (b) $(Me)_2 Si = O$
 - (c) $-[-O-(Me)_2Si-O-]_n$
 - (d) $Me_2SiCl(OH)$
- 8. Which of the following represents a chelating ligand

[JIPMER 2002]

- (a) H_2O
- (b) OH
- (c) DMG
- (d) Cl⁻
- CuSO₄ reacts with KCN solution and forms 9.

[MP PMT 1992; IIT 1996; UPSEAT 2001, 02]

- (a) $Cu(CN)_2$
- (b) Cu(CN)
- (c) $K_2[Cu(CN)_4]$
- (d) $K_3[Cu(CN)_4]$

In the process of extraction of gold. 10.

Roasted gold ore $+CN^- + H_2O \xrightarrow{O_2} [X] + OH^-$

$$[X] + Zn \rightarrow [Y] + Au$$

Identify the complexes [X] and [Y]

- (a) $X = [Au(CN)_2]^-, Y = [Zn(CN)_4]^{2-}$
- (b) $X = [Au(CN)_A]^{3-}, Y = [Zn(CN)_A]^{2-}$
- (c) $X = [Au(CN)_2]^-, Y = Zn(CN)_6]^{4-}$
- (d) $X = [Au(CN)_{4}]^{-}, Y = [Zn(CN)_{4}]^{2-}$
- 11. A solution of potassium ferrocyanide would contains

 [BHU 1999] [KCET 1990]
 - (a) 2

(b) 3

(c) 4

- (d) 5
- 12. Which one of the following complexes is an outer orbital [AIEEE 2004] complex
 - (a) $[Co(NH_3)_6]^{3+}$
- (b) $[Mn(CN)_6]^{4-}$
- (c) $[Fe(CN)_6]^{4-}$
- (d) $[Ni(NH_3)_6]^{2+}$

Atomic nos: Mn = 25, Fe = 26, Co = 27, Ni = 28

Which one of the following has largest number of isomers 13.

[AIEEE 2004]

- (a) $[Ir(PP_3)_2H(CO)]^{2+}$
- (b) $[Co(NH_3)_5 Cl]^{2+}$
- (c) $[Ru(NH_3)_4 Cl_2]^+$
- (d) $[Co(en), Cl_2]^+$


(R = alkyl group; en = ethylenediamine)


- Which kind of isomerism is exhibited by octahedral [IIT-JEE Screening 2005] $Co(NH_3)_4 Br_2 Cl$?
 - (a) Geometrical and Ionization
 - (b) Geometrical and Optical
 - (c) Optical and Ionization
 - (d) Geometrical only
- Which one of the following is expected to exhibit optical isomerism (en = ethylenediamine)
 - (a) $cis [Pt(NH_3)_2Cl_2]$
- (b) $trans [Co(en), Cl_2]$
- (c) $trans [Pt(NH_3)_2Cl_2]$ (d) $cis [Co(en)_2Cl_2]$
- $[EDTA]^{4-}$ is a: 16.
- [UPSEAT 2004]
- (a) Monodentate ligand (c) Quadridentate ligand
- (b) Bidentate ligand (d) Hexadentate ligand
- Which of the following statements is incorrect? 17.


- (a) In $K_3[Fe(CN)_6]$, the ligand has satisfied only the secondary valency of ferric ion.
- (b) In $K_3[Fe(CN)_6]$, the ligand has satisfied both primary and secondary valencies of ferric ion.
- (c) In $K_4[Fe(CN)_6]$, the ligand has satisfied both primary and secondary valencies of ferrous ion.
- (d) In $[Cu(NH_3)_4]SO_4$, the ligand has satisfied only the secondary valency of copper.
- Which of the following is considered as an anticancer 18. species. [CBSE PMT 2004]

UNIVERSAL SELF SCORER

920 Co-ordination Chemistry







(d)
$$\begin{bmatrix} Cl & Cl \\ Pt & Cl \end{bmatrix}$$

19. An aqueous solution of $CoCl_2$ on addition of excess of concentrated HCl turns blue due to formation of

[AIIMS 2005]

- (a) $[Co(H_2O)_4 Cl_2]$
- (b) $[Co(H_2O)2Cl_4]^{2-}$
- (c) $[CoCl_4]^{2-}$
- (d) $[Co(H_2O)2Cl_2]$
- **20.** The correct order for the wavelength of absorption in the visible region is **[AIIMS 2005]**
 - (a) $[Ni(NO_2)_6]^{4-} < [Ni(NH_3)_6]^{2+} < [Ni(H_2O)_6]^{2+}$
 - (b) $[Ni(NO_2)_6]^{4-} < [Ni(H_2O)_6]^{2+} < [Ni(NH_3)_6]^{2+}$
 - (c) $[Ni(H_2O)_6]^{2+} < [Ni(NH_3)_6]^{2+} < [Ni(NO_2)_6]^{4-}$
 - (d) $[Ni(NH_3)_6]^{2+} < [Ni(H_2O)_6]^{2+} < [Ni(NO_2)_6]^{4-}$
- 21. In which of the following pairs both the complexes show optical isomerism [AIIMS 2005]
 - (a) $cis-[Cr(C_2O_4)_2Cl_2]^{3-}$, $cis-[Co(NH_3)_4Cl_2]$
 - (b) $[Co(en)_3]Cl_3$, $cis-[Co(en)_2Cl_2]Cl$
 - (c) [PtCl(dien)]Cl, $[NiCl_2Br_2]^{2-}$
 - (d) $[Co(NO_3)_3(NH_3)_3]$, cis- $[Pt(en)_2Cl_2]$
- **22.** Which of the following compounds shows optical isomerism

[AIEEE 2005; CBSE PMT 2005]

- (a) $[Cu(NH_3)_4]^{2+}$
- (b) $[ZnCl_4]^{2-}$
- (c) $[Cr(C_2O_4)_3]^{3-}$
- (d) $[Co(CN)_6]^{3-}$



Assertion & Reason

For AIIMS Aspirants

Read the assertion and reason carefully to mark the correct option out of the options given below:

- (a) If both assertion and reason are true and the reason is the correct explanation of the assertion.
- (b) If both assertion and reason are true but reason is not the correct explanation of the assertion.
- (c) If assertion is true but reason is false.
- (d) If the assertion and reason both are false.
- (e) If assertion is false but reason is true.
- **1.** Assertion : Potassium ferrocyanide and potassium ferricyanite both are diamagnetic.
 - Reason : Both have unpaired electrons.
- **2.** Assertion : NF_3 is a weaker ligand than $N(CH_3)_3$

Reason : NF_3 ionizes to give F^- ions

ionsiionsioiiioioioioioioioioio

ions in aqueous solution.

Reason

3. Assertion : The $[Ni(en)_3]Cl_2$ (en = ethylene diamine)

has lower stability than $[Ni(NH_3)_6]Cl_2$

: In [Ni(en)₃]Cl₂ the geometry of Ni is trigonal bipyramidal

4. Assertion : The ligands nitro and nitrito are called ambidentate ligands.

Reason : These ligands give linkage isomers.

5. Assertion : Geometrical isomerism is also called *cistrans* isomerism.

Reason : Tetrahedral complexes show geometrical isomerism.

6. Assertion : $\left[(en)_2 Co \bigvee_{OH}^{NH} Co(en)_2 \right]^{3+}$ is named as

tetrakis (ethylene diamine) μ -hydroxoimido dicobalt (III) ion.

Reason : In naming polynuclear complexes *i.e.*, containing two or more metal atoms joined by bridging ligands, the word μ is added with hyphen before the name of such ligands.

7. Assertion : $[Fe(CN)_6]^{3-}$ is weakly paramagnetic while $[Fe(CN)_6]^{4-}$ is diamagnetic.

Reason : $[Fe(CN)_6]^{3^-}$ has +3 oxidation state while $[Fe(CN)_6]^{4^-}$ has +2 oxidation state.

8. Assertion : $H_2N - NH_{2}$ is a chelating ligand.

Reason : A chelating ligand must possess two or more lone pairs at such a distance that it may form suitable strain free rings at the metal ion.

9. Assertion : $[Ti(H_2O)_6]^{3+}$ is coloured while $[Sc(H_2O)_6]^{3+}$ is colourless.

Reason : d-d transition is not possible in $[Sc(H_2O)_6]^{3+}$.

10. Assertion : All the octahedral complexes of Ni^{2+} must be outer orbital complexes.

Reason : Outer orbital octahedral complexes are given by weak ligands.

11. Assertion : Potassium ferrocyanide is diamagnetic

whereas potassium fericyanide is paramagnetic.

Reason : Crystal field splitting in ferrocyanide ion is greater than that of ferricyanide ion.

[AIIMS 2005]



Basic Terms



1	b	2	С	3	b	4	b	5	d
6	а	7	b	8	С	9	b	10	С
11	d	12	С	13	d	14	b	15	b
16	b	17	а	18	С	19	С	20	b
21	b	22	С	23	b	24	d	25	а
26	d	27	а	28	b	29	b	30	С
31	b	32	С	33	b	34	b	35	d
36	d	37	b	38	b	39	d	40	b
41	d	42	b	43	d	44	b	45	а
46	a	47	d	48	С	49	b	50	d
51	a	52	b	53	d	54	а	55	d
56	d	57	a	58	d	59	а	60	d
61	С	62	d						

Nomenclature, oxidation State and EAN number

1	b	2	С	3	С	4	b	5	d
6	С	7	b	8	а	9	а	10	а
11	С	12	а	13	b	14	а	15	b
16	С	17	d	18	b	19	d	20	а
21	b	22	b	23	а	24	b	25	а
26	d	27	С	28	b	29	а	30	а
31	а	32	b	33	а	34	b	35	С
36	b	37	а	38	а	39	b	40	С
41	а	42	b	43	С	44	b	45	b
46	d	47	а	48	b	49	С	50	С
51	b	52	b	53	С	54	d	55	b
56	а	57	b	58	С				

Isomerism and Magnetic properties

1	а	2	d	3	а	4	С	5	С
6	d	7	b	8	а	9	а	10	С
11	d	12	d	13	b	14	С	15	а
16	b	17	а	18	d	19	а	20	b
21	С	22	b	23	b	24	а	25	а
26	d	27	d	28	а	29	С	30	а
31	b	32	а	33	С	34	а	35	d
36	С	37	d	38	а	39	d	40	b
41	а	42	С	43	d	44	а	45	b
46	е	47	а	48	d	49	С	50	С
51	d	52	а	53	С	54	а	55	а
56	С	57	С						

Hybridisation and Geometry

1	а	2	b	3	а	4	b	5	a
			b	8	d	9	С	10	a
11	b	12	С	13	d	14	а	15	b
16	b	17	d	18	d	19	а	20	а

21	b	22	d	23	С	24	С	25	С
26	d	27	d	28	d	29	d	30	a
31	d	32	С	33	а	34	а	35	С
36	а	37	С	38	d	39	d	40	b
41	b	42	а	43	С				

Complexes and complex stability

1	d	2	d	3	d	4	b	5	а
6	а	7	d	8	b	9	С	10	b
11	b	12	b	13	а	14	b	15	b

Application of organometallics

1	d	2	d	3	d	4	С	5	b
6	b	7	b	8	а	9	а	10	d
11	С	12	b	13	а	14	С	15	d
16	С	17	С	18	С	19	d	20	а

Critical Thinking Questions

1	b	2	b	3	а	4	b	5	b
6	b	7	С	8	С	9	d	10	а
11	d	12	d	13	d	14	а	15	d
16	d	17	ad	18	а	19	С	20	а
21	b	22	С						

Assertion & Reason

1	d	2	С	3	d	4	а	5	С
6	е	7	b	8	е	9	а	10	b
11	С								