

Ordinary Thinking Objective Questions

Electrolytes and Electrolysis

- Which of the following will not conduct electricity in aqueous solution [AMU 1982, 83]
 - Copper sulphate
 - Sugar
 - Common salt
 - None of these
- Strong electrolytes are those which [MNR 1983]
 - Dissolve readily in water
 - Conduct electricity
 - Dissociate into ions at high dilution
 - Completely dissociate into ions at all dilutions
- In aqueous solution, strong electrolytes [AMU 1983, 84]
 - Are partially ionized
 - Do not ionise
 - Ionise almost completely
 - Form polymers
- An electrolyte [KCET 1984; MP PET/PMT 1988]
 - Forms complex ions in solution
 - Gives ions only when electricity is passed
 - Possesses ions even in solid state
 - Gives ions only when dissolved in water
- Electrolytes when dissolved in water dissociates into ions because [CPMT 1974, 78; MNR 1983]
 - They are unstable
 - The water dissolves it
 - The force of repulsion increases
 - The forces of electrostatic attraction are broken down by water
- Electrolyte can conduct electricity because
 - Their molecules contain unpaired electrons, which are mobile
 - Their molecules contain loosely held electrons which get free under the influence of voltage
 - The molecules break up into ions when a voltage is applied
 - The molecules are broken up into ions when the electrolyte is fused or is dissolved in the solvent
- Which one of the following metals could not be obtained on electrolysis of aqueous solution of its salts [IIT 1990]
 - Ag
 - Mg
 - Cu
 - Cr
- Which of the following aqueous solution will conduct an electric current quite well [MP PMT 1987]
 - Glycerol
 - HCl
 - Sugar
 - Pure water
- On the electrolysis of aqueous solution of sodium sulphate, on cathode we get [MP PMT 1992, 2002]
 - Na
 - H₂
 - SO₂
 - SO₃
- Electrolysis involves oxidation and reduction respectively at [CPMT 1973; AMU 1983; NCERT 1983, 84; MH CET 2001]
 - Anode and cathode
 - Cathode and anode
 - At both the electrodes
 - None of the above
- Which of the following compounds will not undergo decomposition on passing electricity through aqueous solution
 - Sugar
 - Sodium Chloride
 - Sodium Bromide
 - Sodium Acetate
- During the electrolysis of an electrolyte, the number of ions produced, is directly proportional to the [AFMC 2002]
 - Time consumed
 - Electro chemical equivalent of electrolysis
 - Quantity of electricity passed
 - Mass of electrons
- When the sample of copper with zinc impurity is to be purified by electrolysis, the appropriate electrodes are [AIEEE 2002]

Cathode	Anode
(a) Pure zinc	Pure copper
(b) Impure sample	Pure copper
(c) Impure zinc	Impure sample
(d) Pure copper	Impure sample
- In the electrolytic cell, flow of electrons is from [IIT Screening 2003]
 - Cathode to anode in solution
 - Cathode to anode through external supply
 - Cathode to anode through internal supply
 - Anode to cathode through internal supply
- An electric current is passed through an aqueous solution of the following. Which one shall decompose [NCERT 1972]
 - Urea
 - Glucose
 - AgNO₃
 - Ethyl alcohol
- The electric conduction of a salt solution in water depends on the
 - Shape of its molecules
 - Size of its molecules
 - Size of solvent molecules
 - Extent of its ionization
- A solution of sodium sulphate in water is electrolysed using inert electrodes. The products at the cathode and anode are respectively [IIT 1987, 96]
 - H₂, O₂
 - O₂, H₂
 - O₂, Na
 - O₂, SO₂
- On electrolysis a solution of dilute H₂SO₄ between platinum electrodes, the gas evolved at the anode is [NCERT 1977, 79; MNR 1980; CBSE PMT 1992]
 - SO₂
 - SO₃
 - O₂
 - H₂
- The addition of a polar solvent to a solid electrolyte results in
 - Polarization
 - Association
 - Ionization
 - Non-liberation of heat
- During the electrolysis of fused NaCl, which reaction occurs at anode [NCERT 1973; AFMC 1992; MP PMT 2002]
 - Chloride ions are oxidized
 - Chloride ions are reduced
 - Sodium ions are oxidised
 - Sodium ions are reduced
- The amount of ion discharged during electrolysis is not directly proportional to [NCERT 1973]
 - Resistance
 - Time
 - Current
 - Chemical equivalent of the ion
- Electrolysis of aqueous HCl solution produces [CPMT 1987]
 - H₂ gas at the anode
 - H₂ gas at the cathode
 - Cl₂ gas at the cathode

- (d) Cl_2 and O_2 gases both at the anode
23. During electrolysis of $NaCl$ solution, part of the reaction is $Na^+ + e^- \rightarrow Na$. This is termed as [NCERT 1984]
(a) Oxidation (b) Reduction
(c) Deposition (d) Cathode reaction
24. When a solution of an electrolyte is heated the conductance of the solution [KCET 1991]
(a) Increases because of the electrolyte conducts better
(b) Decreases because of the increased heat
(c) Decreases because of the dissociation of the electrolyte is suppressed
(d) Increases because the electrolyte is dissociated more
25. The passage of current liberates H_2 at cathode and Cl_2 at anode. The solution is [EAMCET 1979,87]
(a) Copper chloride in water (b) $NaCl$ in water
(c) H_2SO_4 (d) Water
26. Pure water does not conduct electricity because it [Manipal MEE 1995]
(a) Has a low boiling point
(b) Is almost totally unionized
(c) Is neutral
(d) Is readily decomposed
27. Which is responsible for electrical conduction of molten sodium chloride [MADT Bihar 1995]
(a) Free electrons
(b) Free ions
(c) Free molecules
(d) Atoms of sodium and chlorine
28. In electrolysis of aqueous copper sulphate, the gas at anode and cathode is [AFMC 1995]
(a) O_2 and H_2 (b) SO_2 and H_2
(c) H_2 and O_2 (d) SO_3 and O_2
29. Use of electrolysis is [AFMC 1995]
(a) Electroplating (b) Electrorefining
(c) (a) and (b) both (d) None of these
30. Sodium is made by the electrolysis of a molten mixture of about 40% $NaCl$ and 60% $CaCl_2$ because [CBSE PMT 1995]
(a) $CaCl_2$ helps in conduction of electricity
(b) This mixture has a lower melting point than $NaCl$
(c) Ca^{++} can displace Na from $NaCl$
(d) Ca^{++} can reduce $NaCl$ to Na
31. Electrolysis is a process in which the cations and anions of the electrolyte are [MP PET 1995]
(a) Hydrated (b) Hydrolysed
(c) Charged (d) Discharged
32. Degree of ionisation of a solution depends upon [BHU 1998]
(a) Temperature (b) Nature of the electrolyte
(c) Nature of the solvent (d) None of these
33. Which of the following is non-electrolytes [KCET (Med.) 1999]
(a) $NaCl$ (b) $CaCl_2$
(c) $C_{12}H_{22}O_{11}$ (d) CH_3COOH
34. When a molten ionic hydride is electrolysed [JIPMER 1999]
(a) Hydrogen is liberated at the cathode
(b) Hydrogen is liberated at the anode
(c) There is no reaction
(d) H^- ions produced migrate to the cathode
35. During electrolysis, the species discharged at cathode are [AFMC 2000]
(a) Ions (b) Cation
(c) Anion (d) All of these
36. Electrolysis of molten anhydrous calcium chloride produces [AIIMS 2000]
(a) Calcium (b) Phosphorus
(c) Sulphur (d) Sodium
37. Which of the following properties of pure metal makes it more useful than the corresponding alloy [RPET 2000]
(a) It is harder than corresponding alloy
(b) It has high density
(c) It can be extracted easily
(d) It conducts heat and electricity easily
38. Which of the following liberate hydrogen on reaction with dilute H_2SO_4 [Roorkee 2000]
(a) Fe (b) Cu
(c) Al (d) Hg
39. Which one of the following material conducts electricity [Kerala (Med.) 2003]
(a) Diamond
(b) Crystalline sodium chloride
(c) Barium sulphate
(d) Fused potassium chloride
(e) Molten sulphur
40. Which of the following metals will give H_2 on reaction with $NaOH$
(a) Mg (b) Ba
(c) Ca (d) Sr
41. Which of the following is not a non electrolyte [J & K 2005]
(a) Acetic acid (b) Glucose
(c) Ethanol (d) Urea

Faraday's law of electrolysis

1. Amount of electricity that can deposit 108 gm of silver from $AgNO_3$ solution is [AFMC 1993; MP PMT 2004]
(a) 1 ampere (b) 1 coulomb
(c) 1 faraday (d) None of the above
2. When 9.65 coulombs of electricity is passed through a solution of silver nitrate (atomic weight of $Ag = 107.87$ taking as 108) the amount of silver deposited is [EAMCET 1992; KCET 2000]
(a) 10.8 mg (b) 5.4 mg
(c) 16.2 mg (d) 21.2 mg
3. Three faradays electricity was passed through an aqueous solution of iron (II) bromide. The weight of iron metal (at. wt. = 56) deposited at the cathode (in gm) is [EAMCET 1991]
(a) 56 (b) 84
(c) 112 (d) 168
4. A silver cup is plated with silver by passing 965 coulombs of electricity, the amount of silver deposited is [EAMCET 1990; MP PET 1994, 97]
(a) 9.89 g (b) 107.87 g

- (c) 1.0787 g (d) 1.002 g
5. The atomic weight of Al is 27. When a current of 5 Faradays is passed through a solution of Al^{+++} ions, the weight of Al deposited is [NCERT 1984]
(a) 27 gm (b) 36 gm
(c) 45 gm (d) 39 gm
6. An apparatus used for the measurement of quantity of electricity is known as a [BHU 1979]
(a) Calorimeter (b) Cathetometer
(c) Coulometer (d) Colorimeter
7. The unit of electrochemical equivalent is [EAMCET 1980]
(a) Gram (b) Gram/ampere
(c) Gram/coulomb (d) Coulomb/gram
8. A certain current liberated 0.504 gm of hydrogen in 2 hours. How many grams of copper can be liberated by the same current flowing for the same time in a copper sulphate solution [NCERT 1973, 77; CPMT 1979, 89; AIIMS 1987]
(a) 12.7 gm (b) 15.9 gm
(c) 31.8 gm (d) 63.5 gm
9. What weight of copper will be deposited by passing 2 Faradays of electricity through a cupric salt (Atomic weight of $Cu = 63.5$) [NCERT 1975; CPMT 1977, 79]
(a) 2.0 gm (b) 3.175 gm
(c) 63.5 gm (d) 127.0 gm
10. If the current is passed into the solution of an electrolyte [AIIMS 1979]
(a) Anions move towards anode, cations towards cathode
(b) Anions and cations both move towards anode
(c) Anions move towards cathode, cations towards anode
(d) No movement of ions takes place
11. Unit of Faraday is
(a) Ampere (b) Coulomb
(c) Coulomb $mole^{-1}$ (d) Coulomb Sec^{-1}
12. On passing 0.1 Faraday of electricity through aluminium chloride, the amount of aluminium metal deposited on cathode is ($Al = 27$) [MP PMT 1991]
(a) 0.9 gm (b) 0.3 gm
(c) 0.27 gm (d) 2.7 gm
13. Which of the following represents the first law of Faraday [MP PMT 1991]
(a) $E = mc^2$ (b) $E = hv$
(c) $m = ect$ (d) $PV = nRT$
14. 5 amperes is passed through a solution of zinc sulphate for 40 minutes. Find the amount of zinc deposited at the cathode [CBSE PMT 1996]
(a) 40.65 gm (b) 4.065 gm
(c) 0.4065 gm (d) 65.04 gm
15. In an electroplating experiment m g of silver is deposited, when 4 amperes of current flows for 2 minutes. The amount (in gms) of silver deposited by 6 amperes of current flowing for 40 seconds will be [MNR 1991]
(a) $4m$ (b) $m/2$
(c) $m/4$ (d) $2m$
16. On passing 3 ampere of electricity for 50 minutes, 1.8 gram metal deposits. The equivalent mass of metal is [MP PMT 1992]
(a) 20.5 (b) 25.8
(c) 19.3 (d) 30.7
17. The desired amount of charge for obtaining one mole of Al from Al^{3+}
(a) $3 \times 96500 C$ (b) 96500 C
- (c) $\frac{96500}{3} C$ (d) $\frac{96500}{2} C$
18. On passing one faraday of electricity through the electrolytic cells containing Ag^+ , Ni^{+2} and Cr^{+3} ions solution, the deposited Ag (At. wt. = 108), Ni (At. wt. = 59) and Cr (At. wt. = 52) is [AIIMS 1982]
- | | Ag | Ni | Cr |
|-----|----------|----------|----------|
| (a) | 108 gm | 29.5 gm | 17.3 gm |
| (b) | 108 gm | 59.0 gm | 52.0 gm |
| (c) | 108.0 gm | 108.0 gm | 108.0 gm |
| (d) | 108 gm | 117.5 gm | 166.0 gm |
19. One Faraday of electricity when passed through a solution of copper sulphate deposits [CPMT 1978]
(a) 1 mole of Cu (b) 1 gm atom of Cu
(c) 1 molecule of Cu (d) 1 gm equivalent of Cu
20. When 1 coulomb of charge is passed through electrolyte solution, then the mass deposited is equal to
(a) Equivalent weight
(b) Atomic weight
(c) Electrochemical equivalent
(d) Chemical equivalent
21. The platinum electrodes were immersed in a solution of cupric sulphate and electric current passed through the solution. After some time it was found that colour of copper sulphate disappeared with evolution of gas at the electrode. The colourless solution contains [NCERT 1984]
(a) Platinum sulphate (b) Copper hydroxide
(c) Copper sulphate (d) Sulphuric acid
22. On passing C ampere of electricity through a electrolyte solution for t second, m gram metal deposits on cathode. The equivalent weight E of the metal is [MP PMT 1990]
(a) $E = \frac{C \times t}{m \times 96500}$ (b) $E = \frac{C \times m}{t \times 96500}$
(c) $E = \frac{96500 \times m}{C \times t}$ (d) $E = \frac{C \times t \times 96500}{m}$
23. How many Faradays are required to generate one gram atom of magnesium from $MgCl_2$ [MADT Bihar 1982]
(a) 1 (b) 2
(c) 3 (d) 4
24. To deposit 0.6354 gm of copper by electrolysis of aqueous cupric sulphate solution, the amount of electricity required (in coulombs) is
(a) 9650 (b) 4825
(c) 3860 (d) 1930
25. In electrolysis of a fused salt, the weight of the deposit on an electrode will not depend on [CPMT 1973]
(a) Temperature of the bath
(b) Current intensity
(c) Electrochemical equivalent of ions
(d) Time for electrolysis
26. Faraday's laws of electrolysis will fail when [NCERT 1971]
(a) Temperature is increased
(b) Inert electrodes are used
(c) A mixture of electrolytes is used
(d) In none of these cases
27. According to the first law of Faraday, the weight of a substance discharge at the electrode is
(a) $W = ZQ$ (b) $W = eF$
(c) $W = \frac{Z}{F} It$ (d) $W = ZI$

28. When 0.04 faraday of electricity is passed through a solution of CaSO_4 , then the weight of Ca^{2+} metal deposited at the cathode is [BHU 1996]
 (a) 0.2 gm (b) 0.4 gm
 (c) 0.6 gm (d) 0.8 gm
29. A current 2.0 A is passed for 5 hours through a molten metal salt deposits 22 g of metal (At. wt. = 177). The oxidation state of the metal in the metal salt is [KCET 1996]
 (a) +1 (b) +2
 (c) +3 (d) +4
30. How many atoms of calcium will be deposited from a solution of CaCl_2 by a current of 25 milliamperes flowing for 60 seconds
 (a) 4.68×10^{18} (b) 4.68×10^{15}
 (c) 4.68×10^{12} (d) 4.68×10^9
31. On passing 0.5 faraday of electricity through NaCl , the amount of Cl deposited on cathode is [BHU 1997; RPET 1999]
 (a) 35.5 gm (b) 17.75 gm
 (c) 71 gm (d) 142 gm
32. What is the amount of chlorine evolved when 2 amperes of current is passed for 30 minutes in an aqueous solution of NaCl [BHU 1998; AIIMS 1999]
 (a) 66 g (b) 1.32 g
 (c) 33 g (d) 99 g
33. On passing a current through KCl solution, 19.5 g of potassium is deposited. If the same quantity of electricity is passed through a solution of aluminium chloride, the amount of aluminium deposited is [EAMCET 1997]
 (a) 4.5 g (b) 9.0 g
 (c) 13.5 g (d) 27 g
 (e) None is correct
34. Electrolysis rules of Faraday's states that mass deposited on electrode is proportional to [CBSE PMT 2000]
 (a) $m \propto I^2$ (b) $m \propto Q$
 (c) $m \propto Q^2$ (d) None of these
35. A current being passed for two hour through a solution of an acid liberating 11.2 litre of oxygen at NTP at anode. What will be the amount of copper deposited at the cathode by the same current when passed through a solution of copper sulphate for the same time [BVP 2003]
 (a) 16 g (b) 63 g
 (c) 31.5 g (d) 8 g
36. In a metal oxide, there is 20% oxygen by weight. Its equivalent weight is [Pb. PMT 2000]
 (a) 40 (b) 64
 (c) 72 (d) 32
37. On the basis of the information available from the reaction $\frac{4}{3}\text{Al} + \text{O}_2 \rightarrow \frac{2}{3}\text{Al}_2\text{O}_3$, $\Delta G = -827 \text{ kJ mol}^{-1}$ of O_2 the minimum *emf* required to carry out an electrolysis of Al_2O_3 is ($F = 96500 \text{ C mol}^{-1}$) [CBSE PMT 2003]
 (a) 8.56 V (b) 2.14 V
 (c) 4.28 V (d) 6.42 V
38. Then during electrolysis of a solution of AgNO_3 , 9650 coulombs of charge pass through the electroplating bath, the mass of silver deposited in the cathode will be [AIEEE 2003]
 (a) 1.08 g (b) 10.8 g
 (c) 21.6 g (d) 108 g
39. Total charge on 1 mole of a monovalent metal ion is equal to [DPMT 2001]
 (a) $9.65 \times 10^4 \text{ Coulomb}$ (b) $6.28 \times 10^{18} \text{ Coulomb}$
 (c) $1.6 \times 10^{-19} \text{ Coulomb}$ (d) None of these
40. When an electric current is passed through acidulated water 112 ml of hydrogen gas at N.T.P. collect at the cathode in 965 seconds. The current passed, in amperes is [MNR 1991; UPSEAT 2001]
 (a) 1.0 (b) 0.5
 (c) 0.1 (d) 2.0
41. How much chlorine will be liberated on passing one ampere current for 30 minutes through NaCl solution [BVP 2003]
 (a) 0.66 mole (b) 0.33 mole
 (c) 0.66 gm (d) 0.33 gm
42. The number of electrons involved in redox reactions when a Faraday of electricity is passed through an electrolyte in solution is [NCERT 1982; Pb. PMT 1999]
 (a) 6×10^{23} (b) 6×10^{-23}
 (c) 96500 (d) 8×10^{19}
43. Coulomb is equal to [Orissa JEE 2002]
 (a) ampere \times second (b) ampere \times minute
 (c) watt \times second (d) volt \times second
44. The energy required to release 1 electron from He^+ is..... [Orissa JEE 2002]
 (a) + 54.4 eV (b) - 13.6 eV
 (c) + 27.2 eV (d) Cannot be predicted
45. Faraday's laws of electrolysis are related to the [IIT 1983]
 (a) Atomic number of cation
 (b) Atomic number of anion
 (c) Equivalent weight of the electrolyte
 (d) Speed of the cation
46. The electric charge for electrode decomposition of one gram equivalent of a substance is [IIT 1984; KCET 1992]
 (a) One ampere per second
 (b) 96500 coulombs per second
 (c) One ampere for one hour
 (d) Charge on one mole of electrons
47. The number of electrons passing per second through a cross-section of copper wire carrying 10^{-6} amperes of current per second is found to be [EAMCET 1985]
 (a) 1.6×10^{-19} (b) 6×10^{-35}
 (c) 6×10^{-16} (d) 6×10^{12}
48. The electrolytic cells, one containing acidified ferrous chloride and another acidified ferric chloride are connected in series. The ratio of iron deposited at cathodes in the two cells when electricity is passed through the cells will be [CPMT 1989]
 (a) 3 : 1 (b) 2 : 1
 (c) 1 : 1 (d) 3 : 2
49. When 96500 coulomb of electricity is passed through a copper sulphate solution, the amount of copper deposited will be [MP PMT 1996]
 (a) 0.25 mol (b) 0.50 mol
 (c) 1.00 mol (d) 2.00 mol
50. During electrolysis of fused aluminium chloride 0.9 gm of aluminium was deposited on the cathode. The volume of chlorine liberated at the anode will be
 (a) 2.24 litres (b) 11.2 litres
 (c) 1.12 litres (d) 5.6 litres
51. Faraday has the dimensions of [MP PET 1995]
 (a) Coulombs
 (b) Coulomb equivalent
 (c) Coulomb per equivalent
 (d) Coulomb per degree Kelvin

52. The required charge for one equivalent weight of silver deposited on cathode is [Roorkee 1995] [AMU 1983; AFMC 1989; MP PET 2001]
(a) $9.65 \times 10^7 C$ (b) $9.65 \times 10^4 C$ (c) $9.65 \times 10^3 C$ (d) $9.65 \times 10^5 C$
53. 96500 coulombs of electric current liberates from $CuSO_4$ solution [MP PMT 1995] 63. The quantity of electricity needed to liberate 0.5 gram equivalent of an element is [CPMT 1988; MP PMT 1997]
(a) 63.5 gm Cu (b) 31.75 gm Cu (c) 96500 gm Cu (d) 100 gm Cu
54. A current of strength 2.5 amp was passed through $CuSO_4$ solution for 6 minutes 26 seconds. The amount of copper deposited is (Atomic weight of Cu = 63.5) (1 faraday = 96500 coulombs) [EAMCET 1989; MP PET 1994]
(a) 0.3175 g (b) 3.175 g (c) 0.635 g (d) 6.35 g
55. A certain quantity of electricity is passed through an aqueous solution of $AgNO_3$ and cupric salt solution connected in series. The amount of Ag deposited is 1.08 gm, the amount of copper deposited is (atomic weight of Cu = 63.5; Ag = 108) [EAMCET 1986]
(a) 0.6454 g (b) 6.354 g (c) 0.3177 g (d) 3.177 g
56. The number of electrons required to deposit 1 gm atom of aluminium (at. wt. = 27) from a solution of aluminium chloride will be (where N is Avogadro's number) [AIIMS 1992]
(a) 1 N (b) 2 N (c) 3 N (d) 4 N
57. Three faradays of electricity are passed through molten Al_2O_3 , aqueous solution of $CuSO_4$ and molten NaCl taken in different electrolytic cells. The amount of Al, Cu and Na deposited at the cathodes will be in the ratio of [BHU 1990]
(a) 1 mole : 2 mole : 3 mole (b) 3 mole : 2 mole : 1 mole (c) 1 mole : 1.5 mole : 3 mole (d) 1.5 mole : 2 mole : 3 mole
58. An electrolytic cell contains a solution of Ag_2SO_4 and have platinum electrodes. A current is passed until 1.6 gm of O_2 has been liberated at anode. The amount of silver deposited at cathode would be [CPMT 1971]
(a) 107.88 gm (b) 1.6 gm (c) 0.8 gm (d) 21.60 gm
59. The aqueous solution of which of the following decomposes on passing electric current [EAMCET 1973]
(a) Cane sugar (b) Urea (c) Methanol (d) Potassium iodide
60. The number of Faradays needed to reduce 4 gram equivalents of Cu^{++} to Cu metal will be [BHU 1981]
(a) 1 (b) 2 (c) 1/2 (d) 4
61. When electricity is passed through the solution of $AlCl_3$, 13.5 gm of Al are deposited. The number of Faraday must be [NCERT 1974; MP PET 1992; MP PMT 1994]
(a) 0.50 (b) 1.00 (c) 1.50 (d) 2.00
62. The value of one Faraday is (a) 95500 C mol⁻¹ (b) 96550 C mol⁻¹ (c) 96500 C mol⁻¹ (d) 98500 C mol⁻¹
64. The number of coulombs required for the deposition of 107.870 g of silver is [MP PET/PMT 1998]
(a) 96,500 (b) 48,250 (c) 1,93,000 (d) 10,000
65. When one of ampere current flows for 1 sec through a conductor, this quantity of electricity is known as [MP PMT 1993]
(a) Faraday (b) Coulomb (c) E.M.F. (d) Ohm
66. The mass deposited at an electrode is directly proportional to [CPMT 1974; AFMC 1987; MP PET 2000]
(a) Atomic weight (b) Equivalent weight (c) Molecular weight (d) Atomic number
67. From the solution of which of the following one faraday of electricity will liberate one gram atom of metal [MH CET 1999; MP PET 1993, 2000; AFMC 2000]
(a) NaCl (b) BaCl₂ (c) CuSO₄ (d) AlCl₃
68. On electrolysis, 1 mole of aluminium will be deposited from its molten salt by [MH CET 2000]
(a) 3 moles of electrons (b) 4 moles of electrons (c) 2 moles of electrons (d) 1 mole of electrons
69. The atomic weight of Fe is 56. The weight of Fe deposited from $FeCl_3$ solution by passing 0.6 Faraday of electricity is [MH CET 2000]
(a) 5.6 g (b) 11.2 g (c) 22.4 g (d) 33.6 g
70. 2.5 F of electricity are passed through a $CuSO_4$ solution. The number of gm equivalent of Cu deposited on anode is [CPMT 1973; DPMT 1982; MP PMT 2001]
(a) Zero (b) 1.25 (c) 2.5 (d) 5.0
71. The equivalent weight of a certain trivalent element is 20. Molecular weight of its oxide is [KCET 2003]
(a) 152 (b) 56 (c) 168 (d) 68
72. Silver is removed electrically from 200 ml of a 0.1 N solution of $AgNO_3$ by a current of 0.1 ampere. How long will it take to remove half of the silver from the solution [AMU 1999]
(a) 16 sec (b) 96.5 sec (c) 100 sec (d) 10 sec
73. In order to separate oxygen from one mole of H_2O the required quantity of coulomb would be [RPET 1999]
(a) 1.93×10^5 (b) 9.6×10^4 (c) 1.8 (d) 3.2
74. A current of 0.25 A is passed through $CuSO_4$ solution placed in voltameter for 45 minutes. The amount of Cu deposited on cathode is (At weight of Cu = 63.6) [BHU 2001]

- (a) 0.20 g (b) 0.22 g
(c) 0.25 g (d) 0.30 g [CPMT 1982; Pb.CET 2003]
75. Faraday constant [KCET (Med.) 2001]
(a) Is a numerical constant
(b) Depends on equivalent
(c) Depends upon the current passed
(d) Depends on the number of electrons
76. If 0.5 amp current is passed through acidified silver nitrate solution for 10 minutes. The mass of silver deposited on cathode, is (eq. wt. of silver nitrate = 108) [AFMC 2001]
(a) 0.235 g (b) 0.336 g
(c) 0.536 g (d) 0.636 g
77. The unit for the electric current is [KCET (Med.) 2001]
(a) Ohm (b) Volt
(c) Ampere (d) Coulomb
78. The quantity of electricity required to liberate 112 cm^3 of hydrogen at STP from acidified water is [KCET (Med.) 2001]
(a) 0.1 Faraday (b) 1 Faraday
(c) 965 Coulomb (d) 96500 Coulomb
79. Which solution will show highest resistance during the passage of current [BHU 2001]
(a) 0.05 N NaCl (b) 2 N NaCl
(c) 0.1 N NaCl (d) 1 N NaCl
80. 4 g of copper was dissolved in concentrated nitric acid. The copper nitrate solution on strong heating gave 5 g of its oxide. The equivalent weight of copper is [KCET 2004]
(a) 23 (b) 32
(c) 12 (d) 20
81. The amount of silver deposited by passing 241.25 coulomb of current through silver nitrate solution is [MHCET 2003]
(a) 2.7 g (b) 2.7 mg
(c) 0.27 g (d) 0.54 g
82. When 1F of electricity is passed through acidulated water, O_2 evolved is [MHCET 2004]
(a) 11.2 dm^3 (b) 5.6 dm^3
(c) 22.4 dm^3 (d) 1.0 dm^3
83. Charge required to liberate 11.5 g sodium is [AIIMS 1992; DCE 2002]
(a) 0.5 F (b) 0.1 F
(c) 1.5 F (d) 96500 coulombs
84. In the electrolysis of water, one Faraday of electrical energy would evolve [DCE 2004]
(a) One mole of oxygen (b) One g atom of oxygen
(c) 8 g of oxygen (d) 22.4 litres of oxygen
85. In a galvanic cell, the electrons flow from [KCET 2004]
(a) Anode to cathode through the solution
(b) Cathode to anode through the solution
(c) Anode to cathode through the external circuit
(d) Cathode to anode through the external circuit
86. An electric current is passed through silver nitrate solution using silver electrodes. 10.79 g of silver was found to be deposited on the cathode if the same amount of electricity is passed through copper sulphate solution using copper electrodes, the weight of copper deposited on the cathode is [Kerala PMT 2004]
(a) 6.4 g (b) 2.3 g
(c) 12.8 g (d) 1.6 g
(e) 3.2 g
87. The law of electrolysis were proposed by
(a) Kohlraush (b) Faraday
(c) Nernst (d) Berthelot
88. How many atoms of calcium will be deposited from a solution of CaCl_2 by a current 0.25 mA following for 60 seconds
(a) 4.68×10^{18} (b) 4.68×10^{15}
(c) 4.68×10^{12} (d) 4.68×10^9
89. The number of coulombs required to reduce 12.3 g of nitrobenzene to aniline [UPSEAT 2003]
(a) 115800 C (b) 5790 C
(c) 28950 C (d) 57900 C
90. During the process of electrolytic refining of copper, some metals present as impurity settle as 'anode mud'. These are [AIEEE 2005]
(a) Sn and Ag (b) Pb and Zn
(c) Ag and Au (d) Fe and Ni
91. A galvanic cell is set up from a zinc bar weighing 50g and 1.0 litre, 1.0M, CuSO_4 solution. How long would the cell run, assuming it delivers a steady current of 1.0 ampere [Roorkee 2000]
(a) 48 hrs (b) 41 hrs
(c) 21 hrs (d) 1 hr
92. On passing electric current through molten aluminium chloride, 11.2 litre of Cl_2 is liberated at NTP at anode. The quantity of aluminium deposited at cathode is (at. wt. of Al = 27)
(a) 9 g (b) 18 g
(c) 27 g (d) 36 g
93. An electric current is passed through silver voltameter connected to a water voltameter. The cathode of the silver voltameter weighed 0.108 g more at the end of the electrolysis. The volume of oxygen evolved at STP is [Kerala (Med.) 2003]
(a) 56 cm^3 (b) 550 cm^3
(c) 5.6 cm^3 (d) 11.2 cm^3
(e) 22.4 cm^3
94. During electrolysis of aqueous NaOH, 4 g of O_2 gas is liberated at NTP at anode, H_2 gas liberated at cathode is [CBSE PMT 1998]
(a) 2.8 litres (b) 5.6 litres
(c) 11.2 litres (d) 22.4 litres

Conductor and conductance

1. Which one of the following statements is correct [MP PET 1997]
(a) The oxidation number of oxygen in KO_2 is zero
(b) The specific conductance of an electrolyte solution decreases with increase in dilution
(c) Sn^{2+} oxidises Fe^{3+}
(d) $\text{Zn} / \text{ZnSO}_4$ is a reference electrode
2. In infinite dilutions, the equivalent conductances of Ba^{2+} and Cl^- are 127 and $76 \text{ ohm}^{-1} \text{ cm}^2 \text{ eqvt}^{-1}$. The equivalent conductivity of BaCl_2 at indefinite dilution is [CBSE 2000]
(a) 101.5 (b) 139.5

- (c) 203.5 (d) 279.5
3. The factor which is not affecting the conductivity of any solution is
(a) Dilution (b) Nature of electrolyte
(c) Temperature (d) None of these
4. Specific conductance of 0.1 M nitric acid is $6.3 \times 10^{-2} \text{ ohm}^{-1} \text{ cm}^{-1}$. The molar conductance of solution is
[Kerala (Med.) 2003]
(a) $630 \text{ ohm}^{-1} \text{ cm}^2 \text{ mole}^{-1}$ (b) $315 \text{ ohm}^{-1} \text{ cm}^2 \text{ mole}^{-1}$
(c) $100 \text{ ohm}^{-1} \text{ cm}^2 \text{ mole}^{-1}$ (d) $6300 \text{ ohm}^{-1} \text{ cm}^2 \text{ mole}^{-1}$
(e) $63.0 \text{ ohm}^{-1} \text{ cm}^2 \text{ mole}^{-1}$
5. The conductivity of strong electrolyte is [CPMT 2003]
(a) Increase on dilution slightly
(b) Decrease on dilution
(c) Does not change with dilution
(d) Depend upon density of electrolytes itself
6. If X is the specific resistance of the solution and M is the molarity of the solution, the molar conductivity of the solution is given by [Kurukshetra CEE 2002]
(a) $\frac{1000X}{M}$ (b) $\frac{1000}{MX}$
(c) $\frac{1000M}{X}$ (d) $\frac{MX}{1000}$
7. Conductivity (unit Siemen's) is directly proportional to area of the vessel and the concentration of the solution in it and is inversely proportional to the length of the vessel then the unit of the constant of proportionality is [AIEEE 2002]
(a) Sm mol^{-1} (b) $\text{Sm}^2 \text{ mol}^{-1}$
(c) $\text{S}^{-2} \text{ m}^2 \text{ mol}$ (d) $\text{S}^2 \text{ m}^2 \text{ mol}^{-2}$
8. If one end of a piece of a metal is heated, the other end becomes hot after some time. This is due to [CBSE PMT 1995]
(a) Energised electrons moving to the other part of the metal
(b) Resistance of the metal
(c) Mobility of atoms in the metal
(d) Minor perturbation in the energy of atoms
9. Conductivity of a solution is directly proportional to [KCET 1984]
(a) Dilution (b) Number of ions
(c) Current density (d) Volume of the solution
10. The increase in equivalent conductance of an electrolyte solution with dilution is due to the increase in [MP PMT 1996]
(a) Ionic attraction
(b) Molecular attraction
(c) Degree of association of the electrolyte
(d) Degree of ionisation of the electrolyte
11. Which of the following conducts electricity [AFMC 1995]
(a) Fused NaCl (b) CO_2
(c) Br_2 (d) Si
12. Which of the following shows electrical conduction [CBSE PMT 1999; AIIMS 1999]
(a) Potassium (b) Graphite
(c) Diamond (d) Sodium
13. The unit of equivalent conductivity is [CPMT 1999; BCECE 2005]
(a) ohm cm
(b) $\text{ohm}^{-1} \text{ cm}^2 (\text{gm equivalent})^{-1}$
(c) $\text{ohm cm}^2 (\text{gm equivalent})$
(d) S cm^{-2}
14. It has been observed that gaseous hydrogen chloride is a very poor conductor of electricity but a solution of hydrogen chloride gas in water is a good conductor of electricity. This is due to the fact that [NCERT 1976]
(a) Water is good conductor of electricity
(b) Hydrogen chloride gas in water solution ionizes
(c) A gas is non-conductor but a liquid conducts electricity
(d) Gas does not obey Ohm's law whereas solution does
15. Electrolytic conduction differs from metallic conduction in that in the case of electrolytic conduction [KCET 1987; Bihar CEE 1992]
(a) The resistance increases with increasing temperature
(b) The resistance decreases with increasing temperature
(c) The flow of current does not generate heat
(d) The resistance is independent of the length of the conductor
16. The electrolytic conductance is a direct measure of [KCET 1990; CPMT 2003]
(a) Resistance (b) Potential
(c) Concentration (d) Dissociation
17. Conductivity of a strong electrolyte [KCET 1993]
(a) Increases on dilution
(b) Does not change considerably on dilution
(c) Decreases on dilution
(d) Depends on density
18. Which of the following statements is not applicable to electrolytic conductors [AIIMS 1991]
(a) New products show up at the electrodes
(b) Ions are responsible for carrying the current
(c) Show a positive temperature coefficient for conductance
(d) A single stream of electrons flows from cathode to anode
19. Which one is not a conductor of electricity [RPET 1999]
(a) NaCl (aqueous) (b) NaCl (solid)
(c) NaCl (molten) (d) Ag metal
20. Solid sodium chloride is bad conductor of electricity because
(a) It contains only molecules
(b) It does not possess ions
(c) The ions present in it are not free to move
(d) It does not contain free molecules
21. Which of the following is a poor conductor of electricity [EAMCET 1992]
(a) CH_3COONa (b) $\text{C}_2\text{H}_5\text{OH}$
(c) NaCl (d) KOH
22. The molar conductivity is maximum for the solution of concentration [DCE 2002]
(a) 0.001 M (b) 0.005 M
(c) 0.002 M (d) 0.004 M
23. The unit of molar conductivity is [DCE 2002]
(a) $\Omega^{-1} \text{ cm}^{-2} \text{ mol}^{-1}$ (b) $\Omega \text{ cm}^{-2} \text{ mol}^{-1}$
(c) $\Omega^{-1} \text{ cm}^2 \text{ mol}^{-1}$ (d) $\Omega \text{ cm}^2 \text{ mol}$
24. The highest electrical conductivity of the following aqueous solutions is of [AIEEE 2005]
(a) 0.1 M acetic acid (b) 0.1 M chloroacetic acid
(c) 0.1 M fluoroacetic acid (d) 0.1 M difluoroacetic acid
25. Given $l/a = 0.5 \text{ cm}^{-1}$, $R = 50 \text{ ohm}$, $N = 1.0$. The equivalent conductance of the electrolytic cell is [Orissa JEE 2005]
(a) $10 \text{ ohm}^{-1} \text{ cm}^2 \text{ gmeq}^{-1}$ (b) $20 \text{ ohm}^{-1} \text{ cm}^2 \text{ gmeq}^{-1}$
(c) $300 \text{ ohm}^{-1} \text{ cm}^2 \text{ gmeq}^{-1}$ (d) $100 \text{ ohm}^{-1} \text{ cm}^2 \text{ gmeq}^{-1}$

26. If equivalent conductance of 1M benzoic acid is $12.8 \text{ ohm}^{-1} \text{ cm}^2$ and if the conductance of benzoate ion and H^+ ion are 42 and $288.42 \text{ ohm}^{-1} \text{ cm}^2$ respectively, its degree of dissociation is
(a) 39% (b) 3.9%
(c) 0.35% (d) 0.039%
27. The unit ohm^{-1} is used for [J & K 2005]
(a) Molar conductivity (b) Equivalent conductivity
(c) Specific conductivity (d) Conductivity
- Cell constant and Electrochemical Cells**
1. When electric current is passed through a cell having an electrolyte, the positive ions move towards the cathode and the negative ions towards the anode. If the cathode is pulled out of the solution [AIIMS 1980]10.
(a) The positive and negative ions will move towards the anode
(b) The positive ions will start moving towards the anode, the negative ions will stop moving
(c) The negative ions will continue to move towards the anode and the positive ions will stop moving
(d) The positive and negative ions will start moving randomly
2. If the half cell reaction $A + e^- \rightarrow A^-$ has a large negative reduction potential, it follows that [MNR 1992; UPSEAT 2000, 02]
(a) A is readily reduced (b) A is readily oxidised
(c) A^- is readily reduced (d) A^- is readily oxidised
3. Mark the false statement [MP PET 1997]
(a) A salt bridge is used to eliminate liquid junction potential
(b) The Gibbs free energy change, ΔG is related with electromotive force (E), as $\Delta G = -nFE$
(c) Nernst equation for single electrode potential is
$$E = E^\circ - \frac{RT}{nF} \ln a_{M^{n+}}$$

(d) The efficiency of a hydrogen oxygen fuel cell is 23%
4. The specific conductance of a 0.1 N KCl solution at 23°C is $0.012 \text{ ohm}^{-1} \text{ cm}^{-1}$. The resistance of cell containing the solution at the same temperature was found to be 55 ohm. The cell constant will be [CBSE PMT 1999, 2000; KCET 2001]
(a) 0.142 cm^{-1} (b) 0.66 cm^{-1}
(c) 0.918 cm^{-1} (d) 1.12 cm^{-1}
5. Which of the following reactions occurs at the cathode of a common dry cell [NCERT 1978]
(a) $Mn \rightarrow Mn^{2+} + 2e^-$
(b) $2MnO_2 + Zn^{2+} + 2e^- \rightarrow ZnMn_2O_4$
(c) $2ZnO_2 + Mn^{2+} + 2e^- \rightarrow MnZn_2O_4$
(d) $Zn \rightarrow Zn^{2+} + 2e^-$
6. In Cu – Zn cell [BHU 1981]
(a) Reduction occurs at the copper cathode
(b) Oxidation occurs at the copper cathode
(c) Reduction occurs at the anode
(d) Chemical energy is converted to light energy
7. Which of the following reaction is used to make a fuel cell [AIIMS 2003]
(a) $Cd(s) + 2Ni(OH)_3(s) \rightarrow CdO(s) + 2Ni(OH) + H_2O(l)$
(b) $Pb(s) + PbO_2(s) + 2H_2SO_4(aq) \rightarrow 2PbSO_4(s) + 2H_2O(l)$
(c) $2H_2(g) + O_2(g) \rightarrow 2H_2O(l)$
(d) $2Fe(s) + O_2(g) + 4H^+(aq) \rightarrow 2Fe^{2+}(aq) + 2H_2O(l)$
8. When lead storage battery is charged [MP PET 2003]
(a) $PbSO_4$ is reduced
(b) H_2SO_4 is regenerated
(c) $PbSO_4$ is deposited on lead electrode
(d) Lead is deposited on lead electrode
9. When lead storage battery is charged [MP PET 1993; MP PMT 2000]
(a) Lead dioxide dissolves
(b) Sulphuric acid is regenerated
(c) The lead electrode becomes coated with lead sulphate
(d) The amount of sulphuric acid decreases
- The electrolytic decomposition of dilute sulphonic acid with platinum electrode in cathodic reaction is [MNR 1988; UPSEAT 1999, 2002]
(a) Oxidation
(b) Reduction
(c) Oxidation and reduction both
(d) Neutralisation
11. Which colourless gas evolves, when NH_4Cl reacts with zinc in a dry cell battery [Orissa JEE 2003]
(a) NH_4 (b) N_2
(c) H_2 (d) Cl_2
12. Which of the substances Na, Hg, S, Pt and graphite can be used as electrodes in electrolytic cells having aqueous solutions
(a) Na, Pt and graphite (b) Na and Hg
(c) Pt and graphite only (d) Na and S only
13. In electrolysis of dilute H_2SO_4 using platinum electrodes [DPMT 1983; IIT 1983; Kurukshetra CET 2002; AFMC 2005]
(a) H_2 is evolved at cathode
(b) NH_3 is produced at anode
(c) Cl_2 is obtained at cathode
(d) O_2 is produced
14. For cell reaction, $Zn + Cu^{2+} \rightarrow Zn^{2+} + Cu$, cell representation is
(a) $Zn | Zn || Cu | Cu$ (b) $Cu | Cu || Zn | Zn$
(c) $Cu | Zn || Zn | Cu$ (d) $Cu | Zn || Zn | Cu$
15. Which one is not called a anode reaction from the following
(a) $Cl^- \rightarrow \frac{1}{2} Cl_2 + e^-$ (b) $Cu \rightarrow Cu^{++} + 2e^-$
(c) $Hg^+ \rightarrow Hg^{++} + e^-$ (d) $Zn^{2+} + 2e^- \rightarrow Zn$
16. A cell from the following which converts electrical energy into chemical energy
(a) Dry cell (b) Electrochemical cell
(c) Electrolytic cell (d) None of these
17. In the cell $Zn | Zn^{2+} || Cu^{2+} | Cu$, the negative electrode is [MP PMT 1995]
(a) Cu (b) Cu^{2+}
(c) Zn (d) Zn^{2+}
18. Which of the following statements is correct? Galvanic cell converts [KCET 1991]
(a) Chemical energy into electrical energy
(b) Electrical energy into chemical energy
(c) Metal from its elemental state to the combined state

- (d) Electrolyte into individual ions
19. Hydrogen-oxygen fuel cells are used in space-craft to supply [MP PMT 1993; MP PET 1999] [Pb. PMT 1999; KCET 1999]
 (a) Power for heat and light
 (b) Power for pressure
 (c) Oxygen
 (d) Water
20. The standard cell potential of $Zn|Zn^{2+}(aq)||Cu^{2+}(aq)|Cu$ cell is 1.10 V. The maximum work obtained by this cell will be
 (a) 106.15 kJ (b) -212.30 kJ
 (c) -318.45 kJ (d) -424.60 kJ
21. The relationship between standard reduction potential of cell and equilibrium constant is shown by [MP PET 2002]
 (a) $E_{cell}^0 = \frac{n}{0.059} \log K_c$ (b) $E_{cell}^0 = \frac{0.059}{n} \log K_c$
 (c) $E_{cell}^0 = 0.059 n \log K_c$ (d) $E_{cell}^0 = \frac{\log K_c}{n}$
22. Consider the Galvanic cell $Zn^{\ominus} | ZnSO_4 || CuSO_4 | Cu^{\oplus}$ the reaction at cathode is [AMU 2000]
 (a) $Zn^{2+} + 2e^- \rightarrow Zn$
 (b) $Cu^{2+} + 2e^- \rightarrow Cu$
 (c) $Cu^{2+} + Zn \rightarrow Cu + Zn^{2+}$
 (d) $Zn^{2+} + Cu \rightarrow Zn + Cu^{2+}$
23. The cell reaction $Cu + 2Ag^+ \rightarrow Cu^{+2} + Ag$ is best represented by [AMU 2000]
 (a) $Cu_{(s)} | Cu^{+2}(aq) || Ag^+(aq) | Ag_{(s)}$
 (b) $Pt | Cu^{+2} || Ag^+(aq) | Ag_{(s)}$
 (c) $Cu^{+2} | Cu || Pt | Ag$
 (d) None of the above representations
24. $Zn_{(s)} | Zn^{2+}(aq) || Cu^{2+}(aq) | Cu_{(s)}$ is [Kerala (Engg.) 2002]
 (anode) (cathode)
 (a) Weston cell (b) Daniel cell
 (c) Calomel cell (d) Faraday cell
 (e) Standard cell
25. The specific conductance of a solution is $0.2 \text{ ohm}^{-1} \text{ cm}^{-1}$ and conductivity is 0.04 ohm^{-1} . The cell constant would be
 (a) 1 cm^{-1} (b) 0 cm^{-1}
 (c) 5 cm^{-1} (d) 0.2 cm^{-1}
26. If the conductance and specific conductance of a solution is one then its cell constant would be [RPET 1999]
 (a) 1 (b) Zero
 (c) 0.5 (d) 4
27. Saturated solution of KNO_3 is used to make 'salt-bridge' because
 (a) Velocity of K^+ is greater than that of NO_3^-
 (b) Velocity of NO_3^- is greater than that of K^+
 (c) Velocities of both K^+ and NO_3^- are nearly the same
 (d) KNO_3 is highly soluble in water
28. In balancing the half reaction $S_2O_3^{2-} \rightarrow S_{(s)}$ the number of electrons that must be added is [DPMT 2000]
 (a) 4 on the left (b) 3 on the right
 (c) 2 on the left (d) 2 on the right
29. Which one of the following statement is true for an electrochemical cell [Pb. PMT 1999; KCET 1999]
 (a) H_2 is cathode and Cu is anode
 (b) H_2 is anode and Cu is cathode
 (c) Reduction occurs at H_2 electrode
 (d) Oxidation occurs at Cu electrode
30. In the reaction [MP PET 2002]
 $Cu(s) + 2Ag^+(aq) \rightarrow Cu^{2+}(aq) + 2Ag(s)$
 The reduction half-cell reaction is [AIIMS 1997]
 (a) $Cu + 2e^- \rightarrow Cu^{2-}$ (b) $Cu - 2e^- \rightarrow Cu^{2+}$
 (c) $Ag^+ + e^- \rightarrow Ag$ (d) $Ag - e^- \rightarrow Ag^+$
31. Which of the following statements about galvanic cell is incorrect [JIPMER 1997]
 (a) Anode is positive
 (b) Oxidation occurs at the electrode with lower reduction potential
 (c) Cathode is positive
 (d) Reduction occurs at cathode
32. The molar conductances of $NaCl$, HCl and CH_3COONa at infinite dilution are 126.45, 426.16 and $91 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$ respectively. The molar conductance of CH_3COOH at infinite dilution is [CBSE PMT 1997]
 (a) $201.28 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$
 (b) $390.71 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$
 (c) $698.28 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$
 (d) $540.48 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$
33. The electrodes of a conductivity cell are 3 cm apart and have a cross-sectional area of 4 cm^2 . The cell constant of the cell (in cm^{-1}) is
 (a) 4×3 (b) $4/3$
 (c) $3/4$ (d) $9/4$
34. The anode half-reaction occurring during the discharge of a lead storage battery is
 (a) $Pb(s) + SO_2 + O_2 \rightarrow PbSO_4(s)$
 (b) $Pb(s) + SO_4^{2-}(aq) \rightleftharpoons PbSO_4(s) + 2e^-$
 (c) $PbO_2(s) + 4H^+(aq) + 2e^- + SO_4^{2-}(aq) \rightleftharpoons PbSO_4(s) + 2H_2O$ [RPET 1999]
 (d) $Pb^{2+}(aq) + SO_4^{2-}(aq) \rightarrow PbSO_4(s)$
35. The unit of cell constant is [MP PET 1996]
 (a) $\text{ohm}^{-1} \text{ cm}^{-1}$ (b) ohm cm
 (c) cm (d) cm^{-1}
36. In dry cell the reaction which takes place at the zinc anode is [MP PET 1996]
 (a) $Zn^{2+} + 2e^- \rightarrow Zn(s)$
 (b) $Zn(s) \rightarrow Zn^{2+} + 2e^-$
 (c) $Mn^{2+} + 2e^- \rightarrow Mn(s)$
 (d) $Mn(s) \rightarrow Mn^{+} + e^- + 1.5V$
37. The chemical reaction taking place at the anode of a cell is [MP PET 1996]
 (a) Ionisation (b) Reduction

- (c) Oxidation (d) Hydrolysis
38. Which of the following reactions occurs at the cathode during the charging of a lead storage battery [Manipal MEE 1995; MP PET 2002]
- (a) $Pb^{2+} + 2e^- \rightarrow Pb$
 (b) $Pb^{2+} + SO_4^{2-} \rightarrow PbSO_4$
 (c) $Pb \rightarrow Pb^{2+} + 2e^-$
 (d) $PbSO_4 + 2H_2O \rightarrow 2PbO_2 + 4SO_4^{2-} + 2e^-$
39. A depolarizer used in dry cell batteries is [NCERT 1981]
- (a) Ammonium chloride (b) Manganese dioxide
 (c) Potassium hydroxide (d) Sodium phosphate
40. When a lead storage battery is discharged [IIT 1987; MP PMT 2004]
- (a) SO_2 is evolved
 (b) Lead sulphate is consumed
 (c) Lead is formed
 (d) Sulphuric acid is consumed
41. In electroplating, the article to be electroplated serves as [AMU 1982, 83]
- (a) Cathode (b) Electrolyte
 (c) Anode (d) Conductor
42. The position of some metals in the electrochemical series in decreasing electropositive character is given as $Mg > Al > Zn > Cu > Ag$. What will happen, if a copper spoon is used to stir a solution of aluminium nitrate [NCERT 1977]
- (a) The spoon will get coated with Al
 (b) An alloy of Cu and Al is formed
 (c) The solution becomes blue
 (d) There is no reaction
43. In a electrochemical cell [AFMC 1989]
- (a) Potential energy changes into kinetic energy
 (b) Kinetic energy changes into potential energy
 (c) Chemical energy changes into electrical energy
 (d) Electrical energy changes into chemical energy
44. In galvanic cell, the salt bridge is used to [MP PMT 2002]
- (a) Complete the circuit
 (b) Reduce the electric resistance in the cell
 (c) Separate cathode from anode
 (d) Carry salts for the chemical reaction
45. If a strip of Cu metal is placed in a solution of ferrous sulphate [NCERT 1974; CPMT 1977; MP PET 2000]
- (a) Copper will precipitate out
 (b) Iron will precipitate out
 (c) Copper will dissolve
 (d) No reaction will take place
46. Which of the following is not used to construct salt bridge [MP PET 2003]
- (a) CH_3COOK (b) KCl
 (c) NH_4NO_3 (d) KNO_3
47. The reference electrode is made by using [MP PMT 2002]
- (a) $ZnCl_2$ (b) $CuSO_4$
 (c) $HgCl_2$ (d) Hg_2Cl_2
48. In a hydrogen – oxygen fuel cell, combustion of hydrogen occurs to
- (a) Produce high purity water
 (b) Create potential difference between the two electrodes
 (c) Generate heat
 (d) Remove adsorbed oxygen from electrode surfaces
49. $\lambda_{ClCH_2COONa} = 224 \text{ ohm}^{-1} \text{ cm}^2 \text{ gmeq}^{-1}$,
 $\lambda_{NaCl} = 38.2 \text{ ohm}^{-1} \text{ cm}^2 \text{ gmeq}^{-1}$,
 $\lambda_{HCl} = 203 \text{ ohm}^{-1} \text{ cm}^2 \text{ gmeq}^{-1}$,
 What is the value of λ_{ClCH_2COOH} [JEE Orissa 2004]
- (a) $288.5 \text{ ohm}^{-1} \text{ cm}^2 \text{ gmeq}^{-1}$
 (b) $289.5 \text{ ohm}^{-1} \text{ cm}^2 \text{ gmeq}^{-1}$
 (c) $388.5 \text{ ohm}^{-1} \text{ cm}^2 \text{ gmeq}^{-1}$
 (d) $59.5 \text{ ohm}^{-1} \text{ cm}^2 \text{ gmeq}^{-1}$
50. Which of the following statement is true for the electrochemical Daniel cell [AIIMS 2004]
- (a) Electrons flow from copper electrode to zinc electrode
 (b) Current flows from zinc electrode to copper electrode
 (c) Cations move toward copper electrode which is cathode
 (d) Cations move toward zinc electrode
51. Which of the following statement is true for an electrochemical cell
- (a) H_2 is cathode and Cu is anode
 (b) H_2 is anode and Cu is cathode
 (c) Reduction occurs at H_2 electrode
 (d) Oxidation occurs at Cu electrode
52. Which of the following statements are true for an fuel cells [DPMT 2004]
- (a) They run till the reactants are active
 (b) They are free from pollution
 (c) They are more efficient
 (d) All of the above
53. For gold plating, the electrolyte used is [Pb.CET 2004]
- (a) $AuCl_3$ (b) $HAuCl_4$
 (c) $k[Au(CN)_2]$ (d) None of these
54. The acid used in lead storage battery is [Pb.CET 2003]
- (a) H_2SO_4 (b) H_3PO_4
 (c) HCl (d) HNO_3
55. At $25^\circ C$ specific conductivity of a normal solution of KCl is 0.002765 mho . The resistance of cell is 400 ohms . The cell constant is [Pb.PMT 2004]
- (a) 0.815 (b) 1.016
 (c) 1.106 (d) 2.016
56. Which of the following is used widely in the manufacture of lead storage battery [BHU 2004]
- (a) Arsenic (b) Lithium
 (c) Bismuth (d) Antimony
57. The chemical reaction, $2AgCl(s) + H_2(g) \rightarrow 2HCl(aq) + 2Ag(s)$ taking place in a galvanic cell is represented by the notation [AIIMS 2005]
- (a) $Pt | H_2(g), 1 \text{ bar} | 1M KCl(aq) | AgCl(s) | Ag(s)$
 (b) $Pt | H_2(g), 1 \text{ bar} | 1M HCl(aq) || 1M Ag^+(aq) | Ag(s)$ [AIIMS 2004]

- (c) $Pt(s) | H_2(g), 1\text{bar} | 1M\ HCl(aq) | AgCl(s) | Ag(s)$
 (d) $Pt(s) | H_2(g), 1\text{bar} | 1M\ HCl(aq) | Ag(s) | AgCl(s)$
58. If the Zn^{2+} / Zn electrode is diluted to 100 times then the change in e.m.f. [DPMT 2005]
 (a) Increase of 59 mV
 (b) Decrease of 59 mV
 (c) Increase of 29.5 mV
 (d) Decrease of 29.5 mV
59. If hydrogen electrode dipped in 2 solution of $pH = 3$ and $pH = 6$ and salt bridge is connected the e.m.f. of resulting cell is [DPMT 2005]
 (a) 0.177 V (b) 0.3 V
 (c) 0.052 V (d) 0.104 V
60. The tendency of an electrode to lose electrons is known as [J & K 2005]
 (a) Electrode potential (b) Reduction potential
 (c) Oxidation potential (d) e.m.f.
61. When electric current is supplied through an ionic hydride of fused state, then [Kerala CET 2005]
 (a) Hydrogen is obtained at anode
 (b) Hydrogen is obtained at cathode
 (c) No change
 (d) Hydride ion moves towards cathode
 (e) hydride ion present in solution
- Electrode potential, E_{Cell} , Nernt equation and ECS**
1. The hydrogen electrode is dipped in a solution of $pH = 3$ at 25°C . The potential of the cell would be (the value of $2.303RT/F$ is 0.059 V) [KCET 1993, 2005]
 (a) 0.177 V (b) -0.177 V
 (c) 0.087 V (d) 0.059 V
2. The standard electrode potentials of Zn^{2+} / Zn and Ag^+ / Ag are -0.763 V and $+0.799\text{ V}$ respectively. The standard potential of the cell is [KCET 1993]
 (a) 1.56 V (b) 0.036 V
 (c) -1.562 V (d) 0.799 V
3. The standard reduction potentials at 298 K for the following half reactions are given against each
 $Zn^{2+}(aq) + 2e \rightleftharpoons Zn(s); -0.762$
 $Cr^{3+}(aq) + 3e \rightleftharpoons Cr(s); -0.740$
 $2H^+(aq) + 2e \rightleftharpoons H_2(g); 0.00$
 $Fe^{3+}(aq) + e \rightleftharpoons Fe^{2+}(aq); 0.770$
 Which is the strongest reducing agent [IIT 1981; MP PET/PMT 1988; MP PMT 1989; MH CET 2001]
 (a) $Zn(s)$ (b) $Cr(s)$
 (c) $H_2(g)$ (d) $Fe^{2+}(aq)$
4. When Zn piece is kept in $CuSO_4$ solution, the copper get precipitated due to standard potential of zinc is
 (a) $>$ copper (b) $<$ copper
 (c) $>$ sulphate (d) $<$ sulphate
5. Which of the following metal does not react with the solution of copper sulphate [CPMT 1999]
 (a) Mg (b) Fe
 (c) Zn (d) Ag
6. A solution containing one mole per litre of each $Cu(NO_3)_2$, $AgNO_3$, $Hg_2(NO_3)_2$ and $Mg(NO_3)_2$, is being electrolysed by using inert electrodes. The values of standard electrode potentials in volts (reduction potentials) are $Ag / Ag^+ = +0.80$, $2Hg / Hg_2^{2+} = +0.79$, $Cu / Cu^{2+} = +0.34$, $Mg / Mg^{2+} = -2.37$ with increasing voltage, the sequence of deposition of metals on the cathode will be [IIT 1984; AMU 1999; Kerala PMT 2004]
 (a) Ag, Hg, Cu, Mg (b) Mg, Cu, Hg, Ag
 (c) Ag, Hg, Cu (d) Cu, Hg, Ag
7. The standard reduction electrode potentials of four elements are
 $A = -0.250\text{ V}$ $B = -0.136\text{ V}$
 $C = -0.126\text{ V}$ $D = -0.402\text{ V}$
 The element that displaces A from its compounds is
 (a) B (b) C
 (c) D (d) None of these
8. The standard oxidation potential of zinc and silver in water at 298 K are
 $Zn(s) \rightarrow Zn^{2+} + 2e^-; E = 0.76\text{ V}$
 $Ag(s) \rightarrow Ag^{2+} + 2e^-; E = -0.80\text{ V}$
 Which of the following reactions actually take place [NCERT 1983, 84; KCET 2003]
 (a) $Zn(s) + 2Ag^+(aq) \rightarrow Zn^{2+}(aq) + 2Ag(s)$
 (b) $Zn^{2+}(aq) + 2Ag(s) \rightarrow 2Ag^+(aq) + Zn(s)$
 (c) $Zn(s) + Ag(s) \rightarrow Zn^{2+}(aq) + Ag^+(aq)$
 (d) $Zn^{2+}(aq) + Ag^+(aq) \rightarrow Zn(s) + Ag(s)$
9. Beryllium is placed above magnesium in the second group. Beryllium dust, therefore when added to $MgCl_2$ solution will [CPMT 1977]
 (a) Have no effect
 (b) Precipitate Mg metal
 (c) Precipitate MgO
 (d) Lead to dissolution of Be metal
10. The name of equation showing relation between electrode potential (E) standard electrode potential (E°) and concentration of ions in solution is
 (a) Kohlrausch's equation (b) Nernst's equation
 (c) Ohm's equation (d) Faraday's equation
11. The correct representation of Nernst's equation is
 (a) $E_{M^{n+}/M} = E^\circ_{M^{n+}/M} + \frac{0.0591}{n} \log(M^{n+})$
 (b) $E_{M^{n+}/M} = E^\circ_{M^{n+}/M} - \frac{0.0591}{n} \log(M^{n+})$

- (c) $E_{M^{n+}/M} = E_{M^{n+}/M}^{\circ} + \frac{n}{0.0591} \log(M^{n+})$
 (d) None of the above
12. Standard electrode potential of *NHE* at 298 K is
 (a) 0.05 V (b) 0.1 V
 (c) 0.00 V (d) 0.11 V
13. When a copper wire is placed in a solution of $AgNO_3$, the solution acquires blue colour. This is due to the formation of
 [Roorkee 1989]
 (a) Cu^{2+} ions
 (b) Cu^{+} ions
 (c) Soluble complex of copper with $AgNO_3$
 (d) Cu^{-} ion by the reduction of Cu
14. Consider the reaction $M_{(aq)}^{n+} + ne^{-} \rightarrow M_{(s)}$. The standard reduction potential values of the elements M_1, M_2 and M_3 are $-0.34V, -3.05V$ and $-1.66V$ respectively. The order of their reducing power will be
 [NCERT 1990]
 (a) $M_1 > M_2 > M_3$ (b) $M_3 > M_2 > M_1$
 (c) $M_1 > M_3 > M_2$ (d) $M_2 > M_3 > M_1$
15. $E^{\circ} = \frac{RT}{nF} \ln K_{eq}$. This is called
 [CPMT 1988; MP PET 2000]
 (a) Gibb's equation
 (b) Gibb's-Helmholtz equation
 (c) Nernst's equation
 (d) Vander Waal's equation
16. Four alkali metals A, B, C and D are having respectively standard electrode potential as $-3.05, -1.66, -0.40$ and 0.80 . Which one will be the most reactive
 [MP PMT/PET 1988; CPMT 1983; MNR 1993; UPSEAT 2002]
 (a) A (b) B
 (c) C (d) D
17. Which one of the following metals cannot evolve H_2 from acids or H_2O or from its compounds
 [MP PET/PMT 1988; CPMT 1996; AFMC 1998, 99; Pb. PET 1999; BVP 2003]
 (a) Hg (b) Al
 (c) Pb (d) Fe
18. Which one of the following reaction is not possible
 [MP PMT 1991]
 (a) $Fe + H_2SO_4 \rightarrow FeSO_4 + H_2$
 (b) $Cu + 2AgNO_3 \rightarrow Cu(NO_3)_2 + 2Ag$
 (c) $2KBr + I_2 \rightarrow 2KI + Br_2$
 (d) $CuO + H_2 \rightarrow Cu + H_2O$
19. When a rod of metal A is dipped in an aqueous solution of metal B (concentration of B^{2+} ion being 1M) at $25^{\circ}C$, the standard electrode potentials are $A^{2+}/A = -0.76$ volts, $B^{2+}/B = +0.34$ volts [KCET 1992]
 (a) A will gradually dissolve
 (b) B will deposit on A
 (c) No reaction will occur
 (d) Water will decompose into H_2 and O_2
20. The reaction $Zn^{2+} + 2e^{-} \rightarrow Zn$ has a standard potential of $-0.76V$. This means [KCET 1992]
 (a) Zn can't replace hydrogen from acids
 (b) Zn is a reducing agent
 (c) Zn is an oxidising agent
 (d) Zn^{2+} is a reducing agent
21. $2H^{+}(aq) + 2e^{-} \rightarrow H_2(g)$. The standard electrode potential for the above reaction is (in volts) [CPMT 1988]
 (a) 0 (b) +1
 (c) -1 (d) None of these
22. K, Ca and Li metals may be arranged in the decreasing order of their standard electrode potentials as [CPMT 1990]
 (a) K, Ca, Li (b) Ca, K, Li
 (c) Li, Ca, K (d) Ca, Li, K
23. The correct order of chemical reactivity with water according to electrochemical series [MP PMT 1991]
 (a) $K > Mg > Zn > Cu$ (b) $Mg > Zn > Cu > K$
 (c) $K > Zn > Mg > Cu$ (d) $Cu > Zn > Mg > K$
24. EMF of cell $Ni | Ni^{2+}(1.0M) || Au^{3+}(1.0M) | Au$ (Where E° for $Ni^{2+} | Ni$ is $-0.25V$; E° for $Au^{3+} | Au$ is $1.50V$) is [MP PET 1993; MP PMT 2000]
 (a) $+1.25V$ (b) $-1.75V$
 (c) $+1.75V$ (d) $+4.0V$
25. Oxidation and reduction take place in a cell, then its electromotive force will be [RPET 1999]
 (a) Positive (b) Negative
 (c) Zero (d) Stable
26. For a spontaneous reaction the ΔG , equilibrium constant (K) and E_{Cell}° will be respectively [AIEEE 2005]
 (a) $-ve, >1, +ve$ (b) $+ve, >1, -ve$
 (c) $-ve, <1, -ve$ (d) $-ve, >1, -ve$
27. The reference electrode is made from which of the following [MP PET/PMT 1988]
 (a) $ZnCl_2$ (b) $CuSO_4$
 (c) Hg_2Cl_2 (d) $HgCl_2$
28. The charge over anode in a galvanic cell is
 (a) Negative
 (b) Positive
 (c) No charge
 (d) Sometimes negative and sometimes positive
29. The standard electrode potential for the two electrode A^{+}/A and B^{+}/B are respectively $0.5V$ and $0.75V$. The emf of the given cell $A | A^{+}(a=1) || B^{+}(a=1) | B$ will be
 (a) $1.25V$ (b) $-1.25V$
 (c) $-0.25V$ (d) $0.25V$
30. The standard reduction potential for Li^{+}/Li ; Zn^{2+}/Zn ; H^{+}/H_2 and Ag^{+}/Ag is $-3.05, -0.762, 0.00$ and $+0.80V$. Which of the following has highest reducing capacity
 (a) Ag (b) H_2
 (c) Zn (d) Li

31. If an iron rod is dipped in CuSO_4 solution [MADT Bihar 1984]
- Blue colour of the solution turns green
 - Brown layer is deposited on iron rod
 - No change occurs in the colour of the solution
 - Blue colour of the solution vanishes
 - None of the above
32. E° values of Mg^{2+}/Mg is -2.37V , of Zn^{2+}/Zn is -0.76V and Fe^{2+}/Fe is -0.44V . Which of the following statements is correct [EAMCET 1989]
- Zn will reduce Fe^{2+}
 - Zn will reduce Mg^{2+}
 - Mg oxidises Fe
 - Zn oxidises Fe
33. The standard reduction potential for Fe^{2+}/Fe and Sn^{2+}/Sn electrodes are -0.44 and -0.14 volt respectively. For the given cell reaction $\text{Fe}^{2+} + \text{Sn} \rightarrow \text{Fe} + \text{Sn}^{2+}$, the standard EMF is [IIT Screening 1990; MP PMT 2003]
- $+0.30\text{V}$
 - -0.58V
 - $+0.58\text{V}$
 - -0.30V
34. Electrode potential of Zn^{2+}/Zn is -0.76V and that of Cu^{2+}/Cu is $+0.34\text{V}$. The EMF of the cell constructed between these two electrodes is [EAMCET 1992; BHU 2001; CBSE PMT 2001; KCET 1990; MHCET 1999, 2003; Pb. CET 2002; AFMC 2001; Pb. PMT 2004]
- 1.10V
 - 0.42V
 - -1.1V
 - -0.42V
35. EMF of a cell whose half cells are given below is
 $\text{Mg}^{2+} + 2e^- \rightarrow \text{Mg}(s); E = -2.37\text{V}$
 $\text{Cu}^{2+} + 2e^- \rightarrow \text{Cu}(s); E = +0.33\text{V}$ [EAMCET 1987; MP PET 1994; Pb. PMT 2000]
- -2.03V
 - 1.36V
 - 2.7V
 - 2.03V
36. A cell constructed by coupling a standard copper electrode and a standard magnesium electrode has emf of 2.7 volts. If the standard reduction potential of copper electrode is $+0.34$ volt that of magnesium electrode is [KCET 1989]
- $+3.04$ volts
 - -3.04 volts
 - $+2.36$ volts
 - -2.36 volts
37. When $E^\circ_{\text{Ag}^+/\text{Ag}} = 0.8$ volt and $E^\circ_{\text{Zn}^{2+}/\text{Zn}} = -0.76$ volt, which of the following is correct [MP PMT 1994]
- Ag^+ can be reduced by H_2
 - Ag can oxidise H_2 into H^+
 - Zn^{2+} can be reduced by H_2
 - Ag can reduce Zn^{2+} ion
38. Adding powdered lead and iron to a solution that is 1.0M in both Pb^{2+} and Fe^{2+} ions, would result a reaction, in which [CPMT 1987]
- More iron and Pb^{2+} ions are formed
 - More lead and Fe^{2+} ions are formed
 - Concentration of both Pb^{2+} and Fe^{2+} ions increases
 - There is no net change
39. Given standard electrode potentials
- $$\text{Fe}^{++} + 2e^- \rightarrow \text{Fe}; E^\circ = -0.440\text{V}$$
- $$\text{Fe}^{+++} + 3e^- \rightarrow \text{Fe}; E^\circ = -0.036\text{V}$$
- The standard electrode potential (E°) for $\text{Fe}^{+++} + e^- \rightarrow \text{Fe}^{++}$ is [AIIMS 1982]
- -0.476V
 - -0.404V
 - $+0.404\text{V}$
 - $+0.771\text{V}$
40. Reduction potential of four elements P, Q, R, S is $-2.90, +0.34, +1.20$ and -0.76 . Reactivity decreases in the order [MP PET 1989; UPSEAT 2001]
- $P > Q > R > S$
 - $Q > P > R > S$
 - $R > Q > S > P$
 - $P > S > Q > R$
41. Which of the following metal can deposit copper from copper sulphate solution [CPMT 1983; MP PMT 1989]
- Mercury
 - Iron
 - Gold
 - Platinum
42. Standard electrode potential of Ag^+/Ag and Cu^+/Cu is $+0.80\text{V}$ and $+0.34\text{V}$ respectively. these electrodes are joint together by salt bridge if [AMU 2002]
- Copper electrode is work like cathode, then E°_{cell} is $+0.45\text{V}$
 - Silver electrode is work like anode then E°_{cell} is -0.34V
 - Copper electrode is work like anode then E°_{cell} is $+0.46\text{V}$
 - Silver electrode is work like cathode then E°_{cell} is -0.34V
 - Silver electrode is work like anode then E°_{cell} will be $+1.14\text{V}$
43. The reaction is spontaneous if the cell potential is [MP PET 1999]
- Positive
 - Negative
 - Zero
 - Infinite
44. Which substance eliminates bromine from KBr solution [IIT 1981]
- I_2
 - Cl_2
 - HI
 - SO_2
45. A standard hydrogen electrode has zero electrode potential because [IIT 1997]
- Hydrogen is easiest to oxidise
 - The electrode potential is assumed to be zero
 - Hydrogen atom has only one electron
 - Hydrogen is the lightest element
46. In the electrochemical cell
 $\text{H}_2(g) | 1\text{atm} | \text{H}^+(1\text{M}) || \text{Cu}^{2+}(1\text{M}) | \text{Cu}(s)$
 Which one of the following statements is true [EAMCET 1997]
- H_2 is cathode; Cu is anode
 - Oxidation occurs at Cu electrode
 - Reduction occurs at H_2 electrode
 - H_2 is anode; Cu is cathode
47. Expression representing the cell potential (E_{cell})
- $E_{\text{cath}} + E_{\text{an}}$
 - $E_{\text{an}} - E_{\text{cath}}$
 - $E_{\text{cath}} - E_{\text{an}}$
 - $E_{\text{c}} - E_{\text{a}}$
48. Iron displaces copper from its salt solution, because [MP PMT 1996]
- Atomic number of iron is less than that of copper

- (b) The standard reduction potential of iron is less than that of copper
(c) The standard reduction potential of iron is more than that of copper
(d) The iron salt is more soluble in water than the copper salt
49. (i) Copper metal dissolves in 1M silver nitrate solution and crystals of silver metal get deposited.
(ii) Silver metal does not react with 1 M zinc nitrate solution
(iii) Zinc metal dissolves in 1M copper sulphate solution and copper metal gets deposited
- Hence the order of decreasing strength of the three metals as reducing agents will be
- (a) $Cu > Ag > Zn$ (b) $Ag > Cu > Zn$
(c) $Zn > Cu > Ag$ (d) $Cu > Zn > Ag$
50. Standard electrode potentials of Zn and Fe are known to be (i) $-0.76 V$ and (ii) $-0.44 V$ respectively. How does it explain that galvanization prevents rusting of iron while zinc slowly dissolves away
- (a) Since (i) is less than (ii), zinc becomes the cathode and iron the anode
(b) Since (i) is less than (ii), zinc becomes the anode and iron the cathode
(c) Since (i) is more than (ii), zinc becomes the anode and iron the cathode
(d) Since (i) is more than (ii), zinc becomes the cathode and iron the anode
51. Amongst the following electrodes the one with zero electrode potential is [MP PMT 1997]
(a) Calomel electrode
(b) Standard hydrogen electrode
(c) Glass electrode
(d) Gas electrode
52. Which of the following is correct expression for electrode potential of a cell [MP PMT 1997]
- (a) $E = E^{\circ} - \frac{RT}{nF} \ln \frac{[\text{product}]}{[\text{reactant}]}$
(b) $E = E^{\circ} + \frac{RT}{F} \ln \frac{[\text{product}]}{[\text{reactant}]}$
(c) $E = E^{\circ} - \frac{RT}{nF} \ln \frac{[\text{reactant}]}{[\text{product}]}$
(d) $E = -\frac{RT}{F} \ln \frac{[\text{product}]}{[\text{reactant}]}$
53. Calculate standard free energy change for the reaction $\frac{1}{2} Cu(s) + \frac{1}{2} Cl_2(g) \rightleftharpoons \frac{1}{2} Cu^{2+} + Cl^{-}$ taking place at $25^{\circ} C$ in a cell whose standard e.m.f. is 1.02 volts [MP PMT 1997]
(a) $-98430 J$ (b) $98430 J$
(c) $96500 J$ (d) $-49215 J$
54. In which cell the free energy of a chemical reaction is directly converted into electricity? [MP PET/PMT 1998]
(a) Leclanche cell (b) Concentration cell
(c) Fuel cell (d) Lead storage battery
55. Nernst equation is related with
(a) The electrode potential and concentration of ions in the solution
(b) Equilibrium constant and concentration of ions
(c) Free energy change and E.M.F. of the cell
(d) None of these
56. The standard reduction potentials of 4 elements are given below. Which of the following will be the most suitable reducing agent
 $I = -3.04 V$, $II = -1.90 V$, $III = 0 V$, $IV = 1.90 V$ [CPMT 1999]
(a) I (b) II
(c) III (d) IV
57. Electrode potential data are given below :
 $Fe^{3+}(aq) + e^{-} \rightarrow Fe^{2+}(aq); E^{\circ} = +0.77 V$
 $Al^{3+}(aq) + 3e^{-} \rightarrow Al(s); E^{\circ} = -1.66 V$
 $Br_2(aq) + 2e^{-} \rightarrow 2Br^{-}(aq); E^{\circ} = +1.08 V$
- Based on the data given above, reducing power of Fe^{2+} , Al and Br^{-} will increase in the order [Pb. PMT 1998]
(a) $Br^{-} < Fe^{2+} < Al$ (b) $Fe^{2+} < Al < Br^{-}$
(c) $Al < Br^{-} < Fe^{2+}$ (d) $Al < Fe^{2+} < Br^{-}$
58. The standard electrode potential (E°) for OCl^{-}/Cl^{-} and $Cl^{-}/\frac{1}{2} Cl_2$ respectively are $0.94 V$ and $-1.36 V$. The E° value for $OCl^{-}/\frac{1}{2} Cl_2$ will be [KCET 1996]
(a) $-0.42 V$ (b) $-2.20 V$
(c) $0.52 V$ (d) $1.04 V$
59. If the reduction potential is more, then [CPMT 1996]
(a) It is easily oxidised
(b) It is easily reduced
(c) It acts as oxidising agent
(d) It has redox nature
60. One of the following is false for Hg [BHU 1998]
(a) It can evolve hydrogen from H_2S
(b) It is a metal
(c) It has high specific heat
(d) It is less reactive than hydrogen
61. E° for the cell $Zn|Zn^{2+}(aq)||Cu^{2+}(aq)|Cu$ is $1.10 V$ at $25^{\circ} C$, the equilibrium constant for the reaction $Zn + Cu^{2+}(aq) \rightleftharpoons Cu + Zn^{2+}(aq)$ is of the order of [CBSE PMT 1997]
(a) 10^{-28} (b) 10^{-37}
(c) 10^{+18} (d) 10^{+17}
62. Standard reduction potentials at $25^{\circ} C$ of $Li^{+}|Li$, $Ba^{2+}|Ba$, $Na^{+}|Na$ and $Mg^{2+}|Mg$ are -3.05 , -2.90 , -2.71 and -2.37 volt respectively. Which one of the following is the strongest oxidising agent [CBSE PMT 1994; JIPMER 2002]
(a) Na^{+} (b) Li^{+}
(c) Ba^{2+} (d) Mg^{2+}
63. Which of the following displaces Br_2 from an aqueous solution containing bromide ions [CBSE PMT 1994; JIPMER (Med.) 2002]
(a) Cl_2 (b) Cl^{-}
(c) I_2 (d) I_3^{-}
64. For the cell reaction
 $Cu^{2+}(C_1 aq) + Zn(s) = Zn^{2+}(C_2 aq) + Cu(s)$

of an electrochemical cell, the change in free energy at a given temperature is a function of [CBSE PMT 1998]


- (a) $\ln (C_1)$ (b) $\ln (C_2)$
(c) $\ln (C_1 + C_2)$ (d) $\ln (C_2 / C_1)$

65. The e.m.f. of the cell in which the following reaction $Zn(s) + Ni^{2+}(a=1.0) \rightleftharpoons Zn^{2+}(a=10) + Ni(s)$ occurs, is found to be 0.5105 V at 298 K. The standard e.m.f. of the cell is

[Roorkee Qualifying 1998]

- (a) 0.5400 (b) 0.4810 V
(c) 0.5696 V (d) -0.5105 V

66. For the redox reaction



E_{cell}° is 1.10 volt. E_{cell} for the cell will be $\left(2.303 \frac{RT}{F} = 0.0591\right)$

[AIEEE 2003]

- (a) 2.14 volt (b) 1.80 volt
(c) 1.07 volt (d) 0.82 volt


67. The emf of a Daniel cell at 298 K is E_1 $Zn | ZnSO_4 || CuSO_4 | Cu$ when the concentration of $ZnSO_4$ (0.01 M) (1.0 M)

is 1.0 M and that of $CuSO_4$ is 0.01 M the emf changed to E_2 . What is the relationship between E_1 and E_2

[CBSE PMT 2003]

- (a) $E_2 = 0 \neq E_1$ (b) $E_1 > E_2$
(c) $E_1 < E_2$ (d) $E_1 = E_2$

68. The oxidation potentials of following half-cell reactions are given



whose cell-reaction is $Fe^{2+}(aq) + Zn \rightarrow Zn^{2+}(aq) + Fe$

[MP PMT 2003]

- (a) -1.20 V (b) +0.32 V
(c) -0.32 V (d) +1.20 V

69. The E° for half cells Fe/Fe^{2+} and Cu/Cu^{2+} are -0.44 V and +0.32 V respectively. Then [MP PMT 2003]

- (a) Cu^{2+} oxidises Fe (b) Cu^{2+} oxidises Fe^{2+}
(c) Cu oxidises Fe^{2+} (d) Cu reduces Fe^{2+}

70. What is E° for electrode represented by $Pt, O_2(1 \text{ atm}) / 2H^+(1 \text{ M})$ [JIPMER 1997]

- (a) Unpredictable (b) Zero
(c) 0.018 V (d) 0.118 V



71. The cell potential of a cell in operation is

- (a) Zero (b) Positive
(c) Negative (d) None of the above

72. Which of the following is displaced by Fe [Roorkee 1995]

- (a) Ag (b) Hg
(c) Zn (d) Na

73. The standard electrode potential of the half cells are given below






The emf of the cell $Fe^{2+} + Zn \rightarrow Zn^{2+} + Fe$ is

[CPMT 2003]

- (a) 1.54 V (b) -1.54 V

- (c) -0.19 V (d) +0.19 V

74. $Zn^{2+} + 2e^- \rightarrow Zn(s); E^{\circ} = -0.76$,



Strongest reducing agent is

[BHU 2003]

- (a) Fe^{2+} (b) Zn
(c) Cr (d) H_2

75. Standard reduction electrode potentials of three metals A, B and C are respectively +0.5 V, -3.0 V and -1.2 V. The reducing powers of these metals are [IIT 1998; AIEEE 2003]


- (a) $B > C > A$ (b) $A > B > C$
(c) $C > B > A$ (d) $A > C > B$

76. For a cell reaction involving a two-electron change, the standard emf of the cell is found to be 0.295 V at 25°C. The equilibrium constant of the reaction at 25°C will be

[Roorkee 1999; AIEEE 2003; CBSE PMT 2004]

- (a) 1×10^{-10} (b) 29.5×10^{-2}
(c) 10 (d) 1×10^{10}

77. For the electrochemical cell,



[IIT-JEE (Screening) 2000]

- (a) $M + X \rightarrow M^+ + X^-$ is the spontaneous reaction
(b) $M^+ + X^- \rightarrow M + X$ is the spontaneous reaction
(c) $E_{cell} = 0.77 \text{ V}$
(d) $E_{cell} = -0.77 \text{ V}$

78. The standard potential at 25°C for the following half reactions are given against them



When zinc dust is added to the solution of $MgCl_2$


[UPSEAT 2001]

- (a) $ZnCl_2$ is formed
(b) Zinc dissolves in the solution
(c) No reaction takes place
(d) Mg is precipitated

79. $KMnO_4$ acts as an oxidising agent in the neutral medium and gets reduced to MnO_2 . The equivalent weight of $KMnO_4$ in neutral medium [AMU 2001]


- (a) mol. wt/2 (b) mol. wt/3
(c) mol. wt/4 (d) mol. wt/7

80. Which of the following condition will increase the voltage of the cell, represented by the equation



- (a) Increase in the concentration of Ag^+ ion
(b) Increase in the concentration of Cu^{2+} ion
(c) Increase in the dimension of silver electrode
(d) Increase in the dimension of copper electrode

81. Which will increase the voltage of the cell



- (a) Increase in the concentration of Ag^+ ions
(b) Increase in the concentration of Sn^{2+} ions
(c) Increase in size of the silver rod
(d) None of these
82. The mass of the proton is 1840 times that of electron, its potential difference is V . The kinetic energy of proton is [DCE 2001]
(a) 1840 KeV (b) 1 KeV
(c) $\frac{1}{1840}$ KeV (d) 920 KeV
83. What will be the emf for the given cell
 $Pt | H_2(P_1) | H^+(aq) || H_2(P_2) | Pt$ [AIEEE 2002]
(a) $\frac{RT}{f} \log \frac{P_1}{P_2}$ (b) $\frac{RT}{2f} \log \frac{P_1}{P_2}$
(c) $\frac{RT}{f} \log \frac{P_2}{P_1}$ (d) None of these
84. What is the potential of a cell containing two hydrogen electrodes the negative one in contact with $10^{-8} M H^+$ and positive one in contact with $0.025 M H^+$ [MP PMT 2000]
(a) 0.18 V (b) 0.28 V
(c) 0.38 V (d) 0.48 V
85. Will $Fe(s)$ be oxidised to Fe^{2+} by the reaction with $1 M HCl$ (E° for $Fe/Fe^{2+} = +0.44 V$) [Pb. PMT 2000]
(a) Yes (b) No
(c) May be (d) Can't say
86. EMF of a cell in terms of reduction potential of its left and right electrodes is [AIEEE 2002]
(a) $E = E_{left} - E_{right}$ (b) $E = E_{left} + E_{right}$
(c) $E = E_{right} - E_{left}$ (d) $E = -(E_{right} + E_{left})$
87. Arrange the following in the order of their decreasing electrode potential Mg, K, Ba, Ca [JIPMER 2002]
(a) K, Ba, Ca, Mg (b) Ca, Mg, K, Ba
(c) Ba, Ca, K, Mg (d) Mg, Ca, Ba, K
88. Which of the following has highest electrode potential [Pb. PMT 2000]
(a) Li (b) Cu
(c) Au (d) Al
89. The cell reaction of a cell is
 $Mg(s) + Cu^{2+}(aq) \rightarrow Cu(s) + Mg^{2+}(aq)$
If the standard reduction potentials of Mg and Cu are -2.37 and $+0.34 V$ respectively. The emf of the cell is [EAMCET 1995; JIPMER (Med.) 2001; AFMC 2002; CBSE PMT 2002]
(a) 2.03 V (b) $-2.03 V$
(c) $+2.71 V$ (d) $-2.71 V$
90. The element which can displace three other halogens from their compound is [EAMCET 1998]
(a) Cl (b) F
(c) Br (d) I
91. Which of the following has been universally accepted as a reference electrode at all temperatures and has been assigned a value of zero volt [AIIMS 1998]
(a) Graphite electrode
(b) Copper electrode
(c) Platinum electrode
(d) Standard hydrogen electrode
92. Aluminium is more reactive than Fe . But Al is less easily corroded than iron because [DCE 1999]
(a) Al is noble metal
(b) Iron forms both mono and divalent ions
(c) Oxygen forms a protective oxide layer
(d) Fe undergoes reaction easily with H_2O
93. Zinc displaces copper from the solution of its salt because [MP PET 1995]
(a) Atomic number of zinc is more than that of copper
(b) Zinc salt is more soluble in water than the copper salt
(c) Gibbs free energy of zinc is less than that of copper
(d) Zinc is placed higher than copper in electro-chemical series
94. An electrochemical cell is set up as follows
 $Pt(H_2, 1 atm) / 0.1 M HCl$
 $|| 0.1 M \text{ acetic acid } / (H_2, 1 atm) Pt$
E.M.F. of this cell will not be zero because [CBSE PMT 1995]
(a) The pH of $0.1 M HCl$ and $0.1 M$ acetic acid is not the same
(b) Acids used in two compartments are different
(c) E.M.F. of a cell depends on the molarities of acids used
(d) The temperature is constant
95. Cu^+ ion is not stable in aqueous solution because of disproportionation reaction. E° value for disproportionation of Cu^+ is
(Given $E^\circ_{Cu^{2+}/Cu^+} = 0.15$, $E^\circ_{Cu^{2+}/Cu} = 0.34 V$) [IIT 1995]
(a) $-0.49 V$ (b) $0.49 V$
(c) $-0.38 V$ (d) $0.38 V$
96. E° of a cell $aA + bB \rightarrow cC + dD$ is [CPMT 1997]
(a) $-\frac{RT}{nF} \log \frac{[C]^c [D]^d}{[A]^a [B]^b}$ (b) $-RT \log \frac{[a]^A [b]^B}{[a]^C [d]^D}$
(c) $-\frac{RT}{nF} \log \frac{[C]^c [d]^D}{[A]^a [B]^b}$ (d) $-\frac{RT}{nF} \log \frac{[C]^c [d]^D}{[a]^A [b]^B}$
97. In the experiment set up for the measurement of EMF of a half cell using a reference electrode and a salt bridge, when the salt bridge is removed, the voltage [NCERT 1984; CPMT 1985]
(a) Does not change
(b) Decreases to half the value
(c) Increase to maximum
(d) Drops to zero
98. Electrode potentials of five elements A, B, C, D and E are respectively -1.36 , -0.32 , 0 , -1.26 and -0.42 . The reactivity order of these elements are in the order of [MP PMT 1995]
(a) A, D, E, B and C (b) C, B, E, D and A
(c) B, D, E, A and C (d) C, A, E, D and B
99. What is wrongly stated about electrochemical series

[DCE 1999]

- (a) It is the representation of element in order of increasing or decreasing standard electrode reduction potential
 (b) It does not compare the relative reactivity of metals
 (c) It compares relative strengths of oxidising agents
 (d) H_2 is centrally placed element

100. Which of the following statements is true for fuel cells

[KCET (Med.) 1999; AFMC 2000]

- (a) They are more efficient
 (b) They are free from pollution
 (c) They run till reactants are active
 (d) All of these

101. What is the potential of a half-cell consisting of zinc electrode in 0.01M $ZnSO_4$ solution at $25^\circ C$ ($E^\circ = 0.763 V$)

[AIIMS 2000; BHU 2000]

- (a) 0.8221 V (b) 8.221 V
 (c) 0.5282 V (d) 9.232 V

102. The *emf* of a galvanic cell, with electrode potentials of silver = +0.80 V and that of copper = +0.34 V, is

[AIIMS 1999]

- (a) -1.1 V (b) +1.1 V
 (c) +0.46 V (d) +0.76 V

103. Copper cannot replace..... from solution [DPMT 2002]

- (a) Fe (b) Au
 (c) Hg (d) Ag

104. The strongest reducing agent of the alkali metal is

[CBSE PMT 2000]

- (a) Li (b) Na
 (c) K (d) Cs

105. Which of the following is the most electropositive element

[Pb. PMT 2000]

- (a) Carbon (b) Calcium
 (c) Chlorine (d) Potassium

106. The metal that forms a self protecting film of oxide to prevent corrosion, is

[BHU 1999]

- (a) Cu (b) Al
 (c) Na (d) Au

107. In a cell that utilises the reaction $Zn_{(s)} + 2H^+(aq) \rightarrow$ $Zn^{2+}(aq) + H_{2(g)}$ addition of H_2SO_4 to cathode compartment,

will

[AIEEE 2004]

- (a) Increase the E and shift equilibrium to the right
 (b) Lower the E and shift equilibrium to the right
 (c) Lower the E and shift equilibrium to the left
 (d) Increase the E and shift equilibrium to the left

108. For the electrochemical cell, $M | M^+ || X^- | X$, $E^\circ(M^+ | M) = 0.44 V$ $E^\circ(X | X^-) = 0.33 V$ From this data, one can deduce that

[Pb.CET 2004]

- (a) $E^\circ_{cell} = -0.77 V$
 (b) $M^+ + X^- \rightarrow M + X$ is the spontaneous reaction
 (c) $M + X \rightarrow M^+ + X^-$ is the spontaneous reaction
 (d) $E^\circ_{cell} = .77 V$

109. The standard *emf* of a cell, involving one electron change is found to be 0.591 V at $25^\circ C$. The equilibrium constant of the reaction is $(F = 96,500 C mol^{-1}; R = 8.314 JK^{-1}mol^{-1})$ [AIEEE 2004]

- (a) 1.0×10^{10} (b) 1.0×10^5
 (c) 1.0×10^1 (d) 1.0×10^{30}

110. Standard electrode potential of cell $H_2 | H^+ || Ag^+ | Ag$ is

[AIEEE 2004]

- (a) 0.8 V (b) -0.8 V
 (c) -1.2 V (d) 1.2 V

111. A galvanic cell with electrode potential of 'A' = +2.23 V and 'B' = -1.43 V. The value of E°_{cell} is [Pb.CET 2003]

- (a) 3.66 V (b) 0.80 V
 (c) -0.80 V (d) -3.66 V

112. The *emf* of a cell whose half cells are given below is $Mg^{2+} + 2e^- \rightarrow Mg(s)$ $E^\circ = -2.37 V$ $Cu^{2+} + 2e^- \rightarrow Cu(s)$ $E^\circ = +0.34 V$

[Pb.CET 2001]

- (a) +1.36 V (b) +2.71 V
 (c) +2.17 V (d) -3.01 V

113. For the cell reaction, $2Ce^{4+} + Co \rightarrow 2Ce^{3+} + Co^{2+}$ E°_{cell} is 1.89 V. If $E^\circ_{Ce^{4+}/Ce^{3+}}$

[Pb.CET 2000]

- (a) -1.64 V (b) +1.64 V
 (c) -2.08 V (d) +2.17 V

114. If the ΔG of a cell reaction $AgCl + e^- \rightarrow Ag + Cl^-$ is -21.20 KJ; the standard *emf*, of cell is

[MP PMT 2004]

- (a) 0.229 V (b) 0.220 V
 (c) -0.220 V (d) -0.110 V

115. The *emf* of the cell $Ag | Ag^+(0.1M) || Ag^+(1M) | Ag$ at 298 K is

[DCE 2003]

- (a) 0.0059 V (b) 0.059 V
 (c) 5.9 V (d) 0.59 V

116. The *emf* of the cell $Zn | Zn^{2+}(0.01M) || Fe^{2+}(0.001M) | Fe$ at 298 K is 0.2905 then the value of equilibrium for the cell reaction is [IIT-JEE Scr

- (a) $\frac{0.32}{e^{0.0295}}$ (b) $\frac{0.32}{10^{0.0295}}$
 (c) $\frac{0.26}{10^{0.0295}}$ (d) $\frac{0.32}{10^{0.0591}}$

117. Aluminium displaces hydrogen from dilute HCl whereas silver does not. The *emf* of a cell prepared by combining Al/Al^{3+} and Ag/Ag^+ is 2.46 V. The reduction potential of silver electrode is +0.80 V. The reduction potential of aluminium electrode is

[KCET 2004]

- (a) +1.66 V (b) -3.26 V
 (c) 3.26 V (d) -1.66 V

118. Consider the following E° values : $E^\circ_{Fe^{3+}/Fe^{2+}} = +0.77 V$ $E^\circ_{Sn^{2+}/Sn} = -0.14 V$

- Under standard conditions the potential for the reaction $\text{Sn}_{(s)} + 2\text{Fe}^{3+}(\text{aq}) \rightarrow 2\text{Fe}^{2+}(\text{aq}) + \text{Sn}^{2+}(\text{aq})$ is [AIEEE 2004]
- (a) 0.91 V (b) 1.40 V
(c) 1.68 V (d) 0.63 V
119. $\text{Cr}_2\text{O}_7^{2-} + \text{I}^- \rightarrow \text{I}_2 + \text{Cr}^{3+}$
 $E^0_{\text{cell}} = 0.79 \text{ V}$
 $E^0_{\text{Cr}_2\text{O}_7^{2-}} = 1.33 \text{ V}$, $E^0_{\text{I}_2}$ is [BVP 2004]
- (a) -0.10 V (b) +0.18 V
(c) -0.54 V (d) 0.54 V
120. $\text{Zn}_{(s)} + \text{Cl}_2(1 \text{ atm}) \rightarrow \text{Zn}^{2+} + 2\text{Cl}^-$. E^0_{cell} of the cell is 2.12 V. To increase E [BVP 2004]
- (a) $[\text{Zn}^{2+}]$ should be increased
(b) $[\text{Zn}^{2+}]$ should be decreased
(c) $[\text{Cl}^-]$ should be decreased
(d) P_{Cl_2} should be decreased
121. The $E^0_{M^{3+}/M^{2+}}$ values for Cr, Mn, Fe and Co are -0.41, +1.57, +0.77 and +1.97 V respectively. For which one of these metals the change in oxidation state from +2 to +3 is easiest [AIEEE 2004]
- (a) Fe (b) Mn
(c) Cr (d) Co
122. The rusting of iron takes place as follows
 $2\text{H} + 2\text{e}^- + \frac{1}{2}\text{O}_2 \longrightarrow \text{H}_2\text{O}(\text{l})$;
 $E^0 = +1.23 \text{ V}$
 $\text{Fe} + 2\text{e}^- \longrightarrow \text{Fe}(\text{s})$; $E^0 = -0.44 \text{ V}$
Calculate ΔG^0 for the net process [IIT 2005]
- (a) -322 kJ mol⁻¹ (b) -161 kJ mol⁻¹
(c) -152 kJ mol⁻¹ (d) -76 kJ mol⁻¹
123. When an acid cell is charged, then [AFMC 2005]
- (a) Voltage of cell increases
(b) Electrolyte of cell dilutes
(c) Resistance of cell increases
(d) None of these
124. The standard electrode potential is measured by [KCET 2005]
- (a) Electrometer (b) Voltmeter
(c) Pyrometer (d) Galvanometer
125. Aluminium displaces hydrogen from acids but copper does not. A galvanic cell prepared by combining Cu/Cu^{2+} and Al/Al^{3+} has an e.m.f. of 2.0 V at 298 K. If the potential of copper electrode is +0.34 V, that of aluminium is [CPMT 2001; KCET 2001]
- (a) +1.66 V (b) -1.66 V
(c) +2.34 V (d) -2.3 V
126. If the standard electrode potential of Cu^{2+}/Cu electrode is 0.34 V, what is the electrode potential of 0.01M concentration of Cu^{2+} ($T = 298 \text{ K}$) [EAMCET 2003]
- (a) 0.399 V (b) 0.281 V
(c) 0.222 V (d) 0.176 V
127. Calculate the electrode potential at 298°K for $\text{Zn}|\text{Zn}^{++}$ electrode in which the activity of zinc ions is 0.001 M and $E^0_{\text{Zn}/\text{Zn}^{++}}$ is -0.74 volts [AMU 2002]
- (a) 0.38 volts (b) 0.83 volts
(c) 0.40 volts (d) 0.45 volts
128. Which of the following expression is correct [Orissa JEE 2005]
- (a) $\Delta G^0 = -nFE^0_{\text{cell}}$ (b) $\Delta G^0 = +nFE^0_{\text{cell}}$
(c) $\Delta G^0 = -2.303RTnFE^0_{\text{cell}}$ (d) $\Delta G^0 = -nF \log K_c$
129. For the feasibility of a redox reaction in a cell, the e.m.f. should be [J & K 2002]
- (a) Positive (b) Fixed
(c) Zero (d) Negative


Corrosion

1. Corrosion is basically a [Kerala (Med.) 2002]
- (a) Altered reaction in presence of H_2O
(b) Electrochemical phenomenon
(c) Interaction
(d) Union between light metal and heavy metal
2. Rusting of iron is catalysed by which of the following [MNR 1990; UPSEAT 2001]
- (a) Fe (b) O_2
(c) Zn (d) H^+
3. Which of the following is a highly corrosive salt [AFMC 2005]
- (a) FeCl_2 (b) PbCl_2
(c) Hg_2Cl_2 (d) HgCl_2
4. Corrosion of iron is essentially an electrochemical phenomenon where the cell reactions are [KCET 2005]
- (a) Fe is oxidised to Fe^{2+} and dissolved oxygen in water is reduced to OH^-
(b) Fe is oxidised to Fe^{3+} and H_2O is reduced to O_2^{2-}
(c) Fe is oxidised to Fe^{2+} and H_2O is reduced to O_2^-
(d) Fe is oxidised to Fe^{2+} and H_2O is reduced to O_2

Critical Thinking

Objective Questions

1. The limiting molar conductivities \wedge^0 for NaCl, KBr and KCl are 126, 152 and 150 $\text{S cm}^2\text{mol}^{-1}$ respectively. The \wedge^0 for NaBr is
- (a) 278 $\text{S cm}^2\text{mol}^{-1}$ (b) 176 $\text{S cm}^2\text{mol}^{-1}$
(c) 128 $\text{S cm}^2\text{mol}^{-1}$ (d) 302 $\text{S cm}^2\text{mol}^{-1}$
2. On the basis of the electrochemical theory of aqueous corrosion, the reaction occurring at the cathode is [MP PET 1994; UPSEAT 2001]
- (a) $\text{O}_{2(\text{g})} + 4\text{H}^+_{(\text{aq})} + 4\text{e}^- \rightarrow 2\text{H}_2\text{O}_{(\text{l})}$
(b) $\text{Fe}_{(\text{s})} \rightarrow \text{Fe}^{2+}_{(\text{aq})} + 2\text{e}^-$
(c) $\text{Fe}^{2+}_{(\text{aq})} \rightarrow \text{Fe}^{3+}_{(\text{aq})} + \text{e}^-$





3. The reaction $\frac{1}{2}H_{2(g)} + AgCl(s) \rightarrow H^+(aq) + Cl^-(aq) + Ag(s)$ occurs in the galvanic cell

[IIT 1985; AMU 2002; KCET 2003]

- (a) $Ag / AgCl(s)KCl(soln) || AgNO_3(soln) / Ag$
 (b) $Pt / H_2(g)HCl(soln) || AgNO_3(soln) / Ag$
 (c) $Pt / H_2(g)HCl(soln) || AgCl(s) / Ag$
 (d) $Pt / H_2(g)KCl(soln) || AgCl(s) / Ag$

4. The standard reduction potential E° for the half reactions are as



The EMF for cell reaction $Fe^{2+} + Zn \rightarrow Zn^{2+} + Fe$ is

[IIT 1988; CBSE PMT 1993, 96; BHU 1995, 2000; CPMT 2000; KCET 2000; AIIMS 2001; Orissa JEE 2002]

- (a) $-0.35 V$ (b) $+0.35 V$
 (c) $+1.17 V$ (d) $-1.17 V$

5. The number of electrons to balance the following equation $NO_3^- + 4H^+ + e^- \rightarrow 2H_2O + NO$ is [IIT Screening 1991]

- (a) 5 (b) 4
 (c) 3 (d) 2

6. The standard EMF for the given cell reaction $Zn + Cu^{2+} = Cu + Zn^{2+}$ is $1.10 V$ at $25^\circ C$. The EMF for the cell reaction, when $0.1 M Cu^{2+}$ and $0.1 M Zn^{2+}$ solutions are used, at $25^\circ C$ is

[MNR 1994; AMU 1999; UPSEAT 2002]

- (a) $1.10 V$ (b) $0.110 V$
 (c) $-1.10 V$ (d) $-0.110 V$

7. A gas X at $1 atm$ is bubbled through a solution containing a mixture of $1 M Y^-$ and $1 M Z^-$ at $25^\circ C$. If the reduction potential of $Z > Y > X$, then [IIT 1999]

- (a) Y will oxidize X and not Z
 (b) Y will oxidize Z and not X
 (c) Y will oxidize both X and Z
 (d) Y will reduce both X and Z


8. The oxidation potential of a hydrogen electrode at $pH = 10$ and $pH_1 = 1$ [JIPMER 2000]

- (a) $0.059 V$ (b) $0.59 V$
 (c) $0.00 V$ (d) $0.51 V$

9. The decomposition of hydrogen peroxide is an example of [Roorkee 2000]

- (a) Exothermic reaction (b) Endothermic reaction
 (c) Negative catalysis (d) Auto-oxidation

10. Aluminium oxide may be electrolysed at $1000^\circ C$ to furnish aluminium metal (At. Mass = $27 amu$; $1 Faraday = 96,500 Coulombs$). The cathode reaction is



To prepare $5.12 kg$ of aluminium metal by this method would require [AIEEE 2005]

- (a) $5.49 \times 10^7 C$ of electricity
 (b) $1.83 \times 10^7 C$ of electricity
 (c) $5.49 \times 10^4 C$ of electricity
 (d) $5.49 \times 10^1 C$ of electricity

11.

Electrolyte :	KCl	KNO_3	HCl	$NaOAc$	$NaCl$
$\Lambda^\infty (Scm^2 mol^{-1})$:	149.9	145.0	426.2	91.0	126.5

Calculate Λ_{HOAc}^∞ using appropriate molar conductances of the electrolytes listed above at infinite dilution in H_2O at $25^\circ C$ [AIEEE 2005]

- (a) 517.2 (b) 552.7
 (c) 390.7 (d) 217.5

12. The mass of carbon anode consumed (giving only carbondioxide) in the production of $270 kg$ of aluminium metal from bauxite by the Hall process is [CBSE PMT 2005]

- (a) $180 kg$ (b) $270 kg$
 (c) $540 kg$ (d) $90 kg$

13. $4.5 g$ of aluminium (at mass $27 amu$) is deposited at cathode from Al^{3+} solution by a certain quantity of electric charge. The volume of hydrogen produced at STP from H^+ ions in solution by the same quantity of electric charge will be [CBSE PMT 2005]

- (a) $22.4 L$ (b) $44.8 L$
 (c) $5.6 L$ (d) $11.2 L$

14. What amount of Cl_2 gas liberated at anode, if $1 amp.$ current is passed for $30 min.$ from $NaCl$ solution. [BHU 2005]

- (a) $0.66 moles$ (b) $0.33 moles$
 (c) $0.66 g$ (d) $0.33 g$

Assertion & Reason

For AIIMS Aspirants

Read the assertion and reason carefully to mark the correct option out of the options given below :

- (a) If both assertion and reason are true and the reason is the correct explanation of the assertion.
 (b) If both assertion and reason are true but reason is not the correct explanation of the assertion.
 (c) If assertion is true but reason is false.
 (d) If the assertion and reason both are false.
 (e) If assertion is false but reason is true.

1. Assertion : Sodium ions are discharged in preference to hydrogen ions at a mercury cathode.
 Reason : The nature of the cathode can effect the order of discharge of ions.
 2. Assertion : In electrolysis, the quantity of electricity needed for depositing 1 mole of silver is different from that required for 1 mole of copper.
 Reason : The molecular weights of silver and copper are different
 3. Assertion : Equivalent conductance of all electrolytes decreases with the increases in concentration.
 Reason : Lesser number of ions are available per gram equivalent at higher concentration.
 4. Assertion : Copper reacts with hydrochloric acid and liberates hydrogen from the solution of dilute hydrochloric acid.
 Reason : Hydrogen is below copper in the electrochemical series.

5. Assertion : Copper liberates hydrogen from a solution of dil. HCl .
Reason : Hydrogen is below copper in the reactivity series.
6. Assertion : Zn metal is formed when a Cu plate is dipped in $ZnSO_4$ solution.
Reason : Cu being placed above Zn in electrochemical series.
7. Assertion : Electrical conductivity of copper increases with increase in temperature.
Reason : The electrical conductivity of metals is due to the motion of electrons.
8. Assertion : A small amount of acid or alkali is added before electrolysis of water.
Reason : Pure water is weak electrolyte.
9. Assertion : Copper reacts with HCl and liberates hydrogen.
Reason : Hydrogen is present above Cu in the reactivity series.
10. Assertion : K and Cs are used in photoelectric cells.
Reason : K and Cs emit electrons on exposure to light.
11. Assertion : A large dry cell has high *e.m.f.*
Reason : The *e.m.f.* of a dry cell is proportional to its size.
12. Assertion : The resistivity for a substance is its resistance when its is one meter long and its area of cross section is one square meter.
Reason : The *S.I.* units of resistivity are ohm metre (Ωm) and ohm centimeter (Ωcm).
13. Assertion : When acidified zinc sulphate solution is electrolysed between zinc electrodes, it is zinc that is deposited at the cathode and hydrogen evolution does not take place.
Reason : The electrode potential of zinc is more negative than hydrogen as the overvoltage for the hydrogen as the evolution on zinc is quite large.
14. Assertion : If $\lambda_{Na^+}^0 + \lambda_{Cl^-}^0$ are molar limiting conductivity of the sodium and chloride ions respectively, then the limiting molar conducting for sodium chloride is given by the equation:
$$\wedge_{NaCl}^0 = \lambda_{Na^+}^0 + \lambda_{Cl^-}^0$$

Reason : This is according to Kohlrausch law of independent migration of ions.
15. Assertion : One coulomb of electric charge deposits weight equal to the electrochemical equivalent of the substance.
Reason : One Faraday deposits one mole of the substance.
16. Assertion : Auric chloride ($AuCl_3$) solution cannot be stored in a vessel made of copper, iron, nickel, chromium, zinc or tin.
Reason : Gold is a very precious metal.
17. Assertion : For a cell reaction $Zn(s) + Cu^{2+}(aq) \rightarrow Zn^{2+}(aq) + Cu(s)$; at the equilibrium, voltmeter gives zero reading.
Reason : At the equilibrium, there is no change in the concentration of Cu^{2+} and Zn^{2+} ions.
18. Assertion : A negative value of standard reduction potential means that reduction take place on this electrode with reference to standard hydrogen electrode.
Reason : The standard electrode potential of a half cell has a fixed value.
19. Assertion : Weston is a standard cell.
Reason : Its *e.m.f.* does not change with temperature.
20. Assertion : Galvanic cells containing hydrogen, methane, methanol etc. as fuels are called fuel cells.
Reason : They are designed to convert the energy of combustion of fuels directly into electrical energy.
21. Assertion : Zinc displaces copper from copper sulphate solution.
Reason : The E^0 of zinc is $-0.76 V$ and that of copper is $+0.34 V$.
22. Assertion : Identification of cathode and anode is done by the use of a thermometer.
Reason : Higher is the value of reduction potential, greater would be its reducing power.
23. Assertion : An electrochemical cell can be set up only if the redox reaction is spontaneous.
Reason : A reaction is spontaneous if free energy change is negative.
24. Assertion : Galvanised iron does not rust.
Reason : Zinc has a more negative electrode potential than iron. [AIIMS 2005]
25. Assertion : In an electrochemical cell anode and cathode are respectively negative and positive electrodes.
Reason : At anode oxidation takes place and at cathode reduction takes place.
26. Assertion : $Ni / Ni^{2+} (1.0M) || Au^{3+} (1.0M) | Au$, for this cell emf is $1.75 V$ if $E_{Au^{3+}/Au}^0 = 1.50$ and $E_{Ni^{2+}/Ni}^0 = 0.25 V$.
Reason : Emf of the cell $= E_{cathode}^0 - E_{anode}^0$.
27. Assertion : Salts like KCl, KNO_3 i.e., inert electrolytes are used in salt bridge.
Reason : An inert electrolyte can easily be filled in the U-tube.
28. Assertion : Emf and potential difference are same for cell.
Reason : Both gives the difference in electrode potential under any condition.

Answers

Electrolytes and Electrolysis

1	b	2	d	3	c	4	c	5	d
6	d	7	b	8	b	9	b	10	a
11	a	12	c	13	d	14	d	15	c
16	d	17	a	18	c	19	c	20	a
21	a	22	b	23	b	24	d	25	b
26	b	27	b	28	a	29	c	30	b
31	d	32	d	33	c	34	b	35	b
36	a	37	d	38	c	39	d	40	a
41	a								

Faraday's law of electrolysis

1	c	2	a	3	b	4	c	5	c
6	c	7	c	8	b	9	c	10	a
11	c	12	a	13	c	14	b	15	b

16	c	17	a	18	a	19	d	20	c
21	d	22	c	23	b	24	d	25	a
26	d	27	a	28	d	29	c	30	a
31	b	32	b	33	a	34	b	35	b
36	b	37	b	38	b	39	a	40	a
41	c	42	a	43	a	44	b	45	c
46	b	47	d	48	d	49	b	50	a
51	c	52	b	53	b	54	a	55	c
56	c	57	c	58	d	59	d	60	d
61	c	62	c	63	b	64	a	65	b
66	b	67	a	68	a	69	b	70	c
71	c	72	b	73	a	74	b	75	c
76	b	77	c	78	c	79	b	80	b
81	c	82	b	83	a	84	c	85	c
86	e	87	b	88	a	89	d	90	c
91	c	92	a	93	c	94	b		

Conductor and Conductance

1	b	2	b	3	d	4	d	5	a
6	b	7	b	8	a	9	b	10	d
11	a	12	b	13	b	14	b	15	b
16	d	17	b	18	d	19	b	20	c
21	b	22	a	23	c	24	d	25	a
26	b	27	d						

Cell constant and Electrochemical cells

1	d	2	d	3	d	4	b	5	b
6	a	7	c	8	c	9	b	10	b
11	c	12	c	13	a	14	a	15	d
16	c	17	c	18	a	19	b	20	b
21	b	22	b	23	a	24	b	25	c
26	a	27	c	28	a	29	b	30	c
31	a	32	b	33	c	34	b	35	d
36	b	37	c	38	d	39	b	40	d
41	a	42	d	43	c	44	a	45	d
46	a	47	d	48	b	49	c	50	c
51	b	52	d	53	c	54	a	55	c
56	d	57	b	58	a	59	a	60	c
61	a								

Electrode potential, E_{Cell} , Nernst equation and ECS

1	b	2	a	3	a	4	b	5	d
6	c	7	c	8	a	9	a	10	b
11	a	12	c	13	a	14	d	15	c

16	a	17	a	18	c	19	b	20	b
21	a	22	b	23	a	24	c	25	a
26	a	27	c	28	a	29	d	30	d
31	b	32	a	33	d	34	a	35	c
36	d	37	a	38	b	39	d	40	d
41	b	42	c	43	a	44	b	45	b
46	d	47	c	48	b	49	c	50	b
51	b	52	a	53	a	54	c	55	a
56	a	57	a	58	a	59	c	60	a
61	b	62	d	63	a	64	d	65	b
66	c	67	b	68	b	69	a	70	b
71	b	72	ab	73	c	74	c	75	a
76	d	77	b	78	c	79	b	80	a
81	a	82	b	83	b	84	c	85	a
86	c	87	a	88	c	89	c	90	b
91	d	92	c	93	d	94	a	95	d
96	a	97	d	98	a	99	b	100	d
101	a	102	c	103	a	104	a	105	d
106	b	107	a	108	b	109	a	110	a
111	a	112	b	113	b	114	b	115	b
116	b	117	d	118	a	119	d	120	b
121	c	122	a	123	a	124	b	125	b
126	b	127	b	128	a	129	a		

Corrosion

1	b	2	d	3	d	4	a		
---	---	---	---	---	---	---	---	--	--

Critical Thinking Questions

1	c	2	a	3	c	4	b	5	c
6	a	7	a	8	b	9	acd	10	a
11	c	12	d	13	c	14	c		

Assertion & Reason

1	a	2	b	3	a	4	d	5	d
6	d	7	e	8	a	9	e	10	a
11	d	12	b	13	a	14	a	15	c
16	b	17	a	18	e	19	a	20	a
21	a	22	d	23	b	24	a	25	a
26	a	27	c	28	d				