			Gaseous State 235
			(a) 1.5 <i>lit.</i> (b) 2.8 <i>lit.</i>
	Ordinary Thinking		(c) 11.2 <i>lit.</i> (d) 22.4 <i>lit.</i>
_		12.	Pressure of a gas in a vessel can be measured by
_			(a) Barometer (b) Manometer
_	Objective Questions		(c) Stalgometer (d) All the baove
Ch	aracteristics and Measurable properties of gases	13.	Volume occupied by a gas at one atmospheric pressure and $0^{o}C$ is V mL. Its volume at 273 K will be
,	Which are of the following statements is not correct shout the three		[Bihar MADT 1982]
1.	states of matter <i>i.e.</i> solid. liquid and gaseous		(a) $V ml$ (b) $V/2 ml$
	 (a) Molecules of a solid possess least energy whereas those of a gas possess highest energy 	14.	(c) 2 V (d) None of these Which one of the following statements is wrong for gases
	(b) The density of solid is highest whereas that of gases is lowest		[CBSE PMT 1999]
	(c) Gases like liquids possess definite volumes		(a) Gases do not have a definite shape and volume
	(d) Molecules of a solid possess vibratory motion		(b) Volume of the gas is equal to the volume of the container
2.	The temperature and pressure at which ice, liquid water and water vapour can exist together are		 (c) Confined gas exerts uniform pressure on the walls of its container in all directions
	(a) $0^{o}C, 1 atm$ (b) $2^{o}C, 4.7 atm$		(d) Mass of the gas cannot be determined by weighing a container in which it is enclosed
	(c) $0^{\circ}C 4.7 mm$ (d) $-2^{\circ}C 4.7 mm$	15.	Which of the following exhibits the weakest intermolecular forces [AIIMS 2000]
_			(a) NH_2 (b) HCl
3.	() The lowing is true about gaseous state		$(b) H_{2} \qquad (b) H_{2} \qquad (c)$
	(a) Thermal energy = Molecular attraction (b) Thermal energy - Molecular attraction		(c) He (d) H_2O
	(b) Thermal energy >> Molecular attraction	16.	N_2 is found in a litre flask under $100 kPa$ pressure and O_2 is
	(d) Molegular foreces >> These in liquids		found in another 3 litre flask under $320 kPa$ pressure. If the two
	(d) Molecular forces >> Those in liquids		flasks are connected, the resultant pressures is
4.	(a) Cases (b) Solida		[Kerala PMT 2004]
	(a) Gases (b) Solutions (d) Solutions		(a) 310 <i>kPa</i> (b) 210 <i>kPa</i>
5	Which of the following statement is correct		(c) $420 \ kPa$ (d) $365 \ kPa$
3.	 (a) In all the three states the molecules possess random translational motion 		(e) 265 kPa
	(b) Gases cannot be converted into solids without passing through		ideal gas equation and related gas laws
	(a) One of the common property of liquids and goess is viscosity	1.	If P , V , T represent pressure, volume and temperature of the gas,
	(c) One of the common property of inquits and gases is viscosity (d) According to Paula's law $1/P_{\rm eff}$ constant at constant T		
6.	A volume of 1 m^3 is equal to () 1000 cm ³		(a) $V \propto \frac{1}{T}$ (at constant <i>P</i>) (b) $PV = RT$
	(a) 1000 cm (b) 100 cm		(c) $V \propto 1 / P$ (at constant <i>T</i>) (d) $PV = nRT$
_	(c) $10 dm^3$ (d) $10^6 cm^3$	2.	At constant temperature, in a given mass of an ideal gas
7.	() Note of the following is not a unit of pressure		[CBSE PMT 1991]
	(a) Newton (b) Forr		(a) The ratio of pressure and volume arways remains constant
	(c) Pascal (d) Bar		(b) volume always remains constant
8.	$1^{o}C$ rise in temperature is equal to a rise of		(c) Pressure always remains constant
	(a) $1^{o} F$ (b) $9 / 5^{o} F$	3.	(d) The product of pressure and volume always remains constant Air at sea level is dense. This is a practical application of
	(c) $5/9^{o}F$ (d) $33^{o}F$		[Kerala CEE 2000]
9.	Which of the following relations for expressing volume of a sample is not correct		(a)Boyle's law(b)Charle's law(c)Avogadro's law(d)Dalton's law
	(a) $1L = 10^3 ml$ (b) $1 dm^3 = 1 L$	4.	If 20 cm^3 gas at 1 <i>atm.</i> is expanded to 50 cm^3 at constant <i>T</i> , then what is the final pressure [CPMT 1988]
	(c) $1L = 10^3 m^3$ (d) $1L = 10^3 cm^3$		(a) $20 \times \frac{1}{20}$ (b) $50 \times \frac{1}{20}$
10.	One atmosphere is numerically equal to approximately		$(0) 50^{-1} \frac{1}{20}$
	(a) $10^6 \text{ dynes } cm^{-2}$ (b) $10^2 \text{ dynes } cm^{-2}$		(c) $1 \times \frac{1}{20} \times 50$ (d) None of these
	(c) $10^4 \text{ dynes } cm^{-2}$ (d) $10^8 \text{ dynes } cm^{-2}$	5.	Which of the following statement is false [BHU 1994]

11. $2gm \text{ of } O_2 \text{ at } 27^{\circ}C \text{ and } 760mm \text{ of } Hg \text{ pressure has volume}[BCECE 2005]$

(a) The product of pressure and volume of fixed amount of a gas is independent of temperature

UNIVERS

- Molecules of different gases have the same K.E. at a given temperature
- The gas equation is not valid at high pressure and low (c) temperature
- (d) The gas constant per molecule is known as Boltzmann constant
- 6. Which of the following graphs represent Boyle's law



- Densities of two gases are in the ratio 1:2 and their temperatures 7. are in the ratio 2 : 1, then the ratio of their respective pressures is [BHU 2000]
 - (a) 1:1 (b) 1:2
 - (c) 2:1 (d) 4:1
- At constant pressure, the volume of fixed mass of an ideal gas is 8. directly proportional to [EAMCET 1985]
 - (a) Absolute temperature (b) Degree centigrade
 - (c) Degree Fahrenheit (d) None
- Which of the following expression at constant pressure represents 9. Charle's law [AFMC 1990]
 - (a) $V \propto \frac{1}{T}$ (b) $V \propto \frac{1}{\tau^2}$
 - (d) $V \propto d$ (c) $V \propto T$
- Use of hot air balloons in sports and meteorological obsevations is 10. an application of [Kerala MEE 2002]
 - (a) Boyle's law (b) Newtonic law (c) Kelvin's law (d) Charle's law
- 11. A 10 g of a gas at atmospheric pressure is cooled from $273^{\circ}C$ to

 $0^{o}C$ keeping the volume constant, its pressure would become

(a)	1/2 <i>atm</i>	(b)	1/273 <i>atm</i>
(c)	2 atm	(d)	273 <i>atm</i>

(a)

13.

Pressure remaining the same, the volume of a given mass of an ideal 12. gas increases for every degree centigrade rise in temperature by definite fraction of its volume at

$0^{o}C$	(b) Its critical temperature	

(c) Absolute zero (d) Its Boyle temperature A certain sample of gas has a volume of 0.2 *litre* measured at 1 *atm*

i certain bain	Sie of guo		ronann			cuo		
pressure and	$0^{o}C$. At	t the	same	pressure	but	at	273°C,	its
volume will be				[EAMO	CET 19	92.	93: BHU 20	005]

(a)	0.4 <i>litres</i>	(b)	0.8 <i>litres</i>	
(c)	27.8 <i>litres</i>	(d)	55.6 <i>litres</i>	

- 400 cm^3 of oxygen at $27^{\circ}C$ were cooled to $-3^{\circ}C$ without 14. change in pressure. The contraction in volume will be
 - (a) 40 cm^3 (b) 30 cm^3
 - (d) 360 cm^3 (c) 44.4 cm^3
- The pressure p of a gas is plotted against its absolute temperature T15. for two different constant volumes, V_1 and V_2 . When $V_1 > V_2$, the
 - (a) Curves have the same slope and do not intersect
 - (b) Curves must intersect at some point other than T = 0
 - (c) Curve for V_2 has a greater slope than that for V_1

- (d) Curve for V_1 has a greater slope than that for V_2
- Two closed vessels of equal volume containing air at pressure P_1 16. and temperature T_1 are connected to each other through a narrow tube. If the temperature in one of the vessels is now maintained at T_1 and that in the other at T_2 , what will be the pressure in the vessels

(a)
$$\frac{2P_1T_1}{T_1 + T_2}$$
 (b) $\frac{T_1}{2P_1T_2}$
2P T 2P

(c)
$$\frac{2T_1T_2}{T_1 + T_2}$$
 (d) $\frac{2T_1}{T_1 + T_2}$

- "One gram molecule of a gas at N.T.P. occupies 22.4 litres." This fact [CPMT 1981, 1995] was derived from
 - (a) Dalton's theory

18.

19.

[CBSE PMT 1989]

- (b) Avogadro's hypothesis
- (c) Berzelius hypothesis
- (d) Law of gaseous volume
- In a closed flask of 5 *litres,* 1.0 g of H_2 is heated from 300 to 600
- K. which statement is not correct [CBSE PMT 1991]
- Pressure of the gas increases (a)
- (b) The rate of collision increases
- The number of moles of gas increases (c)
- The energy of gaseous molecules increases (d)
- Which one of the following statements is false
- [Manipal PMT 1991]
- (a) Avogadro number = 6.02×10^{21}
- (b) The relationship between average velocity (\overline{v}) and root mean square velocity (*u*) is $\overline{v} = 0.9213 u$
- The mean kinetic energy of an ideal gas is independent of the (c) pressure of the gas
- $\left(d\right)$ $% \left(d\right)$ The root mean square velocity of the gas can be calculated by the formula $(3RT/M)^{1/2}$

The compressibility of a gas is less than unity at STP. Therefore [IIT 2000] 20.

- (b) $V_m < 22.4$ litres (a) $V_m > 22.4$ litres
- (d) $V_m = 44.8$ litres (c) $V_m = 22.4$ litres
- In the equation of sate of an ideal gas PV = nRT, the value of the 21. universal gas constant would depend only on
 - [KCET 2005]
 - (a) The nature of the gas
 - The pressure of the gas (b)
 - The units of the measurement (c)
 - (d) None of these
- In the ideal gas equation, the gas constant R has the dimensions of [NCERT 1982 22. mole-atm K (b) *litre mole* (a)
 - (c) litre-atm K mole (d) erg K
- In the equation PV = nRT, which one cannot be the numerical 23. value of R[BIT 1987]
 - (a) $8.31 \times 10^7 erg K^{-1} mol^{-1}$
 - $8.31 \times 10^7 dyne \ cm \ K^{-1} mol^{-1}$ (b)
 - $8.31 JK^{-1}mol^{-1}$ (c)
 - (d) 8.31 *atm*. $K^{-1}mol^{-1}$
- Which one of the following indicates the value of the gas constant R[EAMCET I 24.
 - (a) 1.987 cal K^a mol^a (b) 8.3 cal K mol
 - (c) 0.0821 lit K mol (d) 1.987 Joules K mol [Orissa 1990]
 - The constant R is

25.

(a) Work done per molecule

UNIVERSAL ELF SCORE

UNIVER SELF SC

	(b) Work done per degree absolute	37.	16 g of oxygen and 3 g of hydrogen are mixed and kept at 760 mm
	(c) Work done per degree per mole		pressure and $0^{o}C$. The total volume occupied by the mixture will
	(d) Work done per mole		be nearly [Vellore CMC 1991]
26.	Select one correct statement. In the gas equation, $PV = nRT$ [CBSE PM	AT 1992]	(a) 22.4 <i>litres</i> (b) 33.6 <i>litres</i> (d) 44800 ml
	(a) <i>n</i> is the number of molecules of a gas (b) <i>V</i> denotes volume of one mole of the gas	38.	Pure hydrogen sulphide is stored in a tank of 100 <i>litre</i> capacity at
	(c) n moles of the gas have a volume V	001	$20^{\circ}C$ and 2 atm pressure. The mass of the gas will be [CPMT 1080]
	(d) P is the pressure of the gas when only one mole of gas is		(a) $34 g$ (b) $340 g$
	present		(c) 282.4 g (d) 28.24 g
27.	The correct value of the gas constant <i>R</i> is close to	39.	At N.T.P. the volume of a gas is found to be 273 <i>ml</i> . What will be
	[CBSE PMT 1992]		the volume of this gas at 600 $mm Hg$ and $273^{\circ}C$
			[CPMT 1992]
	(b) 0.082 <i>litre-atmosphere</i> $K^{-1} mol^{-1}$		(a) $391.6 ml$ (b) $380 ml$ (c) $691.6 ml$ (d) $750 ml$
	(c) 0.082 litre-atmosphere ⁻¹ K mole ⁻¹	40.	One <i>litre</i> of a gas weighs 2 g at 300 K and 1 atm pressure. If the
	1 1		pressure is made 0.75 <i>atm</i> , at which of the following temperatures
	(d) 0.082 $litre^{-1} atmosphere^{-1} K mol$		[CBSE PMT 1992]
28.	S.I. unit of gas constant <i>R</i> is [CPMT 1994]		(a) 450 K (b) 600 K
	(a) 0.0821 <i>litre atm K^a mole</i>		(c) 800 K (d) 900 K
	(b) 2 calories K mole	41.	A wheather balloon filled with hydrogen at 1 atm and $27^{o}C$ has
	(c) 8.31 <i>joule K[*] mole</i>		volume equal to 12000 <i>litres</i> . On ascending it reaches a place where
	(d) None		the temperature is $-23^{\circ}C$ and pressure is 0.5 <i>atm</i> . The volume of
29.	Gas equation $PV = nRT$ is obeyed by [BHU 2000]		The Dalloon is
	(a) Only isothermal process (b) Only adiabatic process		(a) 24000 <i>litres</i> (b) 20000 <i>litres</i>
	(c) Both (a) and (b) (d) None of these		(c) 10000 <i>litres</i> (d) 12000 <i>litres</i>
30.	For an ideal gas number of moles per litre in terms of its pressure P ,	42.	The density of a gas at $27^{\circ}C$ and 1 <i>atm</i> is <i>d</i> . Pressure remaining
	[AIEEE 2002]		constant at which of the following temperatures will its density
	(a) <i>PT</i> / <i>R</i> (b) <i>PRT</i>		become 0.75 <i>d</i> [CBSE PMT 1992]
	(c) P/RT (d) RT/P		(a) $20^{\circ}C$ (b) $30^{\circ}C$
31.	If two moles of an ideal gas at 546 <i>K</i> occupy a volume of 44.8 <i>litres</i> , the pressure must be		(c) 400 K (d) 300 K
	[NCERT 1981; JIPMER 1991]	43.	A sample of gas occupies 100 ml at $27^{\circ}C$ and 740 mm pressure.
	(a) 2 <i>atm</i> (b) 3 <i>atm</i>		When its volume is changed to 80 <i>ml</i> at 740 <i>mm</i> pressure, the temperature of the gas will be
	(c) 4 <i>atm</i> (d) 1 <i>atm</i>		[Vellore CMC 1991]
32.	How many moles of <i>He</i> gas occupy 22.4 <i>litres</i> at $30^{\circ}C$ and one		(a) $21.6^{\circ}C$ (b) $240^{\circ}C$
	atmospheric pressure [KCET 1992]		(1) 21000 $(1) 2000$
	(c) 0.11 (d) 1.0	44	$(c) - 55$ C $(d) \delta 9.5$ C The total pressure exerted by a number of non-reacting gases is
33.	Volume of 0.5 <i>mole</i> of a gas at 1 <i>atm.</i> pressure and $273K$ is	44.	equal to the sum of the partial pressures of the gases under the
	[EAMCET 1992]		same conditions is known as [CPMT 1986]
	(a) 22.4 <i>litres</i> (b) 11.2 <i>litres</i> (d) 5.6 <i>litres</i>		(a) Boyle's law (b) Charle's law
	$(c) 44.5 mres \qquad (d) 5.5 mres \qquad (c) b b b b b b b b b $	45	(c) Avogadro's law (d) Dalton's law (d) meressure and pressure
34.	At U C and one <i>atm</i> pressure, a gas occupies 100 <i>cc</i> . If the pressure is increased to one and a half-time and temperature is		contain equal number of particles." This statement is a direct
	increased by one-third of absolute temperature, then final volume of		consequence of [Kerala MEE 2002]
	the gas will be		(a) Avogadro's law (b) Charle's law
	[DCE 2000]	46	(c) recar gas equation (d) Law of partial pressure
	(c) 66.7 cc (d) 100 cc	40.	In three unreactive gases having partial pressures I_A , I_B and I_C and their moles are 1.2 and 2 respectively then their total pressure
35.	Correct gas equation is [CBSE PMT 1989; CPMT 1991]		will be [CPMT 1994]
	(a) $\frac{V_1 T_2}{V_1 T_2} = \frac{V_2 T_1}{V_1 T_2}$ (b) $\frac{P_1 V_1}{V_1 T_2} = \frac{T_1}{T_1}$		$(P_A + P_B + P_C)$
	$\begin{array}{cccc} (a) & P_1 & P_2 \\ \end{array} \qquad \begin{array}{cccc} (b) & P_2 V_2 & T_2 \\ \end{array}$		(a) $P = P_A + P_B + P_C$ (b) $P = \frac{1}{6}$
	(a) $\frac{P_1T_2}{P_1} - \frac{P_2V_2}{P_2}$ (b) $\frac{V_1V_2}{P_1} - P_2P_2$		$\sqrt{P_A + P_B + P_C}$
	(c) $\frac{1}{V_1} - \frac{1}{T_2}$ (d) $\frac{1}{T_1 T_2} - T_1 T_2$		(c) $P = \frac{\sqrt{r_A + r_B + r_C}}{3}$ (d) None
36.	Two separate bulbs contain ideal gases A and B. The density of gas	47.	Dalton's law of partial pressure will not apply to which of the
	A is twice that of gas B. The molecular mass of A is half that of gas B. The two gases are at the same temperature. The setting $f(x)$	т <i>/</i> •	following mixture of gases [Bihar MADT 1981]
	D, the two gases are at the same temperature. The ratio of the pressure of A to that of gas B is		(a) H_2 and SO_2 (b) H_2 and Cl_2
	. [BHU 1994]		(c) H_2 and CO_2 (d) CO_2 and Cl_2
	(a) 2 (b) 1/2		
	(c) 4 (d) $1/4$		

- Which of the following mixtures of gases does not obey Dalton's law 48 of partial pressure [CBSE PMT 1996: Kerala PMT 2000] 60 (a) O_2 and CO_2 (b) N_2 and O_2 (d) NH_3 and HCl(c) Cl_2 and O_2 To which of the following gaseous mixtures is Dalton's law not 49. 61. applicable (a) $Ne + He + SO_2$ (b) $NH_3 + HCl + HBr$ (c) $O_2 + N_2 + CO_2$ (d) $N_2 + H_2 + O_2$ 62. Equal amounts of two gases of molecular weight 4 and 40 are 50. mixed. The pressure of the mixture is 1.1 atm. The partial pressure of the light gas in this mixture is [CBSE PMT 1991] (a) 0.55 atm (b) 0.11 *atm* (d) 0.12 atm (c) 1 *atm* Rate of diffusion of a gas is 51. [IIT 1985; CPMT 1987] (a) Directly proportional to its density (b) Directly proportional to its molecular mass (c) Directly proportional to the square root of its molecular mass 63. (d) Inversely proportional to the square root of its molecular mass Which of the following gas will have highest rate of diffusion 52. [Pb. CET Sample paper 1993; CPMT 1990] (a) NH_3 (b) N_2 64. (c) CO_2 (d) O_2 Which of the following relationship is correct, where r is the rate of 53. diffusion of a gas and d is its density [CPMT 1994] (a) $r \propto \sqrt{1/d}$ (b) $r \propto \sqrt{d}$ (c) r = d(d) $r \propto d$ According to Grahman's law at a given temperature, the ratio of the 54. 65. rates of diffusion r_A / r_B of gases A and B is given by [IIT 1998] (a) $(P_A / P_B)(M_A / M_B)^{1/2}$ (b) $(M_A / M_B)(P_A / P_B)^{1/2}$ 66. (c) $(P_A / P_B)(M_B / M_A)^{1/2}$ (d) $(M_A / M_B)(P_B / P_A)^{1/2}$ 67. (where *P* and *M* are the pressures and molecular weights of gases *A* and *B* respectively) The ratio of the rate of diffusion of a given element to that of 55. helium is 1.4. The molecular weight of the element is 68. [Kerala PMT 1990] (a) 2 (b) 4 (c) 8 (d) 16 A gas diffuse 1/5 times as fast as hydrogen. Its molecular weight is [CPMT 1992; Bih 56. (a) 50 (b) 25 25√2 (d) $50\sqrt{2}$ (c) The molecular weight of a gas which diffuses through a porous plug 57. at 1/6th of the speed of hydrogen under identical conditions is[EAMCET 1990] (a) 27 (b) 72 70. (c) 36 (d) 48 Molecular weight of a gas that diffuses twice as rapidly as the gas 58. with molecular weight 64 is [EAMCET 1994] (a) 16 (b) 8
 - **59.** The densities of hydrogen and oxygen are 0.09 and 1.44 $g L^{-1}$. If the rate of diffusion of hydrogen is 1 then that of oxygen in the same units will be [RPMT 1994]

(d) 6.4

(c) 64

(d) 1/16

(d) 4

- **10.** If rate of diffusion of *A* is 5 times that of *B*, what will be the density ratio of *A* and *B* [AFMC 1994] (a) 1/25 (b) 1/5
 - (a) 1/25 (c) 25
 - (0
- . The densities of two gases are in the ratio of 1 : 16. The ratio of their rates of diffusion is [CPMT 1995] (a) 16 : 1 (b) 4 : 1
 - (a) 16:1 (c) 1:4

(a) 4

(c) 16

- (d) 1:16
- **62.** At constant volume and temperature conditions, the rate of diffusion D_A and D_B of gases A and B having densities ρ_A and ρ_B arerelated by the expression

(a)
$$D_A = \left[D_B \cdot \frac{\rho_A}{\rho_B} \right]^{1/2}$$
 (b) $D_A = \left[D_B \cdot \frac{\rho_B}{\rho_A} \right]^{1/2}$
(c) $D_A = D_B \left(\frac{\rho_A}{\rho_B} \right)^{1/2}$ (d) $D_A = D_B \left(\frac{\rho_B}{\rho_A} \right)^{1/2}$

3. Atmolysis is a process of

- (a) Atomising gas molecules
 - $(b) \quad \text{The breaking of atoms to sub-atomic particles} \\$
 - (c) Separation of gases from their gaseous mixture
- (d) Changing of liquids to their vapour state
- 4. A bottle of ammonia and a bottle of dry hydrogen chloride connected through a long tube are opened simultaneously at both ends, the white ammonium chloride ring first formed will be[IIT 1988]
 - $(a) \quad \text{At the centre of the tube} \\$
 - $(b) \quad \text{Near the hydrogen chloride bottle} \\$
 - (c) Near the ammonia bottle
 - $(d) \quad \text{Throughout the length of the tube} \\$
- 55. Which of the following pairs will diffuse at the same rate through a porous plug [EAMCET 1990]

(a)
$$CO, NO_2$$
 (b) NO_2, CO_2

(c)
$$NH_3, PH_3$$
 (d) NOC_2H_6

- i6. If 4 g of oxygen diffuse through a very narrow hole, how much hydrogen would have diffused under identical conditions [CPMT 1971]
 (a) 16 g
 (b) 1 g
 (c) 1/c
 - (c) 1/4 g (d) 64 g
- A gas diffuse at a rate which is twice that of another gas B. The ratio of molecular weights of A to B is [EAMCET 1986]
 (a) 1.0
 (b) 0.75
 - (c) 0.50 (d) 0.25
- **18.** Two grams of hydrogen diffuse from a container in 10 *minutes*. How many *grams* of oxygen would diffuse through the same container in the same time under similar conditions [MNR 1980] (a) 0.5 g (b) 4 g
 - (d) 8 g
 - The rate of diffusion of methane at a given temperature is twice that of X. The molecular weight of X is
 - [MNR 1995; Kerala CEE 2001]

 (a) 64.0
 (b) 32.0

 (c) 40.0
 (d) 80.0
 - X ml of H_2 gas effuses through a hole in a container in 5 seconds. The time taken for the effusion of the same volume of the gas specified below under identical condition is [11T 1996]
 - (a) 10 seconds : He (b) 20 seconds : O_2
 - (c) 25 seconds : CO (d) 55 seconds : CO_2

times that of SO_2 at $50^o C$

71. At what temperature, the rate of effusion of N_2 would be 1.625

[CBSE PMT 1996]

	(a) 110 K (b) 173 K (c) 373 K (d) 273 K	1	83.	A closed vessel contains equidade molecules at a pressure of P is system then the pressure will be	al nun <i>mm</i> . If	iber of nitro nitrogen is re	gen and oxygen emoved from the
2.	Given the reaction $C(s) + H_2O(l) \rightarrow CO(g) + H_2$	(g) calculate the		system then the pressure win b	C		[MP PMT 1985]
	volume of the gases produced at STP from 48.0 g o	f carbon		(a) <i>P</i>	(b)	2 <i>P</i>	[
	(a) $179.2 L$ (b) $89.6 L$			() D/2	(1)	\mathbf{p}^2	
	(c) 44.8 L (d) 22.4 L 4.4 σ of a gas at STP accupies a volume of 2.24 L t	ha gas can ha litamena (DET o	(c) F/Z	(a)	Г 1 н. н	
	(a) Q_{1} (b) CQ_{2}	ne gas can be[naryana v	Orzącia 2	N_2 O_2 H_2 and N_e separatel	re nileo v then	u to the sam	le pressure with Il be filled first[M ar
					y, then	which one wi	
	(c) NO_2 (d) CO_2			(a) N ₂	(b)	O_2	
	Under what conditions will a pure sample of an id	deal gas not only		(c) <i>H</i> ₂	(d)	Ne	
	litre-1	ration of 1 <i>mole</i>	85.	Which of the following gas mix of partial pressure	ture is	not applicable	e for Dalton's law [Pb. CET 2002]
	$(R = 0.082 litreatmmol^{-1} \deg^{-1})$	[CBSE PMT 1993]		(a) SO_2 and Cl_2	(b)	CO_2 and l	V ₂
	(a) At STP			(c) CO and CO_2	(d)	CO and N	<i>1</i> 2
	(b) When $V = 22.4 \ litres$		0 <i>C</i>	At what pressuret'	(-)	11	2 huma of 60 ml
	(c) When $T = 12 K$	1	80.	At what pressure a quantity of	gas wi	n occupy a vo	nume or $OUml$,
	(d) Impossible under any conditions			if it occupies a volume of 1 (while temperature is constant)	00 <i>ml</i> :	at a pressu	re of 720mm? [Pb. CET 2000]
	There are 6.02×10^{22} molecules each of N_2, O_2	and H_2 which		(a) 700 mm	(b)	800 mm	
	are mixed together at 760 mm and 273 K. The main grams is	ss of the mixture		(c) 100 mm	(d)	1200 mm	
	(a) 6.2 (b) 4.12	[ro.r///1997]	87.	At constant temperature and	pressu	re which gas	will diffuse first
	(c) 3.09 (d) 7			H_2 or O_2 ?		U U	[Pb. CET 2000]
	Volume of 4.4 g of CO_2 at NTP is	[Pb. CET 1997]		(a) Hydrogen			
	(a) 22.4 <i>L</i> (b) 44.8 <i>L</i>			(b) Oxygen			
	(c) 2.24 <i>L</i> (d) 4.48 <i>L</i>			(c) Both will diffuse in same t	ime		
	The energy of an ideal gas depends only on its			(d) None of the above			
	(a) Pressure (b) Volume	1	88.	When a jar containing gaseous	s mixtu	re of equal v	olumes of CO_2
	(c) Number of moles (d) Temperature	2		and H_2 is placed in a solution	on of s	odium hydrox	ide, the solution
	A bottle of cold drink contains 200 <i>ml</i> liquid in w	hich CO_2 is 0.1		level will			[Pb. CET 2001]
	molar. Suppose CO_2 behaves like an ideal gas, the	he volume of the		(a) Rise	(b)	Fall	
	dissolved CO_2 at STP is	[CBSE PMT 1991]		(c) Remain constant	(d)	Become zero	
	(a) 0.224 <i>litre</i> (b) 0.448 <i>litre</i>	1	89.	At S.T.P. $1g CaCO_3$ on decor	npositio	on gives CO ₂	
	(c) 22.4 <i>litre</i> (d) 2.24 <i>litre</i>						[Pb. CET 2000]
	The vapour density of a gas is 11.2. The volume occur	upied by 11.2 g of		(a) 22.4 litre	(b)	2.24 litre	
	this gas at N.T.P. is			(c) 0.224 litre	(d)	11.2 litre	• •
	[MNR 198]	32; CBSE PMT 1991]	90.	At NTP, the density of a gas, wh	ose mo	lecular weight	15 45 15
	(c) $22.4 L$ (d) $20 L$			(a) 44.8 gm/litre	(b)	11.4 gm/litre	[· 0. CD1 2001, 03]
	A pre-weighed vessel was filled with oxygen at N.T	.P. and weighted.		(c) 2 gm/ litre	(d)	3 gm/litre	
	It was then evacuated, filled with SO_2 at the s	ame temperature	91.	What is the ratio of diffusion ra	ate of o	xygen and hyd	lrogen
	and pressure, and again weighted. The weight of oxy	ygen will be[NCERT 1989	9]				[Pb. CET 2003]
	(a) The same as that of SO_2	-		(a) 1:4	(b)	4:1	
	1		~~	(c) 1:8	(d)	8:1	
	(b) $\frac{1}{2}$ that of SO_2		92.	The maximum number of mole	cules is	present in	[CBSE PMT 2004]
	(c) Twice that of <i>SO</i> ₂			(a) 0.5 g of H_2 gas	(b)	10 g of O_2	gas
	(d) One fourth that of SQ			(c) 15 L of H_2 gas at STP	(d)	$5 L \text{ of } N_2$ g	gas at STP
	(u) One fourth that of 50°_{2}	9	93.	One litre oxygen gas at STP wi	l weigh	I	[Pb. CET 2004]
	Five grams each of the following gases at 87° (C and 750 <i>mm</i>		(a) 1.43 g	(b)	2.24 g	
	pressure are taken. Which of them will have the leas	st volume[MNR 1991]	0.4	(c) 11.2 g	(d)	22.4 g	
	(a) <i>HF</i> (b) <i>HCl</i>	9	94.	(a) Fractional distillation tech	or two	gases	[AFMC 2004]
	(c) <i>HBr</i> (d) <i>HI</i>			(b) Grahams law of diffusion	technia	ue	
	Who among the following scientists has not don work on gases	e any important [Bihar MADT 1080]		(c) Osmosis			
	(a) Boyle (b) Charles	[2		(d) Chromatography			
	(c) Avogadro (d) Faradav	9	95.	The rate of diffusion of hydrog	en gas i	S [Add come	
					(1)	[MH CET 20	JU3; PD. CET 2000]
				(a) 1.4 times to 110 gas	(D)	Same as ne	gas

UNIVERSAL SELF SCORER

~	(c) 5 times to He gas (d) 2 times to He gas	1.	Postulate of kinetic theory is [FAMCET 1080]
6.	Hydrogen diffuses six times faster than gas A . The molar mass of		(a) Atom is indivisible
	gas A is [KCET 2004]		(b) Gases combine in a simple ratio
	(a) 72 (b) 6		(c) There is no influence of gravity on the molecules of a gas
_	(c) 24 (d) 36		(d) None of the above
7.	At what pressure will a quantity of gas, which occupies $100 ml$ at	•	(d) None of the above
	a pressue of $720 mm$, occupy a volume of $84 ml$ [DPMT 2004]	4.	(a) There are intermalicular attractions
	(a) 736.18 mm (b) 820.20 mm		(a) There are intermolecular attractions (b) Malaxular have seen identify suburge
	(c) 784.15 mm (d) 857.14 mm		(b) Molecules have considerable volume
8.	Containers A and B have same gases. Pressure, volume and		
	temperature of A are all twice that of B , then the ratio of	•	(d) The velocity of molecules decreases after each collision
	number of molecules of A and B are [AFMC 2004]	3.	In deriving the kinetic gas equation, use is made of the root mean square velocity of the molecules because it is
	(a) 1:2 (b) 2		[Bihar MADT 1080]
	(c) 1:4 (d) 4		(a) The average velocity of the molecules
9.	A mixture of NO_2 and N_2O_4 has a vapour density of 38.3 at		(b) The most probable velocity of the molecules
	300K What is the number of males of NO in $100a$ of the		(c) The square root of the average square velocity of the molecules
	100g of the		(d) The most accurate form in which valegity can be used in these
	$\begin{bmatrix} \text{Kerala PMI 2004} \end{bmatrix}$		calculations
	(a) 0.043 (b) 4.4 (c) 3.4 (d) 3.86	4	Kinetic energy of a gas depends upon its [Bihar MADT 1082]
	(e) 0.437	4.	(a) Molecular mass (b) Atomic mass
)0.	A cylinder of 5 litres capacity, filled with air at NTP is connected		(c) Fauivalent mass (d) None of these
	with another evacuated cylinder of 30 litres of capacity. The	F	The kinetic theory of gases perdicts that total kinetic operation of a
	resultant air pressure in both the cylinders will be [BHU 2004]	5.	gaseous assembly depends on [NCERT 1984]
	(a) 10.8 cm of Hg (b) 14.9 cm of Hg		(a) Pressure of the gas
ท	(c) 21.6 cm of Hg (d) 36.8 cm of Hg A certain mass of gas occupies a volume of 300 c c at 27 C and 620		(b) Temperature of the gas
••	The first structure of 500 c.c. at 27° and 020		(c) Volume of the gas
	mm pressure. The volume of this gas at 47 C and 640 mm		(d) Pressure volume and temperature of the gas
	(a) 400 cc (b) 510 cc	6	According to kinetic theory of gases the energy per mole of a gas is
	(c) 310 c.c. (d) 350 c.c.	0.	equal to [EAMCET 1985]
)2.	What will be the volume of the mixture after the reaction?		(a) 1.5 <i>RT</i> (b) <i>RT</i>
	$NH_3 + HCl \rightarrow NH_4Cl$ [BVP 2004]		(c) 0.5 <i>RT</i> (d) 2.5 <i>RT</i>
	4 litre 1.5 litre (solid)	7.	Internal energy and pressure of a gas per unit volume are related as[CBSE]
	(a) 0.5 litre (b) 1 litre (c) 2.5 litre		(a) $P = \frac{2}{2}F$ (b) $P = \frac{3}{2}F$
	(c) 2.5 ntre (d) 0.1 ntre		(a) $I = \frac{1}{3}E$ (b) $I = \frac{1}{2}E$
)3.	The pressure and temperature of $4dm^3$ of carbon dioxide gas are		1
	doubled. Then the volume of carbon dioxide gas would be KCET 2004		(c) $P = \frac{1}{2}E$ (d) $P = 2E$
	(a) $2 dm^3$ (b) $3 dm^3$	0	The translational linetic energy of an ideal gas depends only on its
	(c) $4 dm^3$ (d) $8 dm^3$	ð.	(a) Pressure (b) Force
м	If the absolute temperature of an ideal gas become double and		(a) Tressure (b) Torce (d) Molar mass
4.	pressure become half, the volume of gas would be	٩	Helium atom is two times beavier than a hydrogen molecule at 298
	[Kerala CET 2005]		<i>K</i> , the average kinetic energy of helium is [11T 1982]
	(a) Remain unchange (b) Will be double		(a) Two times that of a hydrogen molecule
	(c) Will be four time (d) will be half		(b) Same as that of a hydrogen molecule
	(e) Will be one fourth		(c) Four times that of a hydrogen molecule
5.	At what temperature, the sample of neon gas would be heated to		(d) Half that of a hydrogen molecule
	double of its pressure, if the initial volume of gas is/are reduced to	10.	Which of the following is valid at absolute zero
	15% at 75° C [Kerala CET 2005]		[Pb. CET 1985]
	(a) $210^{\circ}C$ (b) $502^{\circ}C$		(a) Kinetic energy of the gas becomes zero but the molecular
	(a) 519 C (b) 592 C		(b) Kingtig anarow of the gas becomes zero and melocular motion
	(c) $128^{\circ}C$ (d) $60^{\circ}C$		also becomes zero
	(e) $90^{\circ}C$		(c) Kinetic energy of the gas decreases but does not become zero
6.	Equation of Boyle's law is [DPMT 2005]		(d) None of the above
	dP dV dP dV	n.	The average <i>K.E.</i> of an ideal gas in <i>calories</i> per mole is
	(a) $\frac{\alpha r}{p} = -\frac{\alpha r}{V}$ (b) $\frac{\alpha r}{p} = +\frac{\alpha r}{V}$		approximately equal to [EAMCET 1989]
	p v P V		(a) Three times the absolute temperature
	$d^2 P = dV$ $d^2 P = d^2 V$		(b) Absolute temperature
	(c) $\frac{P}{P} = -\frac{dT}{dT}$ (d) $\frac{P}{P} = +\frac{dT}{dT}$		(c) Two times the absolute temperature
	* ***		(d) 1.5 times the absolute temperature

Kinetic molecular theory of gases and **Molecular collisions**

UNIVERSAL

12. According to kinetic theory of gases, for a diatomic molecule

[MNR 1991]



(a) T	he pressure exerted by the gas is proportional to the mean		(a) 298 K
(b) T	The pressure exerted by the gas is proportional to the root	21.	Ratio of C_p
(c) T	The root mean square velocity is inversely proportional to the emperature		gas 'X prese
(d) T	The mean translational kinetic energy of the molecules is romortional to the absolute temperature		(a) 6.02>
At STI	P, 0.50 mol H_2 gas and 1.0 mol He gas		(c) 3.01
	[CBSE PMT 1993, 2000]	22.	The density $(a) = 0.0006$
(a) H	lave equal average kinetic energies		(c) 14.4816
(b) F (c) C	lave equal molecular speeds Decupy equal volumes	23.	At $100^{\circ}C$
(d) +	lave equal effusion rates	-3.	and that of
Which	of the following expressions correctly represents the		occupied by
relatio	nship between the average molar kinetic energy, $K.E.$, of CO		is
and <i>I</i>	V_2 molecules at the same temperature		(a) 6 cm^3
			(c) 0.6 <i>cn</i>
(a) .	$KE_{CO} = KE_{N_2}$	24.	The ratio γ
(b) .	$\overline{KE}_{CO} > \overline{KE}_{N_2}$		(a) 1.33
(c)	$\overline{KF}_{CO} < \overline{KF}_N$		(c) 2.13
(d) (d)	Cannot be predicted unless the volumes of the gases are given	25.	The density
(u) C Indica	the the correct statement for a 1-L sample of $N_2(q)$ and		(a) S.T.P.
COal	(a) at 208 K and 1 atm pressure		(c) 273°
()		26.	Absolute ze
(a) I	he average translational <i>KE</i> per molecule is the same in N_2		
a	nd CO ₂		(a) At whi
(b) Т	The rms speed remains constant for both N_2 and CO_2		(b) At whi
(c) T	The density of N_2 is less than that of CO_2		(c) At whi (d) All of t
(d) Т	The total translational KE of both N_2 and CO_2 is the same	27.	Consider the
With i	ncrease of pressure, the mean free path		(1) Joule-T
	[Pb. CET 1985]		(2) А пез
(a) [Decreases (b) Increases		(2) The te
(c) L Which	Ones not change (d) Becomes zero		observe
of an	increase in temperature on the distribution of molecular		Which of th
speeds	in a gas [AIEEE 2005]		(a) 1 and 2
(a) T	The most probable speed increases	28.	Vibrational
(b) I	he fraction of the molecules with the most probable speed		(a) Partiall
(c) T	he distribution becomes broader		(b) Only p
(d) T	he area under the distribution curve remains the same as		(c) Only k
f D 1	under the lower temperature	20	(d) None of At the same
and g	as constant respectively, then for an ideal gas, the density is	-).	will have the
given	[CBSE PMT 1989, 91]		() 1
(a) -	$\frac{RT}{P}$ (b) $\frac{P}{P}$		(a) Hydrog
(-)	PM RT	30.	Dimensions
(c) -	$\frac{M}{d}$ (d) $\frac{PM}{d}$	•••	
	V RT		(a) Energy
An ide	al gas will have maximum density when [CPMT 2000] P = 0.5 atm T = 600 V	01	(c) Energy
(a) .	$\mathbf{r} = 0.3 a i m, \mathbf{I} = 000 \mathbf{K}$	JI.	molecular m
(b)	P = 2 atm, T = 150 K		
(c) .	P = 1 atm, T = 300 K		(a) 3 <i>M</i>
(d)	P = 1.0 atm, T = 500 K		(c) $M/3$
1f the	inversion temperature of a gas is $-80^{o}C$, then it will		
produ	ce cooling under Joule-Thomson effect at		

13.

14.

15.

16.

17.

18.

19.

20.

	(a) 298 K (c) 193 K	(b) (d)	273 K 173 K
21	Batio of C and C of a gas '	(u) Kisi	4. The number of stoms of the
21.	$ratio of C_p$ and C_v of a gas r	13 I. 14 N T	TP is
	gas A present in n.2 <i>intres</i> of it a	1L IN. I	[CBSE PMT 1989]
	(a) 6.02×10^{23}	(b)	1.2×10^{24}
	(c) 3.01×10^{23}	(d)	2.01×10^{23}
22.	The density of air is 0.00130 g/m	. The	e vapour density of air will be[DCE 2000]
	(a) 0.00065	(b)	0.65
	(c) 14.4816	(d)	14.56
23.	At $100^{\circ}C$ and 1 <i>atm</i> , if the de	ensity	of liquid water is 1.0 $g \ cm^{-3}$
	and that of water vapour is occupied by water molecules in is	0.000 1 <i>litr</i>	6 $g m^{-3}$, then the volume e of steam at that temperature [IIT 2000]
	(a) 6 cm^3	(b)	$60 \ cm^3$
	(c) 0.6 cm^3	(d)	$0.06 \ cm^3$
	The action of fear in cost access in	(u)	
24.	The ratio γ for mert gases is	(h)	[AFMC 1990]
	(a) 1.33	(d)	1.00
25	The density of neon will be high	est at	CRSF PMT 1000]
-3.	(a) S.T.P.	(b)	$0^{\circ}C, 2 atm$
	(c) $273^{\circ}C \ 1 atm$	(d)	$273^{\circ}C$ 2 atm
26	Absolute zero is defined as the t	(4)	
20.	Absolute zero is defined as the ti	empe	[CRSE PMT 1000]
27.	 (a) At which all molecular motified (b) At which liquid helium boils (c) At which ether boils (d) All of the above Consider the following statement 	s ts :	rases
	(1) Joule-Thomson experiment	is iso	enthalpic as well as adiabatic.
	(2) A negative value of	u_{JT}	(Joule Thomson coefficient
	(3) The temperature at which	a gas neith	on expansion. er cooling nor heating effect is
	observed is known as invers	sion t	emperature.
	(a) 1 and 2	(b)	1 and 3
	(c) 2 and 3	(d)	1, 2 and 3
28.	Vibrational energy is		[Pb. CET 1985]
	(a) Partially potential and parti	ally k	inetic
	(b) Only potential		
	(c) Only kinetic		
20	(d) None of the above		a which of the following gases
29.	will have the highest kinetic ener	essui ev de	er mole
	5	0, 1	[MNR 1991]
	(a) Hydrogen	(b)	Oxygen
	(c) Methane	(d)	All the same
30.	Dimensions of pressure are the s	ame	as that of
	(a) Energy	(b)	[CBSE PMT 1995]
	(c) Energy ner unit volume	(d)	Force per unit volume
31.	The density of a gas An is the	nree 1	times that of a gas <i>B</i> if the
.	molecular mass of A is M , the m	olecu	lar mass of <i>B</i> is
			[CPMT 1987]
	(a) 3 <i>M</i>	(b)	$\sqrt{3} M$
	(c) $M/3$	(J)	$M/\sqrt{3}$
		(u)	

Molecular speeds

SELF	242 Gaseous state		
I .	The ratio of root mean square velocity to average velocity of gas molecules at a particular temperature is [JIT 198 1]		(a) $\sqrt{\frac{3P}{d}}$
	(a) 1.086 : 1 (b) 1 : 1.086		1 a
	(c) 2:1.086 (d) 1.086:2		(c) $\sqrt{\frac{3RT}{K}}$
	Which is not true in case of an ideal gas [CBSE PMT 1991]		
	(a) It cannot be converted into a liquid	12.	Koot mean so
	(b) There is no interaction between the molecules		(a) $m^{1/2}$
	(c) All molecules of the gas move with same speed		(c) $m^{-1/2}$
	(d) At a given temperature, PV is proportional to the amount of the gas	13.	At constant
	The ratio among most probable velocity, mean velocity and root mean square velocity is given by [CBSE PMT 1993]		(a) Increase (b) Increase
	(a) $1 \cdot 2 \cdot 3$ (b) $1 \cdot \sqrt{2} \cdot \sqrt{3}$		(c) Increase
		14.	(d) Decreas
	(c) $\sqrt{2}:\sqrt{3}:\sqrt{8}/\pi$ (d) $\sqrt{2}:\sqrt{8}/\pi:\sqrt{3}$	-	u_1 and u_2 .
	Which of the following has maximum root mean square velocity at the same temperature [Manipal PMT 2002]		the following
	(a) SO_2 (b) CO_2		(a) $\frac{m_1}{m_1} = \frac{m_1}{m_1}$
	(c) O_2 (d) H_2		u_1^2
	The temperature at which RMS velocity of SO_2 molecules is half		(c) $\frac{m_1}{m_1} = \frac{m_1}{m_1}$
	that of <i>He</i> molecules at 300 <i>K</i> is [NTSE 1991]		u_1
	(a) 150 K (b) 600 K	15.	The temperat
	(c) 900 K (d) 1200 K		root mean sq
	At $27^{o}C$, the ratio of <i>rms</i> velocities of ozone to oxygen is		(a) $\sqrt{927}$ /
	[EAMCET 1992]		(b) Same as
	(a) $\sqrt{3/5}$ (b) $\sqrt{4/3}$		(c) Halved (d) Doubled
	(c) $\sqrt{2/3}$ (d) 0.25	16.	The ratio bet
	The average kinetic energy of an ideal gas per molecule in SI units at		that of O_2 a
	$25^{\circ}C$ will be [CBSE PMT 1996]		(a) 4 (c) 1
	$() (17 \times 10^{-21} \text{ hJ}) (1) (17 \times 10^{-21} \text{ J})$	17.	The root me
	(a) $6.1/\times 10$ KJ (b) $6.1/\times 10$ J		varies density
	(c) $6.17 \times 10^{-20} J$ (d) $7.16 \times 10^{-20} J$		(a) d^2_{-}
	At what temperature the <i>RMS</i> velocity of SO_2 be same as that of		(c) \sqrt{d}
	O_2 at 303 K [KCET 2001]	18.	Consider a n
	(a) $273 K$ (b) $606 K$		Compared to
	(c) $303 K$ (d) $403 K$		wall with
	Among the following gases which one has the lowest root mean		(a) Smaller (c) Greater
	square velocity at $25^{\circ}C$ [EAMCET 1983]	19.	The <i>rms</i> spe
	(a) SO_2 (b) N_2		doubled and
	(c) Q_2 (d) Cl_2		the <i>rms</i> spee
	The root mean square velocity of an ideal gas in a closed container		(a) $u/2$ (c) $4u$
	of fixed volume is increased from $5 \times 10^4 \text{ cm s}^{-1}$ to	20.	Choose the c
			meanings
	10×10^4 cm s ⁻¹ . Which of the following statement correctly		(a) $u > u_p$
	explains how the change is accomplished		(c) $u_p > \overline{u}$
	(a) By heating the gas, the temperature is doubled	21.	The ratio of
	(b) By heating the gas, the pressure is quadrupled (<i>i.e.</i> made four		(a) $\pi/2$
	times)		(c) $\sqrt{\pi}/2$
	(c) By heating the gas, the temperature is quadrupled	22.	The r.m.s.
	(d) By heating the gas, the pressure is doubled		temperature,
	The <i>rms</i> velocity at NTP of the species can be calculated from the expression [FAMCET 1000]		(a) 1200 <i>K</i>

$$\frac{3P}{d}$$
 (b) $\sqrt{\frac{3PV}{M}}$

c)
$$\sqrt{\frac{3 R I}{M}}$$
 (d) All the above

uare velocity of a gas molecule is proportional to[CBSE PMT 1990]

- (b) m^0
- (d) *m*
- volume, for a fixed number of moles of a gas, the ne gas increases with increase in temperature due to[11T 1992] in the average molecular speed
 - d rate of collision amongst molecules
 - in molecular attraction
 - e in mean free path
- locities of the two gases at the same temperature are Their masses are m_1 and m_2 respectively. Which of expressions is correct

[BHU 1994]

(a)
$$\frac{m_1}{u_1^2} = \frac{m_2}{u_2^2}$$
 (b) $m_1 u_1 = m_2 u_2$
(c) $\frac{m_1}{u_1} = \frac{m_2}{u_2}$ (d) $m_1 u_1^2 = m_2 u_2^2$

- ture of the gas is raised from $27^{o}C$ to $927^{o}C$, the [CBSE PMT 1994] uare velocity is
 - 27 times the earlier value
 - before
- ween the root mean square velocity of H at 50 K and nt 800 *K* is [IIT 1996]

an square velocity of an ideal gas at constant pressure (*d*) as [IIT 2000]

(a)
$$d^2$$
 (b) d
(c) \sqrt{d} (d) $1/\sqrt{d}$

- nixture of SO_2 and O_2 kept at room temperature. the oxygen molecule, the SO_2 molecule will hit the average speed (b) Greater average speed
 - kinetic energy (d) Greater mass
- ed of N_2 molecules in a gas is u. If the temperature is the nitrogen molecules dissociate into nitrogen atoms, d becomes (h) 2u

)
$$4u$$
 (d) $14u$

prrect arrangement, where the symbols have their usual

(a)
$$u > u_p > u_{rms}$$

(b) $u_{rms} > u > u_p$
(c) $u_p > \overline{u} > u_{rms}$
(d) $u_p > u_{rms} > \overline{u}$

- nost probable velocity to that of average velocity is [JEE Orissa 200 (b) $2/\pi$
 - (d) $2/\sqrt{\pi}$
- velocity of a certain gas is v at 300K. The at which the r.m.s. velocity becomes double [Pb. CET 2002]
 - (b) 900 K (a) 1200K
 - (c) 600*K* (d) 150K
- The r.m.s. velocity of a gas depends upon [DCE 2002] 23. (a) Temperature only

- (b) Molecular mass only
- (c) Temperature and molecular mass of gas
- (d) None of these
- 24. What is the pressure of 2 mole of NH_3 at $27^{\circ}C$ when its volume is 5 litre in vander Waal's equation (a = 4.17, b = 0.03711)[JEE Orissa,2004] (a) 10.33 atm (b) 9.33 atm (c) 9.74 atm (d) 9.2 atm
- 25. The root mean square velocity of one mole of a monoatomic having molar mass M is U_{ms} . The relation between the average kinetic energy (E) of the U_{ms} is

۵

10.

11.

12.

14.

16.

[CPMT 1991]

(a)
$$U_{mss} = \sqrt{\frac{3E}{2M}}$$
 (b) $U_{mss} = \sqrt{\frac{2E}{3M}}$
(c) $U_{mss} = \sqrt{\frac{2E}{M}}$ (d) $U_{mss} = \sqrt{\frac{E}{3M}}$

26. Ratio of average to most probable velocity is

					[Orissa JEE	2005]
(a)	1.128		(b)	1.224		
(c)	1.0		(d)	1.112		

27. If the v_{ms} is $30R^{1/2}$ at $27^{\circ}C$ then calculate the molar mass of gas in kilogram. [DPMT 2005] (a) 1 (b) 2 (c) 4 (d) 0.001

Real gases and Vander waal's equation

1.	The Vander Waal's equation	on explains the behaviour of	
			[DPMT 1981]
	(a) Ideal gases	(b) Real gases	

(u)	lacal gases	(0)	ricul guses
(c)	Vapour	(d)	Non-real gases

Gases deviate from the ideal gas behaviour because their molecules[NCERT 1981]
 (a) Possess negligible volume

- (b) Have forces of attraction between them
- (c) Are polyatomic
- (d) Are not attracted to one another
- **3.** The compressibility factor of a gas is defined as Z = PV/RT. The compressibility factor of ideal gas is

[Pb. CET 1986] (a) 0 (b) Infinity (c) 1 (d) -1

4. In Vander Waal's equation of state for a non-ideal gas, the term that accounts for intermolecular forces is
[CBSE PMT 1990]. IIT 1988]

(a)
$$(V-b)$$
 (b) $(RT)^{-1}$
(c) $\left(P + \frac{a}{V^2}\right)$ (d) RT

5. Vander Waal's equation of state is obeyed by real gases. For *n* moles of a real gas, the expression will be

[IIT 1992; Pb. CET 1986; DPMT 1986]

(a)
$$\left(\frac{P}{n} + \frac{na}{V^2}\right) \left(\frac{V}{n-b}\right) = RT$$

(b) $\left(P + \frac{a}{V^2}\right) (V-b) = nRT$
(c) $\left(P + \frac{na}{V^2}\right) (nV-b) = nRT$
(d) $\left(P + \frac{n^2a}{V^2}\right) (V-nb) = nRT$

6. Any gas shows maximum deviation from ideal gas at

(a) $0^{o}C$ and 1 atmospheric pressure

- (b) $100^{o}C$ and 2 atmospheric pressure
- (c) $-100^{\circ} C$ and 5 atmospheric pressure
- (d) $500^{\circ}C$ and 1 atmospheric pressure
- The temperature at which the second virial coefficient of real gas is zero is called [AFMC 1993] (a) Critical temperature
- (b) Eutetic point
- (c) Boiling point
- (d) Boyle's temperature
- **8.** When is deviation more in the behaviour of a gas from the ideal gas equation PV = nRT

[DPMT 1981; NCERT 1982; CBSE PMT 1993]

[CPMT 1996]

- (a) At high temperature and low pressure
- (b) At low temperature and high pressure
- (c) At high temperature and high pressure
- (d) At low temperature and low high pressure
- Vander Waal's constants 'a' and 'b' are related with..... respectively[**RPMT 1994**]
 - (a) Attractive force and bond energy of molecules
 - (b) Volume and repulsive force of molecules
 - (c) Shape and repulsive forces of molecules
 - (d) Attractive force and volume of the molecules
- Gas deviates from ideal gas nature because molecules
 - ()
 - (a) Are colourless
 - (b) Attract each other
 - (c) Contain covalent bond
 - (d) Show Brownian movement
- The Vander Waal's equation reduces itself to the ideal gas equation at [Kerala MEE 2001; CBSE PMT 2002]
- (a) High pressure and low temperature
- (b) Low pressure and low temperature
- (c) Low pressure and high temperature
- (d) High pressure and high temperature
- The compressibility factor for an ideal gas is [11T 1997]
- (a) 1.5 (b) 1.0(c) 2.0 (d) ∞
- (c) 2.0 (d) ∞
 13. When an ideal gas undergoes unrestrained expansion, no cooling occurs because the molecules [11T 1984, 89]
 - (a) Are above the inversion temperature
 - (b) Exert no attractive force on each other
 - (c) Do work equal to loss in kinetic energy
 - (d) Collide without loss of energy
 - A gas is said to behave like an ideal gas when the relation $PV/T = {\rm constant}$. When do you expect a real gas to behave like an ideal gas

[IIT 1999; CBSE PMT 1990; CPMT 1991]

[AFMC 1993; IIT 1981, 94]

- (a) When the temperature is low
- (b) When both the temperature and pressure are low
- (c) When both the temperature and pressure are high
- (d) When the temperature is high and pressure is low
- 15. A real gas most closely approaches the behaviour of an ideal gas at[KCET 1992]
 - (a) 15 *atm* and 200 *K* (b) 1 *atm* and 273 *K*
 - (c) 0.5 *atm* and 500 *K* (d) 15 *atm* and 500 *K*
 - The temperature at which real gases obey the ideal gas laws over a
 - wide range of pressure is called
 - (a) Critical temperature
 - (b) Boyle temperature
 - (c) Inversion temperature

UNIVERSAL

244 Gaseous state

Reduced temperature

At low pressure, the Vander Waal's equation is reduced to 17.

(a)
$$Z = \frac{pV_m}{RT} = 1 - \frac{ap}{RT}$$
 (b) $Z = \frac{pV_m}{RT} = 1 + \frac{b}{RT}p$
(c) $pV_m = RT$ (d) $Z = \frac{pV_m}{RT} = 1 - \frac{a}{RT}$

18. At high temperature and low pressure, the Vander Waal's equation is reduced to

(a)
$$\left(p + \frac{a}{V_m^2}\right)(V_m) = RT$$

(b)
$$pV_m = RT$$

(c)
$$p(V_m - b) = RT$$

(d)
$$\left(p + \frac{a}{V_m^2}\right)(V_m - b) = RT$$

- When helium is allowed to expand into vacuum, heating effect is 19. [CPMT 1987] observed. Its reason is that
 - (a) Helium is an ideal gas
 - (b) Helium is an inert gas
 - (c) The inversion temperature of helium is very low
 - The boiling point of helium is the lowest among the elements (d)
- 20. In van der Waal's equation of state of the gas law, the constant 'b' [AIEEE 2004] is a measure of
 - (a) Volume occupied by the molecules
 - Intermolecular attraction (b)
 - (c) Intermolecular repulsions
 - Intermolecular collisions per unit volume (d)
- In which molecule the vander Waal's force is likely to be the most 21. important in determining the m.pt. and b.pt.

(a)	нс	(1-) Br	
(a)	$\pi_2 s$	(D) Dr_2	

- (c) HCl (d) *CO*
- Pressure exerted by 1 mole of methane in a 0.25 litre container at 22. 300K using vander Waal's equation (given $1 = 2.253 atml^2 mol^{-2}, b = 0.0428 litmol^{-1})$ is

[Orissa JEE 2005]

(a) 82.82 atm (b) 152.51 atm (c) 190.52 atm (d) 70.52 atm

Critical state and Liquefaction of gases

- Which set of conditions represents easiest way to liquefy a gas[NCERT 1983] 1. (a) Low temperature and high pressure
 - (b) High temperature and low pressure
 - (c) Low temperature and low pressure
 - (d) High temperature and high pressure
- Adiabatic demagnetisation is a technique used for 2.
 - (a) Adiabatic expansion of a gas
 - (b) Production of low temperature
 - (c) Production of high temperature
 - (d) None
- An ideal gas can't be liquefied because [CBSE PMT 1992] 3
 - (a) Its critical temperature is always above $0^{o}C$

- Its molecules are relatively smaller in size (b)
- It solidifies before becoming a liquid (c)
- (d) Forces operative between its molecules are negligible
- However great the pressure, a gas cannot be liquefied above its
 - Boyle temperature (a)
 - Inversion temperature (b)
 - (c) Critical temperature
 - (d) Room temperature
- An ideal gas obeying kinetic theory of gases can be liquefied if[CBSE PMT 1995]
 - Its temperature is more than critical temperature T_c (a)
 - (b) Its pressure is more than critical pressure P_c
 - Its pressure is more than P_c at a temperature less than T_c (c)
 - (d) It cannot be liquefied at any value of P and T

6. The Vander Waal's parameters for gases W, X, Y and Z are b (L mol Gas a (atm L' mol·) w 4.0 0.027 х 8.0 0.030 γ 6.0 0.032 Z 12.0 0.027 Which one of these gases has the highest critical temperature (a) W (b) X (d) (c) *Y* ZThe Vander Waal's constant 'a' for the gases O_2, N_2, NH_3 and CH_4 are 1.3, 1.390, 4.170 and 2.253 $L^2 atmmol^{-2}$ respectively. The gas which can be most easily liquefied is [IIT 1989] (a) O_2 (b) N_{2} (c) NH_3 (d) CH_A 8. A gas can be liquefied [AFMC 2005] (a) Above its critical temperature At its critical temperature (b) (c) Below its critical temperature (d) At any temperature 9. Which of the following is correct for critical temperature (a) It is the highest temperature at which liquid and vapour can coexist Beyond the critical temperature, there is no distinction between (b) the two phases and a gas cannot be liquefied by compression (c) At critical temperature (T_c) the surface tension of the system is zero (d) At critical temperature the gas and the liquid phases have different critical densities A gas has a density of 2.68 g/L at stp. Identify the gas 10. (a) NO_2 (b) *Kr* (c) COS (d) SO_2 Weight of 112 *ml* of oxygen at NTP on liquefaction would be[DPMT 1984] (b) 0.64 g (a) 0.32 g (c) 0.16 g (d) 0.96 g Critical Thinking **Objective Questions** As the temperature is raised from $20^{\circ}C$ to $40^{\circ}C$ the average

1. kinetic energy of neon atoms changes by a factor of which of the [AIEEE 2004] following

(a) 313/293

 $\sqrt{(313/293)}$ (b)

7.

Δ.

5.

[DPMT 2000]

[BHU 1984]

				Contraction
(c) 1/2 (d) 2	2	11.	If C_1, C_2, C_3 represent the	The speeds of n_1, n_2, n_3 molecules,
A gas is found to have a formula $\left[CO\right]_{x}$. If its vapour density is 70,		then the root mean square spee	ed is [11T 1993]
the value of x is	[DCE 2004]		$(-\alpha^2 + \alpha^2 - \alpha^2)$	$\sum^{1/2}$
(a) 2.5 (b) 3	3.0		(a) $\left n_1 C_1^2 + n_2 C_2^2 + n_3 C_3^2 + n_3 C_3$	+
(c) 5.0 (d) 6	5.0		$(n_1 + n_2 + n_3 + \dots)$)
Which of the given sets of temperature a to exhibit the greatest deviation from ide	and pressure will cause a gas eal gas behavior [DCE 2003]		(b) $\frac{(n_1C_1^2 + n_2C_2^2 + n_3C_3^2 + n_3C_3^2)}{(n_1C_1^2 + n_2C_2^2 + n_3C_3^2)}$) ^{1/2}
(a) $100^{\circ} C$ and 4 atm (b)	$100^{\circ} C$ and 2 atm		$n_1 + n_2 + n_3 + \dots$	
(c) $-100^{\circ} C$ and 4 atm (d)	$0^{o} C$ and 2 atm		(c) $\frac{(n_1C_1^2)^{1/2}}{(n_1C_2^2)^{1/2}} + \frac{(n_2C_2^2)^{1/2}}{(n_1C_2^2)^{1/2}}$	$+ \frac{(n_3C_3^2)^{1/2}}{1} + \dots$
The molecular weight of O_2 and SO_2	are 32 and 64 respectively.		n_1 n_2	n ₃
If one <i>litre</i> of O_2 at 15^oC and 750	0 mm pressure contains 'N		$(n_1C_1 + n_2C_2 + n_3C_3 + n_3C_3)$	$+ \dots)^2]^{1/2}$
molecules, the number of molecules in t	wo <i>litres</i> of SO_2 under the		(d) $(n_1 + n_2 + n_3 +)$)
same conditions of temperature and pres	ssure will be [CBSE 1990; MNR 19	91]		
(a) $N/2$ (b) /	N	12.	20 <i>ml</i> of hydrogen diffuses out	through a small hole from a vessel in or 40 <i>ml</i> of oxygen to diffuse out is CBSE
$(c) 2N \qquad (d) $	· 1 N/		(a) 12 <i>min</i>	(b) 64 min
$(c) = 2iv \qquad (d) = 4$	ни 1 ч. ()		(c) 8 min	(d) 32 min
What is the relationship between the averaginary velocity (u) and most probable u''	erage velocity (<i>v</i>), root mean	10	At what termometers will al	warage append of CH melecular he
square velocity (a) and most probable w	[AEMC 1004]	13.	At what temperature will the a	verage speed of $C\pi_4$ molecules have
()	[רימי שמאר ביי איירי]		the same value as O_2 has at 3	,00 K
(a) $\alpha: v: u::1:1.128:1.224$				[CBSE PMT 1989]
(b) $\alpha: v: u:: 1.128: 1: 1.224$			(a) 1200 K (a) 600 K	(D) 150 K (d) 200 K
(c) $\alpha: v: u:: 1.128: 1.224: 1$			(c) 000 K	(d) 300 K
(1) a		14.	A sample of O_2 gas is col	llected over water at $23^{o}C$ at a
(u) $u \cdot v \cdot u = 1.124 \cdot 1.228 \cdot 1$, 10, 0 .1 i		barometric pressure of 751 <i>m</i>	nm Hg (vapour pressure of water at
Consider the following statements : F	or diatomic gases, the ratio		$23^{o}C$ is 21 <i>mm Hg</i>). The	partial pressure of O_2 gas in the
C_p / C_v is equal to			sample collected is	[CBSE PMT 1993]
(1) 1.40 (lower temperature)			(a) 21 <i>mm Hg</i>	(b) 751 <i>mm Hg</i>
(2) 1.66 (moderate temperature)			(c) 0.96 <i>atm</i>	(d) 1.02 <i>atm</i>
(3) 1.29 (higher temperature)		۱Ľ	In an experiment during the a	nalveis of a carbon compound 145 1
which of the above statements are correc	ct		of H	$\frac{1}{270}$
(a) 1, 2 and 3 (b) 1	and 2		or π_2 was collected at 70	by min rig pressure and 27 C
(c) 2 and 3 (d) 1	and 3		temperature. The mass of H_2	is nearly
The compressibility factor for an ideal ga	as is [MP PET 2004]			[MNR 1987]
(a) 1.5 (b) 1	.0		(a) 10 g	(b) 12 g
(c) 2.0 (d)	∞		(c) 24 g	(d) 6 g
The compressibility factor of a gas is le	ess than 1 at STP. Its molar	16.	The volume of $1 g$ each of	methane (CH_4) , ethane (C_2H_6) ,
volume V_m will be	[MP PET 2004]		propane (C_2H_0) and hutane	(C_4H_{10}) was measured at 350 K
(a) $V_m > 22.42$ (b)	<i>V_m</i> < 22.42		and 1 <i>atm.</i> What is the volume	me of butane [NCERT 1981]
(c) $V_m = 22.42$ (d) N	None		(a) 495 cm^3	(b) 600 cm^3
If some <i>moles</i> of On diffuse in 18 sec.	and same <i>moles</i> of other gas		(c) 000 cm^3	(d) $1700 \ cm^3$
diffuse in 45 <i>sec</i> then what is the molec	cular weight of the unknown [CPMT 1988]	17.	The ratio of the rate of diffu	usion of helium and methane under
	[(a) 4	(b) 2
(a) $\frac{45^2}{2} \times 32$ (b)	$\frac{18^2}{2} \times 32$		(a) 4	(0) 2 (1) 2 (2)
18 ²	45 ²		(c) I	(d) 0.5
18^{2}	45^{2}	18.	At what temperature in the c	elsius scale, V (volume) of a certain
(c) $\frac{10}{45^2 \times 32}$ (d)	$\frac{10}{18^2 \times 32}$		mass of gas at $27^{o}C$ will constant	ll be doubled keeping the pressure [Orissa 1993]
The ratio of rates of diffusion of SO_2 , O_2	\mathcal{O}_2 and CH_4 is		(a) $54^{0}C$	(b) $327^{\circ}C$
-	[BHU 1992]		(a) J+ C	(0) 327 C
$()$ 1. $\sqrt{2}$ · 2	[עניני שיויכ]		(c) $427^{o}C$	(d) $527^{o}C$
(a) $1: \sqrt{2}: 2$ (b) 1	:2:4	19.	Pressure of a mixture of 4 g o	of O_2 and 2 g of H_2 confined in a
(c) $2:\sqrt{2}:1$ (d)	1:2:√2		hulb of 1 litre at $0^{\circ}C$ is	[41146 2000]
				[AIIIVIS 2000]
			(a) 23.213 dlm	(0) 31.205 $d(11)$

2.

3.

4.

5.

6.

7.

8.

9.

10.

UNIVERSAL SELF SCORER

SELF	246 Gaseous	state
	(c) 45.215 <i>atm</i>	(d) 15.210 <i>atm</i>
20.	If pressure becomes doub	le at the same absolute temperature on 2
	LCO_2 , then the volume of	of CO_2 becomes
		[A11MS 1992]
	(a) 2 <i>L</i>	(b) 4 <i>L</i>
	(c) 25 <i>L</i>	(d) 1 <i>L</i>
21.	Volume of the air that w	<i>i</i> ll be expelled from a vessel of 300 cm^3
	when it is heated from 2 be	$7^{o}C$ to $37^{o}C$ at the same pressure will
	(a) 310 cm ³	(b) 290 <i>cm</i> ³
	(c) 10 cm^3	(d) 37 cm^3
22.	300 <i>ml</i> of a gas at 2 pressure, the final volume	$7^{o}C$ is cooled to $-3^{o}C$ at constant is
		[NCERT 1981, MP PMT 1992]
	(a) 540 <i>ml</i>	(b) 135 <i>ml</i>
	(c) 270 <i>ml</i>	(d) 350 <i>ml</i>
	Assert	ion & Reason
		For AIIMS Aspirants

Read the assertion and reason carefully to mark the correct option out of the options given below :

- (a) If both assertion and reason are true and the reason is the correct explanation of the assertion. If both assertion and reason are true but reason is not the correct
- (b) explanation of the assertion.
- If assertion is true but reason is false. (c)
- If the assertion and reason both are false. (d)
- (e) If assertion is false but reason is true.

1.	Assertion :	Plot of P Vs. $1/V$ (volume) is a straight line.
	Reason :	Pressure is directly proportional to volume.
2.	Assertion :	Jet aeroplane flying at high altitude need pressurization of the cabin.
	Reason :	Oxygen is not present at higher altitude.
3.	Assertion :	1 mol of ${\cal H}_2$ and ${\cal O}_2$ each occupy 22.4 L of
		volume at $0^{o}C$ and 1 bar pressure.
	Reason :	Molar volume for all gases at the same temperautre and pressure has the same volume.
4.	Assertion :	Pressure exerted by a mixture of reacting gases is equal to the sum of their partial pressures.
	Reason :	Reacting gases react to form a new gas having pressure equal to the sum of both.
5.	Assertion :	Greater the value of Vander Waal's constant $'a'$ greater is the liquefaction of gas.
	Reason :	' <i>a</i> ' indirectly measures the magnitude of attractive forces between the molecules.
6.	Assertion :	Carbondioxide has greater value of root mean square velocity μ_{ms} than carbon monoxide.
	Reason :	$\mu_{\it ms}$ is directly proportional to molar mass.
7.	Assertion :	4.58 mm and $0.0098^{\circ}C$ is known to be triple
	Reason :	At this pressure and temperature all the three states i.e., water, ice and vapour exist simultaneously.

8.	Assertion	:	$1/4^{\circ}$ of the gas is expelled if air present in an							
			open vessel is heated from $27^{\circ}C$ to $127^{\circ}C$.							
	Reason	:	Rate of diffusion of a gas is inversely proportional to the square root of its molecular mass.							
9.	Assertion	:	Compressibility factor for hydrogen varies with pressure with positive slope at all pressures.							
	Reason	:	Even at low pressures, repulsive forces dominate hydrogen gas. [AIIMS 2005]							
10.	Assertion	:	vander Waal's equation is applicable only to non- ideal gases.							
	Reason	:	Ideal gases obey the equation $PV = nRT$.							
11.	Assertion	:	Pressure exerted by gas in a container with increasing temperature of the gas.							
	Reason	:	With the rise in temperature, the average speed of gas molecules increases.							
			[A11MS 1995]							
12.	Assertion	:	Gases do not settle to the bottom of container.							
	Reason	:	Gases have high kinetic energy.							
			[AIIMS 1997]							
13.	Assertion	:	A mixture of He and O_2 is used for respiration for deep sea divers.							
	Reason	:	He is soluble in blood. [AIIMS 1998]							
14.	Assertion	:	Wet air is heavier than dry air.							
	Reason	:	The density of dry air is more than density of water. [AllMS 1999]							
15.	Assertion	:	All molecules in a gas have some speed.							
	Reason	:	Gas contains molecules of different size and shape. [AIIMS 2001]							
16.	Assertion	:	Effusion rate of oxygen is smaller than nitrogen.							
	Reason	:	Molecular size of nitrogen is smaller than oxygen.[AIIMS							

Answers

Characteristics and Measurable properties of gases

1	С	2	с	3	b	4	а	5	с
6	d	7	а	8	b	9	С	10	а
11	a	12	b	13	a	14	d	15	С
16	е								

Ideal gas equation and Related gas laws

1	c	2	d	3	a	4	a	5	a
6	bc	7	a	8	a	9	C	10	d
11	а	12	а	13	а	14	а	15	С
16	C	17	b	18	C	19	a	20	b
21	C	22	C	23	d	24	a	25	с
26	с	27	b	28	с	29	с	30	с

21	h	22	а			
	~		ŭ			

31	а	32	а	33	b	34	b	35	b
36	c	37	d	38	c	39	c	40	a
41	b	42	c	43	c	44	d	45	a
46	a	47	b	48	d	49	b	50	c
51	d	52	а	53	a	54	c	55	a
56	а	57	b	58	а	59	b	60	a
61	b	62	d	63	c	64	b	65	d
66	b	67	d	68	а	69	а	70	b
71	c	72	а	73	d	74	c	75	a
76	c	77	d	78	b	79	b	80	b
81	d	82	d	83	c	84	c	85	a
86	d	87	а	88	а	89	С	90	c
91	а	92	С	93	а	94	b	95	а
96	а	97	d	98	b	99	е	100	а
101	с	102	с	103	с	104	с	105	а
106	а								

Kinetic molecular theory of gases and Molecular collisions

1	d	2	с	3	d	4	d	5	b
6	а	7	а	8	С	9	b	10	b
11	а	12	d	13	а	14	а	15	acd
16	а	17	b	18	d	19	b	20	d
21	а	22	d	23	С	24	b	25	b
26	а	27	d	28	а	29	d	30	С
31	C								

Molecular speeds

1	а	2	c	3	d	4	d	5	d
6	C	7	b	8	b	9	d	10	b
11	d	12	C	13	а	14	d	15	d
16	C	17	d	18	d	19	b	20	b
21	C	22	а	23	C	24	b	25	C
26	а	27	d						

Real gases and Vander waal's equation

1	b	2	b	3	с	4	C	5	d
6	С	7	d	8	b	9	d	10	b
11	с	12	b	13	b	14	d	15	C
16	b	17	а	18	b	19	C	20	а