	686 Hydrogen and	d Its	compounds
7.	Among the following, identify both oxidising and reducing ago		mpound which cannot act as [AMU 2002]
	(a) H_2O_2	(b)	H_2
	(c) SO_2	(d)	Cl_2
8.	Which of the following reaction	produ	ces hydrogen
	C	•	[AllMS 2002]
	(a) $Mg + H_2O$	(b)	$BaO_2 + HCl$
	(c) $H_2S_4O_8 + H_2O$	(d)	$Na_2O_2 + 2HCl$
9.	Hydrogen resembles in many of	f its pro	operties
			[MH CET 2001]
	(a) Halogen	(b)	Alkali metals
	(c) Both (a) and (b)	(d)	None of these
10.	Ortho and para hydrogen differ	in	[AFMC 2001]
	(a) Proton spin	(b)	Electron spin
	(c) Nuclear charge	(d)	Nuclear reaction
11.	Action of water or dilute miner	al acids	on metals can give
			[Kerala (Med.) 2002]
	(a) Monohydrogen	(b)	Tritium
	(c) Dihydrogen	(d)	Trihydrogen
12.	Hydrogen from HCl can be pre	pared b	
	(a) <i>Mg</i>	(b)	Си
	(c) <i>P</i>	(d)	Pt.
13.	Which of the following can adso		
			Finely divided nickel
	(c) Colloidal palladium		Colloidal platinum
14.	The nuclei of tritium (H^3) ato	m wou	lld contain neutrons
	(a) 1	(b)	2
	(c) 3	(d)	4
15.	The colour of hydrogen is		[MP PET 2004]
	(a) Black	(b)	Yellow
	(c) Orange	(d)	Colourless
16.	Ordinary hydrogen at room ten	•	
	(a) 75% of <i>o</i> -Hydrogen + 25%	,	, ,
	(b) 25% of <i>o</i> -Hydrogen + 75%	of <i>p</i> -1	lydrogen

Hydrogen cannot reduce

(a) Hot CuO

(a)

(c)

(c) Hot SnO_2

Antimony

Bismuth

(a) Dehydrogenation

Occlusion

(a) Mg

(c) Cu

solution is

(c) Mg

metal is known as

(a) As

17.

18.

19.

20.

21.

22.

The difference between heat of adsorption of ortho and para 28. 971 hydrogen is (b) 0.8 kJ mol (a) 0.4 kJ mol (d) None of these (c) Zero Hydrogen ion H^- is isoelectronic with 29. (a) *Li* (b) He (d) Li (c) H Hydrogen can be fused to form helium at [AFMC 2005] (a) High temperature and high pressure (b) High temperature and low pressure 004 (c) Low temperature and high pressure (d) Low temperature and low pressure Hydrogen can be prepared by mixing steam, and water gas at $500^{\circ}C$ 31. in the presence of Fe_3O_4 and Cr_2O_3 . This process is called (a) Nelson process (b) Serpeck's process (c) Bosch process (d) Parke's process (c) 50% of o-Hydrogen + 50% of p-Hydrogen Which of the following metal do not liberate hydrogen from dilute (d) 1% of o-Hydrogen + 99% of p-Hydrogen hydrochloric acid (a) Zn (b) Mg (c) Fe (d) Au (b) Fe_2O_3 An element reacts with hydrogen to form a compound A which on 33. (d) Hot Al_2O_3 treatment with water liberates hydrogen gas. The element can be (a) Nitrogen (b) Chlorine Hydrogen does not combine with (c) Selenium (d) Calcium (b) Sodium Hydrogen combines with other elements by 34. (d) Helium (a) Losing an electron The adsorption of hydrogen by metals is called

Gaining an electron

Sharing an electron

with alkali metals or halogen

metals or halogen

(a) Its molecule is diatomic

halogens

(d) Losing, gaining or sharing electron

Hydrogen can form compounds

Hydrogen atom does not contain any neutron

Which of the following terms is not correct for hydrogen

Which of the following explanation is best for not placing hydrogen

The ionization energy of hydrogen is high for group of alkali

Hydrogen is a much lighter element than the alkali metals or

(c)

35.

36.

Adsorbed hydrogen

Which is poorest reducing agent

(d) All have same reducing strength

Number of nucleons in D_2 molecule is

simple water. Its solubility is (a) Larger in heavy water

(c) Solubility is same in both

following property (a) Thermal conductivity

(c) Chemical properties

Reactive hydrogen

(a) Nascent hydrogen

Dihydrogen

of hydrogen is (a) 6

Atomic hydrogen

23.

24.

25.

26.

27.

[EAMCET 1999; Manipal PMT 1999]

(b) Hydrogenation

(d) Adsorption

(b) A1

(d) *Ca*

(b) Zn

(d) Fe

The metal which displaces hydrogen from a boiling caustic soda

Metals like platinum and palladium can absorb large volumes of

hydrogen under special conditions. Such adsorbed hydrogen by the

Which of the following produces hydrolith with dihydrogen

(b)

(c)

(c) 4

(c) 3

Occluded hydrogen

(b) Smaller in heavy water

(d) Smaller in simple water

(b) Magnetic properties

(d) Heat capacity

(d) Atomic hydrogen

The sum of protons, electrons and neutrons in the heaviest isotope

(b) 5

(d) 3

(d) 4

An ionic compound is dissolved simultaneously in heavy water and

Ortho-hydrogen and para-hydrogen resembles in which of the



Alkali metals only and H^- (b) It exists both as H^+ in different chemical (b) Halogen only compounds It is the only species which has no neutrons in the nucleus Both alkali metals and halogens (c) (c) Heavy water is unstable because hydrogen is substituted by its Neither alkali metals nor halogens (d) isotope deuterium Hydrogen molecule differs from chlorine molecule in the following 50. When electric current is passed through an ionic hydride in the respect molten state (a) Hydrogen molecule is non-polar but chlorine molecule is polar (a) Hydrogen is liberated at the anode Hydrogen molecule is polar while chlorine molecule is non-(b) Hydrogen is liberated at the cathode (c) No reaction takes place Hydrogen molecule can form intermolecular hydrogen (d) Hydride ion migrates towards cathode bonds but chlorine molecule does not Which of the halogen has maximum affinity for hydrogen Hydrogen molecule cannot participate in coordination bond formation but chlorine molecule can (a) F_2 Which of the following statements concerning protium, deuterium 51. (c) Br_2 (d) I_2 and tritium is not true Which of the following statements is most applicable to hydrogen (a) They are isotopes of each other (a) It can act as a reducing agent (b) They have similar electronic configurations (b) It can act as an oxidising agent They exist in the nature in the ratio of 1:2:3(c) It can act both as oxidising and reducing agent (d) Their mass numbers are in the ratio of 1:2:3 (d) It can neither act as oxidising nor as a reducing agent When SO_3 is treated with heavy water the product is/are 52. Hydrogen is (a) Deuterium and sulphuric acid (a) Electropositive Deuterium and sulphurous acid (b) (b) Electronegative (c) Only deuterium Both electropositive as well as electronegative Dideuterosulphuric acid Neither electropositive nor electronegative Hydrogen has three isotopes, the number of possible diatomic 53. Ionization energy of hydrogen is molecules will be (a) Equal to that of chlorine (a) 2 (b) 6 (b) Lesser than that of chlorine (c) 9 (d) 12 Slightly higher than that of chlorine In which of the compounds does hydrogen have an oxidation state of 54. (d) Much higher than that of chlorine -1 Hydrogen acts as a reducing agent and thus resembles (a) CH_4 (b) NH_3 (a) Halogen (b) Noble gas (c) HCl (d) CaH_2 (c) Radioactive elements (d) Alkali metals Which position for hydrogen explain all its properties Pure hydrogen is obtained by carrying electrolysis of (a) At the top of halogen Water containing H_2SO_4 (b) At the top of alkali metals (b) Water containing NaOH (c) At the top of carbon family $Ba(OH)_2$ solution (c) (d) None of these KOH solution Hydrogen readily combines with non-metals and thus it shows its In Bosch's process which gas is utilised for the production of 56. (a) Electronegativity character hydrogen gas (b) Electropositive character (a) Producer gas (b) Water gas (c) Both (a) and (b) (c) Coal gas (d) None of these (d) None of these 57. Deuterium differs from hydrogen in The oxidation states shown by hydrogen are Chemical properties (a) -1 only (b) Zero only Physical properties (c) +1, -1, 0(d) +1 only Both physical and chemical properties Hydrogen readily combines with metals and thus shows its Radioactive properties (a) Electropositive character (b) Electronegative character 58. Tritium undergoes radioactive decay giving (c) Both (a) and (b) (d) None of these (a) α -particles (b) β -particles Electrolysis of fused sodium hydride liberate hydrogen at the (c) Neutrons (d) γ-rays (a) Anode The gas used in the hydrogenation of vegetable oils in the presence 59. (b) Cathode of nickel as catalyst is (c) Cathode and anode both (a) Methane (b) Ethane (d) None of these (c) Ozone (d) Hydrogen Protonic acid is The conversion of atomic hydrogen into ordinary hydrogen is 60. (a) A compound that form solvated hydrogen ion in polar solvent

37.

38.

39.

40.

41.

42.

43.

45.

47.

48.

49.

(b) An acid which accepts the proton

An acid which donates the proton

In all its properties, hydrogen resembles

A compound that forms hydride ion in polar solvent

- (a) Exothermic change
- - - Endothermic change (b)
 - (c) Nuclear change
 - Photochemical change
- 61. The name hydrogen was given by

	688 Hydrogen and Its compounds		
	(a) Cavendish (b) Lavoisier		(c) Anhydrous calcium chloride
	(c) Urey (d) None of these		(d) Na_3PO_4
62.	The ratio $C_p/C_ u$ for H_2 is	75.	Which is the lightest gas [CPMT 1993]
	(a) 1.40 (b) 1.67		(a) Nitrogen (b) Helium
	(c) 1.33 (d) None of these		(c) Oxygen (d) Hydrogen
63.	Triatomic hydrogen is called	76.	The composition of tritium is [UGET Manipal 1995]
	(a) Deuterium (b) Hyzone (c) Ortho form (d) Hydronium ion		(a) 1 electron, 1 proton, 1 neutron (b) 1 electron, 2 protons, 1 neutron
			(c) 1 electron, 1 proton, 2 neutrons
64.	$LiAlH_4$ is obtained by reacting an excess of With an ethereal solution of $AlCl_3$		(d) 1 electron, 1 proton, 3 neutrons
	(a) LiCl (b) LiH	77.	The property of hydrogen which distinguishes it from alkali metals is
	(c) Li (d) LiOH		(a) Its electropositive character
65.	Alkali metal hydrides react with water to give		(b) Its affinity for non metal
	(a) Acidic solution (b) Basic solution		(c) Its reducing character
	(c) Neutral solution (d) Hydride ion		(d) Its non-metallic character
66.	Ionic hydrides are usually	78.	The hydride ion H^{-} is a stronger base than its hydroxide ion
	(a) Good electrically conductors when solid		OH^- . Which of the following reactions will occur if sodium
	(b) Easily reduced (c) Good reducing agents		hydride (NaH) is dissolved in water
	(d) Liquid at room temperature		[CBSE PMT 1997]
67.	When $NaBH_4$ is dissolved in water		(a) $H^-(aq) + H_2O \rightarrow H_3O^-(aq)$
	(a) It decomposes with the evolution of H_2		(b) $H^{-}(aq) + H_{2}O(l) \rightarrow OH^{-}(aq) + H_{2}(g)$
	(b) Na^+ and BH_4^- are formed which are stable		(c) $H^{-}(aq) + H_2O(l) \rightarrow \text{No reaction}$
	(c) BH_4^- ions formed initially decompose to produce OH^- ions,		(d) None of these
	which prevent further decomposition	79.	Hydrogen accepts an electron to form inert gas configuration. In this
	(d) NaH and B_2H_6 are produced		it resembles [Pb. PMT 1997] (a) Halogen (b) Alkali metals
68.	Systematic name of H_2O (oxide of hydrogen) is		(c) Chalcogens (d) Alkaline earth metals
00.		80.	Which of the following is correct for hydrogen
	(a) Water (b) Hydrogen oxide (c) Oxidane (d) None of these		[AFMC 1997; BHU 1997]
69.	Group 2 hydrides with significant covalent character is/are		(a) It can form bonds in +1 as well as -1 oxidation state
-	(a) BeH_2 (b) MgH_2		(b) It is always collected at cathode
	(c) Both (a) and (b) (d) None of these		(c) It has a very high ionization potential
70.	Limiting compositions of <i>F</i> -block hydrides are		(d) It has same electronegativity as halogens
,	(a) MH_2 and MH_3 (b) MH_3 and MH_5	81.	Which of the following will not displace hydrogen
			[Pb. PMT 1999] (a) Ba (b) Pb
_			(c) Hg (d) Sn
71.	Hydrogen directly combines with [Roorkee Entrance 1990] (a) Au (b) Cu	82.	Which of the following gas is insoluble in water
	(a) Au (b) Cu (c) Ni (d) Ca		[Pb. CET 2003]
72.	Chemical <i>A</i> is used for water softening to remove temporary		(a) SO_2 (b) NH_3
	hardness. A reacts with sodium carbonate to generate caustic soda.		(c) H_2 (d) CO_2
	When CO_2 is bubbled through a solution of A , it turns cloudy.	83.	Which element forms maximum compound in chemistry
	What is the chemical formula of <i>A</i>	ს ე.	(D). CTT pood

[Pb. CET 1990; AIIMS 1999]

(a) O

(c) Si

(a) Cold water

 $(c)\quad \text{Conc. sulphuric acid}$

(b) *H*

(d) C

(b) Hot NaOH solution

(d) dilute HCl

Hydrogen is not obtained when zinc reacts with

Water or hydride of oxygen

(a) $CaCO_3$

(b) CaO

(c) $Ca(OH)_2$

(d) $Ca(HCO_3)_2$

73. When same amount of zinc is treated separately with excess of sulphuric acid and excess of sodium hydroxide solution the ratio of volumes of hydrogen evolved is [CPMT 1991]

(a) 1:1

(b) 1:2

(c) 2:1

(d) 9:4

74. Which one of the following substances is used in the laboratory for a fast drying of neutral gases

[CBSE PMT 1992]

- (a) Phosphorus pentoxide
- (b) Active charcoal

Which of the following will cause softening of hard water

(a) Passing it through cation exchange resin

(a) Heavy water (D_2O)



(b) Ordinary water (H_2O)

	because [AMU 2002]		
	(a) They are highly soluble in water	12.	(c) Zinc rod (d) Fused caustic soda Temporary hardness of water is due to the presence of
	(b) Their Ca^{++} and Mg^{++} salts are water soluble	12.	(a) Magnesium bicarbonate (b) Calcium chloride
			(c) Magnesium sulphate (d) Calcium carbonate
	(c) Their Ca^{++} and Mg^{++} salts are insoluble in water	13.	Which of the following is not true
	(d) None of these		(a) Hardness of water depends on its behaviour towards soap
2.	D_2O is used more in [BHU 1997; CPMT 1997]		(b) The temporary hardness is due to the presence of <i>Ca</i> and <i>Mg</i> bicarbonates
	(a) Chemical industry (b) Nuclear reactor		(c) Permanent hardness is due to the presence of soluble <i>Ca</i> and <i>Mg</i> sulphates, chlorides and nitrates
	(c) Pharmaceutical preparations		(d) Permanent hardness can be removed by boiling the water
	(d) Insecticide preparation	14.	The molarity of pure water at $4^{\circ}C$ is
3.	Heavy water (D_2O) is [RPET/PMT 2000; CPMT 2000]		(a) 1 M (b) 2.5 M
	(a) A product of oxygen and hydrogen		(c) 5 M (d) 55.5 M
	(b) Water of mineral springs	15.	Which of the following is not a hard water
	(c) Water obtained by repeated distillation and condensation		(a) Water containing $CaCl_2$
	(d) Ordinary water containing dissolved salts heavy metals		(b) Water containing dil. HCl
4.	Temporary hardness may be removed from water by adding		(c) Water containing $MgSO_4$
	[Pb. PMT 2002]		(d) None of these
	(a) $CaCO_3$ (b) $Ca(OH)_2$	16.	Heavy water is used in atomic reactor as
	(c) CaSO ₄ (d) HCl		(a) Coolant
5.	Heavy water is [AFMC 1997; UPSEAT 2003		(b) Moderator
	MH CET 2003; Pb. CET 2001]		(c) Both moderator and coolant
	(a) Water containing Fe, Cr, Mn		(d) Neither coolant nor moderator
	(b) Water at 0°C	17.	Heavy water freezes at
	(c) D_2O		(a) $0^{\circ}C$ (b) $3.8^{\circ}C$
	(d) Water obtained after a number of distillations		(c) $38^{\circ}C$ (d) $-0.38^{\circ}C$
6.	Heavy water is compound of [DPMT 2001; DCE 2002]	18.	The pH of D_2O and H_2O at 298 K is
	(a) Oxygen and heavier isotopes of hydrogen		(a) 7.0, 7.0 (b) 7.35, 7.0
	(b) Hydrogen and heavier isotopes of oxygen		(c) 7.0, 6.85 (d) 6.85, 7.35
	(c) Heavier isotopes of oxygen and hydrogen	19.	Which of the following is not true
	(d) None of these		(a) Ordinary water is electrolysed more rapidly than D_2O
7.	Which of the following pair of ions makes the water hard [AMU 2002]		(b) Reaction between H_2 and Cl_2 is much faster than D_2 and
	(a) Na^+, SO_4^{2-} (b) K^+, HCO_3^-		Cl_2
	(c) Ca^{2+} , NO_3^- (d) NH_4^+ , Cl^-		(c) D_2O freezes at lower temperature than H_2O
_			(d) Bond dissociation energy for D_2 is greater than H_2
8.	Temporary hardness of water can be removed by [Pb. PMT 2001]	20.	Which of the following will determine whether the given colourless
	(a) Addition of potassium permagenate		liquid is water or not
	(b) Boiling		(a) Melting
	(c) Filtration		(b) Tasting
	(d) Addition of chlorine		(c) Phosphthalein
9.	When zeolite (Hydrated sodium aluminium silicate) is treated with		(d) Adding a pinch of anhydrous $CuSO_4$
	hard water the sodium ions are exchanged with	21.	Lead pipes are not used for carrying drinking water because
	[DPMT 2000]		(a) They are covered with a coating of lead carbonate
	(a) OH^- ions (b) SO_4^{2-} ions		(b) They are corroded by air and moisture
	(c) Ca^{2+} ions (d) H^{+} ions		(c) Water containing dissolved air attacks lead forming soluble hydroxide
10.	Which of the following statements do not define the characteristic		(d) None of these
	property of water "Water is a universal solvent"	22.	Which one of the following removes temporary hardness of water
	(a) It can dissolve maximum number of compounds (b) It has very low dielectric constant		(a) Slaked lime (b) Plaster of Paris
	(c) It has high liquid range		(c) Cuprous (d) Hydrolith
			Mari Cal Cil 1 1 1 C 1 C 1 1 .

23.

Synthetic detergents are more effective in hard water than soaps

(d) None of these

11.

The velocity of neutrons in nuclear reactor is slowed down by



(d) Water molecule is not linear

given below the lists

34.

Match list 1 with list 11 and select the correct answer using the codes

/4	\ D	. 1 .						
(b					List 1		List	
(c (d				1.	Heavy water) Bicarbonates a in water	s of <i>Mg</i> and
. w		rocess permanent hardness of water can be		2.	Temporary hard water	er (b) No foreign io	ons in water
	emoved, by adding	[AFMC 2005]		3.	Soft water	(c) D ₂ O	
(a) Sodalime	(b) Sodiumbicarbonate		4.	Permanent hard water	er (d	l) Sulphates an	d chlorides of
(c	e) Washing soda	(d) Sodium chloride				\mathcal{N}	<i>lg</i> and <i>Ca</i> in w	ater
Pe	ermutit is technical nan	ne given to		Cod				
(a) Aluminates of calciu	ım and sodium			1- <i>c</i> , 2- <i>d</i> , 3- <i>b</i> , 4- <i>a</i>		1- <i>b</i> , 2- <i>a</i> , 3- <i>c</i> ,	
(b) Silicates of calcium	and sodium			1- <i>b</i> , 2- <i>d</i> , 3- <i>c</i> , 4- <i>a</i>		1- <i>c</i> , 2- <i>a</i> , 3- <i>b</i> ,	4- <i>d</i>
(c	e) Hydrated silicates of	f aluminium and sodium	35.	The	H-O-H angle in	water mol	ecule is about	[ATMO accel
(d				(a)	90°	(b)	180°	[AFMC 2001]
ΤÌ		tritium oxide molecule is		(c)	102°	(d)		
) 18 <i>amu</i>	(b) 20 <i>amu</i>	36.	. ,	en two ice cubes are pr	. ,	-	ney unite to form
	e) 22 <i>amu</i>	(d) 24 <i>amu</i>			cube. Which of the fol			•
	lolecular weight of heav	* /		toge				[AFMC 2001]
					Hydrogen bond forma	tion		
	1) 19	· /			Van der Waals forces Covalent attraction			
•	e) 17	(d) 20		(c) (d)	lonic interaction			
W	•	anently hard when it contains	37.	. ,	nt is formed when calcin	um carbid	e reacts with h	eavv water[Manipal P
(a	, .		•	(a)	C_2D_2	(b)	~ ~	, , , , , , , , , , , , , , , , , , , ,
(b	,					` '	-	
(c	e) Sulphates of <i>Cu</i> and	l Hg		(c)	Ca_2D_2O		CD_2	
(d	 Carbonates and bica 	arbonates of <i>Mg</i> and <i>Ca</i>	38.	Pure	e water can be obtained	l from sea	water by	[CDCC DATE con]
	odium sulphate is solub ecause	le in water but barium sulphate is insoluble [Pb. PMT 1995]		(a)	Centrifugation	. ,	Plasmolysis	[CBSE PMT 2001]
(a	The hydration energy	rgy of Na_2SO_4 is more than its lattice	39.	(c) Acti	Reverse osmosis on of water or dilute m	` '	Sedimentations on metals ca	
/L	0,5	of Dago is some than its bulleting		(a)	Monohydrogen	(b)	Tritium	[Refula 17711 2002]
(b		of $BaSO_4$ is more than its hydration		(c)	Dihydrogen	()	Trihydrogen	
,	energy	1 . 1 . 1.146		(e)	D_2			
(c		as no role to play in solubility	40		_	. 9.1		1 11 51
(d	l) The hydration ene	rgy of Na_2SO_4 is less than its lattice	40.		al which does not reac	et with co	ld water but e	
	energy			stea		(L)	К	[DCE 2002]
(e	e) Both (a) and (b)			(a) (c)	Pt	()	Fe	
TÌ	he alum used for purify	ing water is [EAMCET 1999]	41.	` '	of neutral water at rooi			
(a) Ferric alum	(b) Chrome alum	4	(a)		•	•	
(c	e) Potash alum	(d) Ammonium alum					10^{-7}	
w	hich of the following m	netal will not reduce H_2O		(c)				
	g		42.	Max	imum number of hydro			
(-) C-	[CPMT 1999]		()	-			Г 2004; ВНU 2004]
(a		(b) Fe		(a)		. ,	2	
(c	•	(d) Li	4.0	(c)	3	` '	4	
W	hich of the following is	correct about heavy water	43.	The	low density of ice com	pared to v	ater is due to	[DL CET noo4]
		[DCE 2002]		(2)	Induced dipole-induce	d dinala ir	ntaractions.	[Pb. CET 2004]
(a	•	ng maximum density is known as heavy		(a) (b)	•	-		
	water			(b)	Dipole-induced dipole		11	
(b) It is heavier than wa	ater (H_2O)		(c)	Hydrogen bonding int Dipole-dipole interacti			
(c	t) It is formed by the and oxygen	combination of heavier isotope of hydrogen	44.	(d) Whi	ch of the following acid		I when SiF_4 1	eacts with water
(d								[BHU 2004]
,	•	er is exceptionally high because		(a)	SiF_4	(b)	H_2SiF_4	
	0,	[KCET 2001]		(2)	H_2SO_4	(4)		
(a) There is covalent bo	and between H and O		(c)		(d)	1125116	
(b	o) Water molecule is li	near	45.	Trip	le point of water is			[AFMC 2004]
(c) Water molecules ass	sociate due to hydrogen bonding		(a)	273 <i>K</i>	(b)	373 <i>K</i>	

(c) 203*K*

46.

[SCRA 2001]

[BHU 2005]

(d) 193*K*

Hardness of water is due to presence of salts of



- (a) Na and K
- (b) Ca and Mg
- (c) Ca and K
- (d) Ca and Na

Hydrogen peroxide

- In which of the following reaction hydrogen peroxide is a reducing agent [BHU 1995]
 - (a) $2FeCl_2 + 2HCl + H_2O_2 \rightarrow 2FeCl_3 + 2H_2O$
 - (b) $Cl_2 + H_2O_2 \rightarrow 2HCl + O_2$
 - (c) $2HI + H_2O_2 \rightarrow 2H_2O + I_2$
 - (d) $H_2SO_3 + H_2O_2 \to H_2SO_4 + H_2O_3$
- There is a sample of 10 volume of hydrogen peroxide solution. Calculate its strength [UPSEAT 2001]
 - (a) 3.00%
- (b) 4.045%
- (c) 2.509%
- (d) 3.035%
- 3. In lab H_2O_2 is prepared by

[CPMT 2002; MH CET 2003;

- Pb. PMT 2004; BCECE 2005]
- (a) Cold $H_2SO_4 + BaO_2$ (c) Conc. $H_2SO_4 + Na_2O_2$
- (b) $HCl + BaO_2$
- (d) $H_2 + O_2$
- **4.** The structure of H_2O_2 is

[UPSEAT 2001]

(a)
$$H O - OH$$

(b)
$$H O - O H$$

H

- (c) H-O-O-H
- (d) $O O \searrow$
- 5. HCl is added to the following oxides which one would give H_2O_2 [Kurukshetra CEE 1998] 8
 - (a) MnO_2
- (b) PbO_2
- (c) BaO_2
- (d) None of these
- **6.** The oxide that gives hydrogen peroxide (H_2O_2) on the treatment with a dilute acid (H_2SO_4) is **[Pb. PMT 1999]**
 - (a) MnO_2
- (b) PbO_2
- (c) Na_2O_2
- (d) TiO_2
- 7. Hydrogen peroxide is reduced by

[CPMT 2000; CBSE PMT 2000; KCET 2002]

- (a) Ozone
 - (b) Barium peroxide
 - (c) Acidic solution of KMnO4
 - (d) Lead sulphide suspension
- **8.** The reaction of $H_2S + H_2O_2 \rightarrow S + 2H_2O$ manifests

[UPSEAT 2000]

- (a) Acidic nature of H_2O_2
- (b) Alkaline nature of H_2O_2
- (c) Oxidising nature of H_2O_2
- (d) Reducing action of H_2O_2
- **9.** What is the product of the reaction of H_2O_2 with Cl_2
 - (a) $O_2 + HOCl$
- (b) $HCl + O_2$
- (c) $H_2O + HCl$
- (d) $HCl + H_2$
- 10. H_2O_2 will oxidise

[Roorkee 1995]

- (a) $KMnO_{\Lambda}$
- (b) PbS
- (c) MnO_2

11.

12.

13.

- (d) H_2S
- Fenton's reagent is

[MP PET 2000; RPET 2000]

- (a) $FeSO_4 + H_2O_2$
- (b) Zn + HCl
- (c) Sn + HCl
- (d) None of these
- The structure of H_2O_2 is (a) Planar
- (b) Linear
- (c) Spherical
- (d) Non-planar
- The volume strength of 1.5 N H_2O_2 solution is

[BHU 2004; Pb. CET 2004]

[CBSE 1999; AFMC 2004]

- (a) 8.4 *litres*
- (b) 4.2 *litres*
- (c) 16.8 litres
- (d) 5.2 litres
- The volume of oxygen liberated from 15 $\it ml$ of 20 volume $\it H_2O_2$ is[MH CET 20]
 - (a) 250 *ml*
- (b) 300 ml
- (c) 150 ml
- (d) 200 ml
- The strength in volumes of a solution containing 30.36 *g/litre* of H_2O_2 is [UPSEAT 2004]
 - (a) 10 volume
- (b) 20 volume
- (c) 5 volume
- (d) None of these
- 16. Hydrogen peroxide is used as
 - (a) Oxidising agent
 - (b) Reducing agent
 - (c) Both as oxidising and reducing agent
 - (d) Drying agent
- 17. Equivalent weight of H_2O_2 is
 - (a) 17 FE 10081 -
- (b) 34 (d) 18
- 20 volume H_2O_2 solution has a strength of about
- (a) 30%
- (b) 6%
- (c) 3%

19.

- (d) 10%
- H_2O_2 is manufactured these days
 - ays [DCE 2004]
 - (a) By the action of H_2O_2 on BaO_2
 - (b) By the action of H_2SO_4 on Na_2O_2
 - (c) By electrolysis of 50% H_2SO_4
 - (d) By burning hydrogen in excess of oxygen
- **20.** Which one of the following is a true peroxide
 - (a) NO_2
- (b) MnO_2
- (c) BaO_2
- (d) SO_2
- 21. 1 \emph{ml} of H_2O_2 solution gives 10 \emph{ml} of O_2 at NTP. It is
 - (a) 10 vol. H_2O_2
- (b) 20 vol. H_2O_2
- (c) 30 vol. H_2O_2
- (d) 40 vol. H_2O_2
- **22.** Which substance does not speed up decomposition of H_2O_2
 - (a) Glycerol
- (b) *Pt*
- (c) Gold
- (d) MnO_2
- **23.** Which of the following cannot be oxidised by H_2O_2
 - (a) O_3
- (b) KI / HCl
- (c) PbS
- (d) Na_2SO_3
- **4.** Which substance cannot be reduced by H_2O_2

- $KMnO_4 / H_2SO_4$
- $K_2Cr_2O_7 / H_2SO_4$ (b)
- (c) Ag_2O
- Which of the following statements is incorrect 25.
 - (a) H_2O_2 can act as an oxidising agent
 - (b) H_2O_2 can act as a reducing agent
 - (c) H_2O_2 has acidic properties
 - (d) H_2O_2 has basic properties
- H_2O_2 is 26.
 - (a) Poor polar solvent than water
 - (b) Better polar solvent than H_2O
 - (c) Both have equal polarity
 - Better polar solvent but its strong auto oxidising ability limits
- H_2O_2 used in rockets has the concentration 27.
 - (a) 50%
- (b) 70%
- (c) 30%
- (d) 90%
- 28 H_2O_2 is a
 - (a) Weak acid
- (b) Weak base
- (c) Neutral
- (d) None of these
- Nitrates of all metals are 29.
 - (a) Soluble in water
- (b) Insoluble
- (c) Coloured
- (d) Unstable
- Decomposition of H_2O_2 is prevented by 30.
 - (a) NaOH
- (b) MnO_2
- (c) Acetanilide
- (d) Oxalic acid
- H_2O_2 is always stored in black bottles because 31.
 - (a) It is highly unstable
 - (b) Its enthalpy of decomposition is high
 - (c) It undergo autooxidation on prolonged standing
 - (d) None of these
- H_2O_2 on reacting with ethene gives 32.
 - (a) Ethane
- (b) Ethanal
- (c) Ethylene glycol
- (d) Ethanol
- Which of the following is wrong about H_2O_2 ? It is used 33.
 - As aerating agent in production of spong rubber
 - (b) As an antichlor
 - (c) For restoring white colour of blackened lead painting
 - (d) None of these
- $H_2O_2 \rightarrow 2H^+ + O_2 + 2e^-$; $E^{\circ} = -0.68 \text{ V}$. This equation 34. represents which of the following behaviour of H_2O_2
 - (a) Reducing
- (b) Oxidising
- (c) Acidic
- (d) Catalytic
- The structure of H_2O_2 is 35
 - (a) Open book like
- (b) Linear
- (c) Closed book
- (d) Pyramidal
- 36. On shaking H_2O_2 with acidified potassium dichromate and ether, ethereal layer becomes
 - (a) Green
- (b) Red
- Blue
- (d) Black
- K_a of H_2O_2 is of the order of 37.
- [MP PMT 1994]


- 10^{-12} (a)
- (b) 10^{-14}
- 10^{-16} (c)
- (d) 10^{-10}
- 38. In which of the following reactions, $H_2 O_2$ acts as a reducing agent[EAMCET 2
 - $PbO_2(s) + H_2O_2(aq) \rightarrow PbO(s) + H_2O(l) + O_2(g)$
 - $Na_2SO_3(aq) + H_2O_2(aq) \rightarrow Na_2SO_4(aq) + H_2O(l)$
 - $2Kl(aq) + H_2O_2(aq) \rightarrow 2KOH(aq) + I_2(s)$
 - $KNO_2(aq) + H_2O_2(aq) \rightarrow KNO_3(aq) + H_2O(l)$
- H_2O_2 acts as an oxidising agent in 39.

[Kerala PMT 2004]

- Neutral medium
- (b) Acidic medium
- Alkaline medium (c)
- Alkaline and neutral medium
- Acidic and alkaline medium
- The H-O-O bond angle in H_2O_2 is 40. [Kerala PMT 2004]
 - 107.28^{o} (a)
- 109.28°
- (c) 104.5^{o}
- (d) 106^{o}
- 97^{o} (e)
- The volume of oxygen liberated from 0.68gm of H_2O_2 is 41.

[Pb. PMT 2004]

- 112ml(a)
- $224 \, ml$
- (c) 56 ml
- (d) 336 ml



Objective Questions

- Polyphosphates are used as water softening agents because they
 - (a) Form soluble complexes with anionic species
 - Precipitate anionic species
 - Forms soluble complexes with cationic species
 - (d) Precipitate cationic species
- The critical temperature of water is higher than that of O_2 because

 H_2O molecule has

[IIT 1997]

- (a) Fewer electrons than oxygen
- (b) Two covalent bonds
- (c) V-shape
- (d) Dipole moment
- One mole of calcium phosphide on reaction with excess water gives[IIT 1999] 3.
 - (a) One mole of phosphene
 - (b) Two moles of phosphoric acid
 - Two moles of phosphene (c)
 - (d) One mole of phosphorus pentaoxide
- When zeolite, which is hydrated sodium aluminium silicate, is 4. treated with hard water the sodium ions are exchanged with
 - (a) H^+ ions
- (b) Ca^{2+} ions
- (c) Mg^{2+} ions
- (d) Both Ca^{2+} and Mg^{2+}
- Hydrogen peroxide is
 - (a) A stronger acid than water
 - (b) A weaker acid than water
 - (c) An oxidising agent
 - (d) A reducing agent



5.	Hydrogen can be obtained from water by		(c) Para (d) All have same energy
	(a) Reaction with metal oxides	19.	Saline hydrides react explosively with water, such fires can b
	(b) Reaction with non-metal oxides		extinguished by
	(c) Reaction with metals		(a) Water (b) Carbon dioxide
	(d) Reaction with metal hydrides		(c) Sand (d) None of these
7.	Which of the following is/are hard water(s)	20.	Metals of groups 7, 8 and 9 do not form metallic hydrides. This i termed as
	(a) Water containing some potash alum		(a) Hydride gap (b) Hydride shift
	(b) Water containing a few drops of HCl(c) Water containing common salt		(c) Anhydride (d) Dehydride
	(c) Water containing common salt (d) Water containing calcium nitrate	21	()
8.	Pick the odd one out	21.	When temporary hard water containing $Mg(HCO_3)_2$ is boiled th ppt. formed is of
	(a) Sodium borohydride reacts very slowly with cold water		•••
	(b) Sodium borohydride reacts very violently with cold water to		
	produce H_2		(c) $Mg(OH)_2$ (d) None of these
	(c) Solubility of sodium borohydride in water at $25^{\circ}C$ is 10.05 g/mL	22.	Permanent hardness due to Mg^{2+} ions is best removed by
	(d) Melting point of sodium borohydride is 500°C		(a) $Ca(OH)_2$ (b) Na_2CO_3
9.	Hydrogen can be obtained from water, by the action of water on		- · · · · · · · · · · · · · · · · · · ·
	(a) Calcium carbide (b) Calcium hydride		(c) $Na_2CO_3 + Ca(OH)_2$ (d) None of these
	(c) Calcium oxide (d) Calcium	23.	The most abundant element in the universe is
0.	What is true about ice		(a) Carbon (b) Silicon (c) Hydrogen (d) Helium
	(a) Its density is more than water	24.	(c) Hydrogen (d) Helium Pick out the correct statement
	(b) It is a good conductor of heat		(a) By decreasing the temperature pure para-hydrogen can b
	(c) It is a thermal insulator		obtained
	(d) Its density is less than water		(b) By increasing the temperature pure ortho-hydrogen can b
1.	Hydrogen will not reduce [IIT 1985]		obtained
	(a) Heated cupric oxide (b) Heated ferric oxide		(c) By decreasing the temperature pure ortho-hydrogen can b obtained
	(c) Heated stannic oxide (d) Heated aluminium oxide		(d) By increasing the temperature pure para-hydrogen can b
2.	$\it HCl$ is added to following oxides. Which one would give H_2O_2 [11T 19	801	obtained
		25.	Hydrogen can be produced by heating
	(a) MnO_2 (b) PbO_2		(a) Cu with H_2SO_4 (b) Sodium formate
	(c) BaO (d) None of these		(c) Sodium oxalate (d) None of these
3.	Which of the following pair will not produce dihydrogen gas	26.	Plumbosolvency is a health hazard in the transportation of
	[IIT 1994]		(a) Hard water only
	(a) $Cu + HCl(dil.)$ (b) $Fe + H_2SO_4$		(b) Soft water only (c) Both (a) and (b)
	(c) $Mg + \text{steam}$ (d) $Na + \text{alcohol}$		(d) Water containing plum juice
	()	27.	A sample of water contains sodium chloride. It is
4.	The amount of H_2O_2 present in 1 $\it L$ of 1.5 $\it NH_2O_2$ solution is	Į	(a) Hard water (b) Soft water
	(a) $2.5 g$ (b) $25.5 g$		(c) Moderately hard (d) None of these
	(c) 3.0 g (d) 8.0 g	28.	Hardness producing salt, whose solubility in water decreases with
5.	Hydrogen is evolved by the action of cold dil. HNO_3 on		rise of temperature is
	[IIT 1998]		(a) $CaCl_2$ (b) $CaSO_4$
	(a) Fe (b) Mn		(c) $Ca(HCO_3)_2$ (d) $MgSO_4$
	(c) Cu (d) Al	29.	A sample of water containing some dissolved table sugar and
6.	Hydrogen can behave as a metal		common salt is passed through organic ion exchange resins. The resulting water will be
	(a) At very high temperature (b) At very low temperature		(a) Tasteless (b) Sweet
	(c) At very high pressure (d) At very low pressure		(c) Salty (d) None of these
7.	$D_2{\cal O}$ is preferred to $H_2{\cal O}$, as a moderator, in nuclear reactors	30.	Water obtained by purification with organic ion exchange resins is
-	because		(a) Pure water
	(a) D_2O slows down fast neutrons better		(b) Free from only Ca^{2+} , Mg^{2+} ions
	· · · · · · · · · · · · · · · · · · ·		
	(b) D_2O has high specific heat		
	(c) D_2O is cheaper	31.	(d) None of theseWhich of the following can effectively remove all types of hardnes
	(d) None of these	ال	of water
8.	Out of the two allotropic forms of dihydrogen, the form with lesser		(a) Soap (b) Washing soda
	molecular energy is		(c) Slaked lime (d) None of these
	(a) Ortho (b) Meta		



- 32. A commercial sample of hydrogen peroxide is labelled as 10 volume. Its percentage strength is nearly [KCET 2005]
 - (a) 1%
- (b) 3%
- (c) 10%
- (d) 90%



Assertion & Reason

For AIIMS Aspirants

Read the assertion and reason carefully to mark the correct option out of the options given below :

- (a) If both assertion and reason are true and the reason is the correct explanation of the assertion.
- (b) If both assertion and reason are true but reason is not the correct explanation of the assertion.
- (c) If assertion is true but reason is false.
- (d) If the assertion and reason both are false.
- (e) If assertion is false but reason is true.
- Assertion : Hydrogen combines with other elements by losing, gaining or sharing of electrons.
 - Reason : Hydrogen forms electrovalent and covalent bonds

with other elements.

2. Assertion : Calgon is used for removing Ca^{2+} and Mg^{2+}

ions from hard water.

Reason : Calgon forms precipitates with Ca^{2+} and

 Mg^{2+} .

3. Assertion : Decomposition of $H_2{\cal O}_2$ is a disproportionation

reaction.

Reason : H_2O_2 molecule simultaneously undergoes

oxidation and reduction.

- **4.** Assertion : H_2O_2 has higher boiling point than water.
 - Reason : H_2O_2 has stronger dipole-dipole interactions

than water.

- **5.** Assertion : H_2O_2 is not stored in glass bottles.
 - Reason : Alkali oxides present in glass catalyse the

decomposition of H_2O_2 .

6. Assertion : H_2O_2 reduces Cl_2 to HCl.

Reason : H_2O_2 is called antichlor.

7. Assertion : In acidic medium, H_2O_2 reacts with MnO_2 to

give O_2 .

Reason : H_2O_2 is a strong oxidising agent.

8. Assertion : In alkaline solution, H_2O_2 reacts with

potassium ferricyanide.

Reason : H_2O_2 is a strong reducing agent.

9. Assertion : Acidulated water is an example of hard water.

Reason : In the presence of an acid, soap is converted into

insoluble free fatty acids.

10. Assertion : Hydrogen peroxide forms only one series of salts

called peroxides.

Reason : Hydrogen peroxide molecule has only one

replaceable hydrogen atom.

	Hydrogen												
1	а	2	d	3	а	4	а	5	d				
6	d	7	b	8	а	9	С	10	а				
11	С	12	а	13	С	14	b	15	d				
16	а	17	d	18	d	19	С	20	d				
21	b	22	b	23	С	24	С	25	d				
26	b	27	С	28	а	29	b	30	а				
31	С	32	d	33	d	34	d	35	С				
36	d	37	а	38	а	39	С	40	С				
41	С	42	d	43	d	44	b	45	С				
46	b	47	а	48	а	49	С	50	d				
51	С	52	d	53	b	54	d	55	С				
56	b	57	b	58	b	59	d	60	а				
61	b	62	а	63	b	64	b	65	b				
66	С	67	С	68	С	69	С	70	а				
71	d	72	С	73	а	74	С	75	d				
76	С	77	d	78	b	79	а	80	а				
81	С	82	С	83	b	84	С						

Water or hydride of oxygen

1	b	2	b	3	С	4	b	5	С
6	а	7	b	8	b	9	С	10	b
11	а	12	а	13	d	14	d	15	d
16	С	17	b	18	b	19	С	20	d
21	С	22	а	23	а	24	С	25	С
26	С	27	d	28	а	29	е	30	С
31	С	32	С	33	С	34	d	35	d
36	а	37	а	38	С	39	С	40	d
41	С	42	d	43	С	44	b	45	a
46.	b								

Hydrogen peroxide

1	В	2	d	3	а	4	b	5	С
6	С	7	d	8	С	9	b	10	b
11	а	12	d	13	а	14	b	15	а
16	С	17	а	18	b	19	С	20	С
21	а	22	a	23	а	24	d	25	d
26	d	27	d	28	а	29	а	30	С
31	С	32	С	33	d	34	а	35	а
36	С	37	а	38	а	39	е	40	е





41 b

Critical Thinking Questions

1	С	2	d	3	С	4	d	5	acd
6	cd	7	abd	8	b	9	bd	10	cd
11	d	12	d	13	a	14	b	15	b
16	С	17	d	18	С	19	С	20	а
21	С	22	С	23	С	24	b	25	b
26	b	27	b	28	b	29	b	30	d
31	а	32	b						

Assertion & Reason

1	а	2	d	3	а	4	С	5	а
6	а	7	b	8	а	9	а	10	d

Answers and Solutions

Hydrogen and its preparation

- 1. (a) $NaH + H_2O \rightarrow NaOH + H_2 \uparrow$
- 2. (d) $KH + H_2O \rightarrow KOH + H_2 \uparrow$
- **3.** (a) Hydrogen burns in air with a light bluish flame.
- **4.** (a) Ortho and para hydrogen show different spin in a hydrogen molecule it does not show hydrogen isotopes.
- 5. (d) Boiling point of liquid hydrogen is lowest of given substances so it is distilled first.
- **6.** (d) $Mg + 2HNO_3 \rightarrow Mg(NO_3)_2 + H_2 \uparrow$
- **8.** (a) $Mg + 2H_2O \rightarrow Mg(OH)_2 + H_2 \uparrow$
- 10. (a) Ortho and para hydrogen differ in proton spin.
- 11. (c) $Mg + 2H_2O \rightarrow Mg(OH)_2 + H_2 \uparrow$
- 12. (a) $Mg + 2HCl \rightarrow MgCl_2 + H_2 \uparrow$
- 13. (c) Order of adsorption of $\ H_2$ (occlusion) is ${\it Colloidal\ Palladium > Palladium > Platinum > Gold > Nickel}$
- **14.** (b) Number of neutrons = Mass number Atomic number = 3 1 = 2
- 17. (d) Because AI has more affinity for oxygen than hydrogen.
- 18. (d) Helium is a noble gas and does not combine with hydrogen.
- 19. (c) Occlusion is the phenomenon of adsorption of hydrogen by metal.
- **20.** (d) CaH_2 is known as hydrolith.
- **21.** (b) Zn displaces hydrogen from the boiling solution of NaOH. $Zn + 2NaOH + 2H_2O \rightarrow Na_2[Zn(OH)_4] + H_2 \uparrow$
- **22.** (b) Occluded hydrogen is the hydrogen absorbed by the metal.
- 23. (c) Because dihydrogen is less reactive.

- **24.** (c) ${}_1H^3$ has 3 nucleons (1 proton + 2 neutrons) and one electron so sum of these is 3 + 1 = 4.
- **25.** (d) ${}_{1}^{2}D_{2} = (2 \text{ neutrons} + 2 \text{ protons}) = 4 \text{ nucleons}.$
- **26.** (b) Solubility of ionic compound is lower in heavy water.
- 27. (c) These allotropic forms have similar chemical properties.
- **28.** (a) It is 0.4 kl/mol.
- **29.** (b) $H^- = 1s^2$; $He = 1s^2$
- 30. (a) A fusion reaction is difficult to occur because positively charged nuclei repel each-other. At very high temperatures of the order of 10^6 to $10^7 \, K$, the nuclei may have sufficient energy to overcome the repulsive forces and fuse. It is for this reason, fusion reactions are also called thermonuclear reactions. Hence, hydrogen can be fused to form helium at high temperature and high pressure.
- 31. (c) It is Bosch process.
- **32.** (d) Gold is a noble metal.
- **33.** (d) $Ca + H_2 \rightarrow CaH_2 \xrightarrow{2H_2O} Ca(OH)_2 + 2H_2$
- **34.** (d) Hydrogen can loose one electron (*e.g. HF*). It can gain one electron (*e.g. NaH*), Hydrogen can also share one electron (*e.g. H _ H*)
- **35.** (c) Hydrogen is a much lighter element than alkali metals or halogen.
- **36.** (d) Heavy water is not unstable.
- 37. (a) $M^+H^- \to M^+ + H^-$

$$\boldsymbol{H}^{-} \rightarrow \frac{1}{2}\,\boldsymbol{H}_{2} + \boldsymbol{e}^{-} \, (\mathrm{At \ anode})$$

- **38.** (a) F_2 has maximum tendency to react with hydrogen. the decreasing order of reactivity is $F_2 > Cl_2 > Br_2 > I_2$.
- **39.** (c) It acts both as a reducing agent and oxidising agent.
- **40.** (c) $H \to H^+ + e^-$

$$H + e^- \rightarrow H^-$$

- 41. (c) IE of H is 1312 kJ/mole.
- 1E of *Cl* is 1255 *kJ*/*mole*.
- **42.** (d) Alkali metals are good reducing agents because of low ionization energy and hydrogen also shows same character.
- 43. (d) Position of hydrogen in the periodic table is not fully justified.
- **44.** (b) $H_2 + Cl_2 \rightarrow H^+Cl^-$. In this hydrogen has positive oxidation state.
- **45.** (c) For example HF, NaH, H_2