

Ordinary Thinking

Objective Questions

Nucleus (Stability and Reaction)

- Nucleons are [CPMT 1982]
 - Protons and electrons
 - Protons and neutrons
 - Electrons and neutrons
 - Electrons, protons and neutrons
- A deuteron contains [NCERT 1982; CPMT 1994]
 - A neutron and a positron
 - A neutron and a proton
 - A neutron and two protons
 - A proton and two neutrons
- The nucleus of radioactive element possesses
 - Low binding energy
 - High binding energy
 - Zero binding energy
 - High potential energy
- On bombarding ${}_7N^{14}$ with α -particles, the nuclei of the product formed after the release of a proton will be or In nuclear reaction ${}_7N^{14} + {}_2He^4 \rightarrow {}_Z X^A + {}_1H^1$, the term ${}_Z X^A$ represents [NCERT 1979; MP PMT 1989; JIPMER 1991; BHU 1995; MP PET 1996; BHU 1996]
 - ${}_8O^{17}$
 - ${}_9F^{18}$
 - ${}_9F^{17}$
 - ${}_8O^{18}$
- Nuclear energy is based on the conversion of
 - Protons into neutrons
 - Mass into energy
 - Neutrons into protons
 - Uranium into radium
- Positron has nearly the same weight as that of [NCERT 1975; JIPMER 1991; BHU 1995]
 - α -particle
 - Proton
 - Neutron
 - Electron
- In the reaction ${}_3Li^6 + (?) \rightarrow {}_2He^4 + {}_1H^3$. The missing particle is [CPMT 1983, 84]
 - Electron
 - Neutron
 - Proton
 - Deuteron
- The ${}_6C^{14}$ in upper atmosphere is generated by the nuclear reaction [MP PET 1993]
 - ${}_7N^{14} + {}_1H^1 \longrightarrow {}_6C^{14} + {}_{+1}e^0 + {}_1H^1$
 - ${}_7N^{14} \longrightarrow {}_6C^{14} + {}_{+1}e^0$
 - ${}_7N^{14} + {}_0n^1 \longrightarrow {}_6C^{14} + {}_1H^1$
 - ${}_7N^{14} + {}_1H^3 + {}_0n^1 \longrightarrow {}_6C^{14} + {}_2He^4$
- Deuterons when bombarded on a nuclide produce ${}_{18}Ar^{38}$ and neutrons. The target is [CPMT 1982, 87]
 - ${}_{17}Cl^{35}$
 - ${}_{19}K^{27}$
 - ${}_{17}Cl^{37}$
 - ${}_{19}K^{39}$
- Which can be used for carrying out nuclear reaction [AFMC 2003]
 - Uranium – 238
 - Neptunium – 239
 - Thorium – 232
 - Plutonium – 239
- On comparing chemical reactivity of C^{12} and C^{14} , it is revealed that
 - C^{12} is more reactive
 - C^{14} is more reactive
 - Both are inactive
 - Both are equally active
- The radionuclide ${}_{90}^{234}Th$ undergoes two successive β -decays followed by one α -decay. The atomic number and the mass number respectively of the resulting radionuclide are
 - 92 and 234
 - 94 and 230
 - 90 and 230
 - 92 and 230
- Hydrogen and deuterium differ in [CPMT 1980]
 - Reactivity with oxygen
 - Reactivity with chlorine
 - Melting point
 - Reducing action
- A nuclear reaction must be balanced in terms of
 - Only energy
 - Only mass
 - Mass and energy
 - None of these
- In the following nuclear reaction, the other product is ${}_{52}Te^{130} + {}_1H^2 \longrightarrow {}_{53}I^{131} + ?$ [MP PET 1991]
 - Positron
 - Alpha particle
 - One neutron
 - Proton
- ${}_{96}^{248}Po \longrightarrow {}_{88}^{208}Pb + {}_8^{40}He + {}_0^1e^0$ is due to [MP PMT 1991]
 - Loss of α -particles
 - Loss of β -particles
 - Loss of positron
 - Electron loss
- Positronium is the name given to an atom-like combination formed between [NCERT 1980; JIPMER 1991]
 - A positron and a proton
 - A positron and a neutron
 - A positron and α -particle
 - A positron and an electron
- An electrically charged atom or a group of atoms is known as
 - A meson
 - A proton
 - An ion
 - A cyclotron
- The charge on positron is equal to the charge on which one of the following [NCERT 1977]
 - Proton
 - Electron
 - α -particle
 - Neutron
- In the nuclear reaction ${}_{12}Mg^{24} + {}_2He^4 \longrightarrow {}_{10}Ne^{20} + ?$ The product nucleus is [BHU 1987]
 - ${}_{13}Al^{27}$
 - ${}_{14}Si^{27}$
 - ${}_{13}Al^{28}$
 - ${}_{12}Mg^{25}$
- ${}_6C^{14}$ is formed from ${}_7N^{14}$ in the upper atmosphere by the action of the fundamental particle [Orissa JEE 2002]
 - Positron
 - Neutron
 - Electron
 - Proton
- In the nuclear reaction ${}_{92}U^{238} \longrightarrow {}_{82}Pb^{206} + x {}_2He^4 + y {}_{-1}\beta^0$ the value of x and y are respectively [Orissa JEE 2002]
 - 8, 6
 - 6, 4
 - 6, 8
 - 8, 10
- If an isotope of hydrogen has two neutrons in its atom, its atomic number and atomic mass number will respectively be


[CBSE 1992]

- (a) 2 and 1 (b) 3 and 1
(c) 1 and 1 (d) 1 and 3

24. Which one of the following nuclear transformation is (n, p) type [AIIMS 1980, 83]

- (a) ${}_3\text{Li}^7 + {}_1\text{H}^1 \longrightarrow {}_4\text{Be}^7 + {}_0n^1$
(b) ${}_{33}\text{As}^{75} + {}_2\text{He}^4 \longrightarrow {}_{35}\text{Br}^{78} + {}_0n^1$
(c) ${}_{83}\text{Bi}^{209} + {}_1\text{H}^2 \longrightarrow {}_{84}\text{Po}^{210} + {}_0n^1$
(d) ${}_{21}\text{Sc}^{45} + {}_0n^1 \longrightarrow {}_{20}\text{Ca}^{45} + {}_1\text{H}^1$

25. What is X in the following nuclear reaction



- (a) ${}_{+1}e^0$ (b) ${}_0n^1$
(c) γ (d) ${}_{-1}e^0$

26. In the reaction ${}_{93}\text{Np}^{239} \longrightarrow {}_{94}\text{Pu}^{239} + (?)$, the missing particle is [MNR 1987]

- (a) Proton (b) Positron
(c) Electron (d) Neutron

27. According to the nuclear reaction ${}_4\text{Be} + {}_2\text{He}^4 \rightarrow {}_6\text{C}^{12} + {}_0n^1$, mass number of (Be) atom is

[AFMC 2002]

- (a) 4 (b) 9
(c) 7 (d) 6

28. Which of the following nuclides has the magic number of both protons and neutrons [EAMCET 1989]

- (a) ${}_{50}\text{Sn}^{115}$ (b) ${}_{82}\text{Pb}^{206}$
(c) ${}_{82}\text{Pb}^{208}$ (d) ${}_{50}\text{Sn}^{118}$

29. In the carbon cycle, from which hot stars obtain their energy, the ${}_6\text{C}^{14}$ nucleus is

- (a) Completely converted into energy
(b) Regenerated at the end of the cycle
(c) Combined with oxygen to form carbon monoxide
(d) Broken up into its constituents protons and neutrons

30. The atomic mass of lead is 208 and atomic number is 82. The atomic mass of bismuth is 209 and atomic number is 83. The ratio of n/p in the atom is [EAMCET 1982]

- (a) Higher of lead (b) Higher of bismuth
(c) Same (d) None of these

31. Which of the following is an n, p reaction [BHU 1995]


- (a) ${}_5\text{C}^{13} + {}_1\text{H}^1 \longrightarrow {}_6\text{C}^{14}$
(b) ${}_7\text{N}^{14} + {}_1\text{H}^1 \longrightarrow {}_8\text{O}^{15}$
(c) ${}_{13}\text{Al}^{27} + {}_0n^1 \longrightarrow {}_{12}\text{Mg}^{27} + {}_1\text{H}^1$
(d) ${}_{92}\text{U}^{235} + {}_0n^1 \longrightarrow {}_{54}\text{Xe}^{140} + {}_{38}\text{Sr}^{93} + 3{}_0n^1$

32. Which one of the following statements is incorrect

[MP PET 1997]

- (a) Mass defect is related with binding energy
(b) 'Meson' was discovered by Yukawa
(c) The size of the nucleus is of the order of $10^{-12} - 10^{-13} \text{ cm}$
(d) Magnetic quantum number is a measure of 'orbital angular momentum' of the electron

33. In the sequence of following nuclear reactions

The value of n will be

[MP PMT 1999]

- (a) 3 (b) 4
(c) 5 (d) 6

34. The introduction of a neutron into the nuclear composition of an atom would lead to a change in [MNR 1995]

- (a) The number of the electrons also
(b) The chemical nature of the atom
(c) Its atomic number
(d) Its atomic weight

35. The composition of tritium (${}_1\text{H}^3$) is

[Manipal MEE 1995; DPMT 1982,96]

- (a) 1 electron, 1 proton, 1 neutron
(b) 1 electron, 2 protons, 1 neutron
(c) 1 electron, 1 proton, 2 neutrons
(d) 1 electron, 1 proton, 3 neutrons

36. Identify 'X' in ${}_{16}\text{S}^{32} + \text{X} \rightarrow {}_{15}\text{P}^{30} + {}_2\text{He}^4$

- (a) ${}_1\text{H}^1$ (b) ${}_1\text{D}^2$
(c) ${}_0n^1$ (d) e^-


37. In terms of energy 1 a. m.u. is equal to

[MP PET/PMT 1998]

- (a) 100 J (b) 931.1 MeV
(c) 931.1 kcal (d) 10^7 erg

38. Positron is [AIIMS 1997]

- (a) Electron with +ve charge
(b) A helium nucleus
(c) A nucleus with two protons
(d) A nucleus with one neutron and one proton



In the above sequence of reaction, the elements which are isotopes of each other are [JIPMER 1997]

- (a) X and W (b) Y and Z
(c) X and Z (d) None of these

40. Stable nuclides are those whose n/p ratio is [MP PMT 1993]

- (a) $n/p = 1$ (b) $n/p = 2$
(c) $n/p > 1$ (d) $n/p < 1$

41. Neutrino has [NCERT 1981]

- (a) Charge +1, mass 1 (b) Charge 0, mass 0
(c) Charge -1, mass 1 (d) Charge 0, mass 1

42. Which one of the following nuclear reaction is correct

[CPMT 1997]

- (a) ${}_6\text{C}^{13} + {}_1\text{H}^1 \rightarrow {}_7\text{N}^{13} + \beta^- + \nu^-$
(b) ${}_{11}\text{Na}^{23} + {}_1\text{H}^1 \rightarrow {}_{10}\text{Ne}^{20} + {}_2\text{He}^4$
(c) ${}_{13}\text{Al}^{23} + {}_0n^1 \rightarrow {}_{11}\text{Na}^{23} + e^0$
(d) None of these

43. Formation of nucleus from its nucleons is accompanied by

[NCERT 1975; RPET 2000]

- (a) Decrease in mass (b) Increase in mass
(c) No change of mass (d) None of them

44. A particle having the same charge and 200 times greater mass than that of electron is

- (a) Positron (b) Proton
(c) Neutrino (d) Meson

45. The positron is [AFMC 1997]
 (a) ${}_{-1}e^0$ (b) ${}_{+1}e^0$
 (c) ${}_1H^1$ (d) ${}_0n^1$
46. Which of the following is the most stable atom [AFMC 1997]
 (a) Bi (b) Al
 (c) U (d) Pb
47. The positron is discovered by [RPMT 1997]
 (a) Pauling (b) Anderson
 (c) Yukawa (d) Segar
48. The nucleus of an atom is made up of X protons and Y neutrons. For the most stable and abundant nuclei [NCERT 1980]
 (a) X and Y are both even (b) X and Y are both odd
 (c) X is even and Y is odd (d) X is odd and Y is even
49. Atom A possesses higher values of packing fraction than atom B. The relative stabilities of A and B are
 (a) A is more stable than B
 (b) B is more stable than A
 (c) A and B both are equally stable
 (d) Stability does not depend on packing fraction
50. How many neutrons are present in the nucleus of Ra [CPMT 1980]
 (a) 88 (b) 226
 (c) 140 (d) 138
51. In a nuclear explosion, the energy is released in the form of [CPMT 1994]
 (a) Kinetic energy (b) Electrical energy
 (c) Potential energy (d) None of these
52. In equation ${}_{11}Na^{23} + {}_1H^1 \rightarrow {}_{12}Mg^{23} + x$, x represents [MP PMT 1990; MP PET 1999]
 (a) Neutron (b) Deuteron
 (c) α -particle (d) Positron
53. Which of the following atomic mass of uranium is the most radioactive [AFMC 1997]
 (a) 238 (b) 235
 (c) 226 (d) 248
54. Which of the following particle is emitted in the reaction ${}_{13}Al^{27} + {}_2He^4 \rightarrow {}_{14}P^{30} + \dots$ [DCE 1999]
 (a) ${}_0n^1$ (b) ${}_{-1}e^0$
 (c) ${}_1H^1$ (d) ${}_1H^2$
55. Which of the following sub-atomic particles is not present in an atom [JIPMER 1999]
 (a) Neutron (b) Proton
 (c) Electron (d) Positron
56. Electromagnetic radiation with maximum wave length is [DCE 2000; UPSEAT 2000]
 (a) Ultraviolet ray (b) Radiowave
 (c) X-ray (d) Infrared
57. Neutrons are obtained by [JIPMER 1999]
 (a) Bombardment of Ra with β -particles
 (b) Bombardment of Be with α -particles
 (c) Radioactive disintegration of uranium
 (d) None of these
58. In the reaction, $Po \xrightarrow{-\alpha} Pb \xrightarrow{-\beta} Bi$, if Bi, belongs to group 15, to which Po belongs [DCE 2000]
 (a) 14 (b) 15
 (c) 13 (d) 16
59. In the nuclear reaction ${}_4Be(p, \alpha)X$, the X is [MP PMT 2000]
 (a) ${}_2He^4$ (b) ${}_3Li^6$
 (c) ${}_3Li^7$ (d) ${}_4Be^8$
60. Which of the following does not contain number of neutrons equal to that of ${}^{40}_{18}Ar$ [MP PMT 2000]
 (a) ${}^{41}_{19}K$ (b) ${}^{43}_{21}Sc$
 (c) ${}^{40}_{21}Sc$ (d) ${}^{42}_{20}Ca$
61. Nuclear reactivity of Na and Na^+ is same because both have
 (a) Same electron and proton
 (b) Same proton and same neutron
 (c) Different electron and proton
 (d) Different proton and neutron
62. Which of the following is the heaviest metal [MH CET 2001]
 (a) Hg (b) Pb
 (c) Ra (d) U
63. In the following reaction, x will be ${}_{29}Cu^{64} \rightarrow {}_{28}Ni^{64} + x$
 (a) A proton (b) An electron
 (c) A neutron (d) A positron
64. Which one out of the following statements is not correct for ortho and para hydrogen [Orissa JEE 2002]
 (a) They have different boiling point
 (b) Ortho form is more stable than para form
 (c) They differ in the spin of their protons
 (d) The ratio of ortho to para hydrogen increases with increase in temperature and finally pure ortho form is obtained
65. For the nuclear reaction, ${}^{24}_{12}Mg + {}^2_1D \rightarrow \alpha + ?$, the missing nucleide is [Kurukshetra CEE 2002]
 (a) ${}^{22}_{11}Na$ (b) ${}^{23}_{11}Na$
 (c) ${}^{23}_{12}Mg$ (d) ${}^{26}_{12}Mg$
66. ${}_ZX^M + {}_2He^4 \rightarrow {}_{15}P^{30} + {}_0n^1$. Then [KCET 2002]
 (a) $Z = 12, M = 27$ (b) $Z = 13, M = 27$
 (c) $Z = 12, M = 17$ (d) $Z = 13, M = 28$
67. An element ${}_{96}X^{227}$ emits 4α and 5β particles to form new element Y. Then atomic number and mass number of Y are
 (a) 93; 211 (b) 211; 93
 (c) 212; 88 (d) 88; 211
68. Meson was discovered [MH CET 2004]
 (a) Yukawa (b) Austin
 (c) Moseley (d) Einstein

Radioactivity and α , β and γ rays

1. Which of the following does not contain material particles [BHU 2002]
 (a) Alpha rays (b) Beta rays
 (c) Gamma rays (d) Canal rays
2. Radioactive substances emit γ -rays, which are

- (a) +ve charged particle (b) -ve charged particle
(c) Massive particle (d) Packet of energy
[Orissa JEE 2002]
3. Which statement is incorrect [CPMT 1982]
(a) α -rays have more penetrating power than β -rays
(b) α -rays have less penetrating power than γ -rays
(c) β -rays have less penetrating power than γ -rays
(d) β -rays have more penetrating power than α -rays
4. The velocity of α -rays is approximately [CPMT 1982]
(a) Equal to that of the velocity of light
(b) 1/10 of the velocity of light
(c) 10 times more than the velocity of light
(d) Uncomparable to the velocity of light
5. The radiations having high penetrating power and not affected by electrical and magnetic field are [Kerala CET 1992]
(a) Alpha rays (b) Beta rays
(c) Gamma rays (d) Neutrons
6. Alpha particles are times heavier (approximately) than neutrons [CPMT 1971]
(a) 2 (b) 4
(c) 3 (d) $2\frac{1}{2}$
7. Uranium ${}_{92}\text{U}^{235}$ on bombardment with slow neutrons produces [CPMT 1982]
(a) Deutrons (b) Fusion reaction
(c) Fission reaction (d) Endothermic reaction
8. α -particles can be detected using [AIIMS 2005]
(a) Thin aluminum sheet (b) Barium sulphate
(c) Zinc sulphide screen (d) Gold foil
9. Alpha rays consist of a stream of [BHU 1979]
(a) H^+ (b) He^{+2}
(c) Only electrons (d) Only neutrons
10. Which is the correct statement [CPMT 1971]
(a) Isotopes are always radioactive
(b) β -rays are always negatively charged particles
(c) α -rays are always negatively charged particles
(d) γ -rays can be deflected in magnetic field
11. The α -particle is identical with [CPMT 1972, 82, 86; BHU 1984; MP PMT 1990, 91, 93; MP PET 1999]
(a) Helium nucleus
(b) Hydrogen nucleus
(c) Electron
(d) Proton
12. If by mistake some radioactive substance gets inside the human body, then from the point of view of radiation damage, the most harmful will be the one which emits [DPMT 1986]
(a) γ -rays (b) Neutrons
(c) β -particles (d) α -particles
13. Radioactivity was discovered by [CPMT 1983, 88; DPMT 1982; AMU 1983; MADT Bihar 1982]
(a) Henry Becquerel (b) Rutherford
(c) J. J. Thomson (d) Madam Curie
14. Which of the following is radioactive element [CPMT 1988]
(a) Sulphur (b) Polonium
(c) Tellurium (d) Selenium
15. Penetrating power of α -particle is [MP PMT 2002]
(a) More than γ -rays (b) More than β -rays
(c) Less than β -rays (d) None of these
16. β -particle is emitted in radioactivity by [AIEEE 2002; MP PMT 2004]
(a) Conversion of proton to neutron
(b) Form outermost orbit
(c) Conversion of neutron to proton
(d) β -particle is not emitted
17. α -rays have [CPMT 1973, 78; NCERT 1977]
(a) Positive charge
(b) Negative charge
(c) No charge
(d) Sometimes positive charge and sometimes negative charge
18. X-rays are produced due to [JIPMER 2002]
(a) Bombarding of electrons on solids
(b) Bombarding of α -particle on solids
(c) Bombarding of γ -rays on solids
(d) Bombarding of neutron on solids
19. Choose the element which is not radioactive [CPMT 1988]
(a) Cm (b) No
(c) Mo (d) Md
20. A magnet will cause the greatest deflection of [MP PMT 1991]
(a) γ -rays (b) β -rays
(c) α -rays (d) Neutrons
21. Of the following radiations, the one most easily stopped by air is [MP PMT 1990]
(a) α -rays (b) β -rays
(c) γ -rays (d) X-rays
22. Uranium ultimately decays into a stable isotope of [MP PET 1995]
(a) Radium (b) Carbon
(c) Lead (d) Neptunium
23. Which leaves no track on Wilson cloud chamber [AFMC 1988]
(a) Electrons (b) Protons
(c) α -particles (d) Neutrons
24. Which has the least penetrating power [CPMT 1994]
(a) β -rays (b) α -rays
(c) γ -rays (d) X-rays
25. There exists on γ -rays [MP PMT 1996; Pb. PMT 2004; EAMCET 2004]
(a) Positive charge
(b) Negative charge
(c) No charge
(d) Sometimes positive charge, sometimes negative charge
26. Which is not emitted by radioactive substance [AIIMS 1997]
(a) α -rays (b) β -rays
(c) Positron (d) Proton
27. The radiations from a naturally occurring radio element, as seen after deflection in a magnetic field in one direction, are [IIT 1984; MP PMT 1986; MP PET/PMT 1988 JIPMER 1999]
(a) Definitely α -rays (b) Definitely β -rays
(c) Both α and β -rays (d) Either α or β -rays
28. The ${}_{88}\text{Ra}^{226}$ is [AIIMS 2001]

- (a) n -mesons (b) u -mesons (c) 0.0018 amu (d) 1.8 amu
(c) Radioactive (d) Non-radioactive (e) 18 amu
29. During β -decay [UPSEAT 2001]
(a) An atomic electron is ejected
(b) An electron which is already present within the nucleus is ejected
(c) A neutron in the nucleus decays emitting an electron
(d) A part of binding of the nucleus is converted into an electron
30. The element californium belongs to the family of [UPSEAT 2002]
(a) Actinide series (b) Alkali metal family
(c) Alkaline earth family (d) Lanthanide series
31. Which of the following is not deflected by magnetic field [MP PMT 2001]
(a) Deuteron (b) Positron
(c) Proton (d) Photon
32. Which of the following can be used to convert ${}^{14}_7\text{N}$ into ${}^{17}_8\text{O}$ [MP PMT 2001]
(a) Deuteron (b) Proton
(c) α -particle (d) Neutron
33. The amount of energy, which is required to separate the nucleons from a nucleus. The energy is called [UPSEAT 2001]
(a) Binding energy (b) Lattice energy
(c) Kinetic energy (d) None of these
34. What happens when α -particle is emitted [CBSE PMT 1989; JIPMER 2002]
(a) Mass number decreases by 12 unit, atomic number decreases by 4 unit
(b) Mass number decreases by 4 unit, atomic number decreases by 2 unit
(c) Only mass number decreases
(d) Only atomic number decreases
35. The charge on gamma rays is [Pb. PMT 2004; EAMCET 2004]
(a) Zero (b) +1
(c) -1 (d) +2
36. A nuclear reaction is accompanied by loss of mass equivalent to 0.01864 amu. Energy liberated is [DCE 2002]
(a) 931 MeV (b) 186.6 MeV
(c) 17.36 MeV (d) 460 MeV
37. Nuclear theory of the atom was put forward by [KCET 2004]
(a) Rutherford (b) Aston
(c) Neils Bohr (d) J.J. Thomson
38. Decrease in atomic number is observed during [IIT 1998]
(a) Alpha emission (b) Beta emission
(c) Positron emission (d) Electron capture
39. Calculate mass defect in the following reaction
 ${}_1\text{H}^2 + {}_1\text{H}^3 \rightarrow {}_1\text{He}^4 + {}_0\text{n}^1$
(Given : mass $\text{H}^2 = 2.014$, $\text{H}^3 = 3.016$, $\text{He} = 4.004$, $n = 1.008 \text{ amu}$) [Kerala CET 2005]
(a) 0.018 amu (b) 0.18 amu

Causes of radioactivity and Group displacement law

1. ${}_{95}\text{Am}^{241}$ and ${}_{90}\text{Th}^{234}$ belong respectively to [MP PMT 1999]
(a) $4n$ and $4n+1$ radioactive disintegration series
(b) $4n+1$ and $4n+2$ radioactive disintegration series
(c) $4n+1$ and $4n+3$ radioactive disintegration series
(d) $4n+1$ and $4n$ radioactive disintegration series
2. Group displacement law states that the emission of α or β particles results in the daughter element occupying a position, in the periodic table, either to the left or right of that of the parent element. Which one of the following alternatives gives the correct position of the daughter element
- | On emission of α particles | On emission of β particles |
|-----------------------------------|----------------------------------|
| (a) 2 groups to the right | 1 group to the right |
| (b) 2 groups to the right | 1 group to the left |
| (c) 2 groups to the left | 1 group to the left |
| (d) 2 groups to the left | 1 group to the right |
3. The nuclides (A nuclide is the general name for any nuclear species) ${}^{12}_{26}\text{Fe}^{56}$ and ${}^{238}_{92}\text{U}$ have 12, 56 and 238 nucleons respectively in the nuclei. The total number of nucleons in a nucleus is equal to [NCERT 1975]
(a) The total number of neutrons in the nucleus
(b) The total number of neutrons in the atom
(c) The total number of protons in the nucleus
(d) The total number of protons and neutrons in the nucleus
4. Radioactivity is due to [DPMT 1983, 89; AIIMS 1988]
(a) Stable electronic configuration
(b) Unstable electronic configuration
(c) Stable nucleus
(d) Unstable nucleus
5. Radioactive disintegration differs from a chemical change in being [MNR 1991]
(a) An exothermic change
(b) A spontaneous process
(c) A nuclear process
(d) A unimolecular first order reaction
6. ${}^{238}_{92}\text{U}$ emits 8 α -particles and 6 β -particles. The neutron/proton ratio in the product nucleus is [AIIMS 2005]
(a) 60/41 (b) 61/40
(c) 62/41 (d) 61/42
7. The element with atomic number 84 and mass number 218 change to other element with atomic number 84 and mass number 214. The number of α and β -particles emitted are respectively [CPMT 1989]
(a) 1, 3 (b) 1, 4
(c) 1, 2 (d) 1, 5
8. A radium ${}_{88}\text{Ra}^{224}$ isotope, on emission of an α -particle gives rise to a new element whose mass number and atomic number will be [CPMT 1980; EAMCET 1985; MP PMT 1993]
(a) 220 and 86 (b) 225 and 87
(c) 228 and 88 (d) 224 and 86

9. ${}_{89}\text{Ac}^{231}$ gives ${}_{82}\text{Pb}^{207}$ after emission of some α and β -particles. The number of such α and β -particles are respectively [MP PMT 1993; UPSEAT 2001]
- (a) 5, 6 (b) 6, 5
(c) 7, 5 (d) 5, 7
10. The number of α and β -particles emitted in the nuclear reaction ${}_{90}\text{Th}^{228} \rightarrow {}_{83}\text{Bi}^{212}$ are respectively [MNR 1992; MP PMT 1993; AFMC 1998, 2001; MH CET 1999; UPSEAT 2000, 01; AMU 2001; CPMT 2002]
- (a) 4, 1 (b) 3, 7
(c) 8, 1 (d) 4, 7
11. The number of neutrons in the parent nucleus which gives N^{14} on β -emission and the parent nucleus is [EAMCET 1985; MNR 1992; Kurukshetra CEE 1998; UPSEAT 2000, 01]
- (a) $8, \text{C}^{14}$ (b) $6, \text{C}^{12}$
(c) $4, \text{C}^{13}$ (d) None of these
12. After the emission of α -particle from the atom ${}_{92}\text{X}^{238}$, the number of neutrons in the atom will be [MNR 1993; UPSEAT 1999, 2001, 02]
- (a) 138 (b) 140
(c) 144 (d) 150
13. When a radioactive element emits an electron the daughter element formed will have [EAMCET 1988; MP PET 1994]
- (a) Mass number one unit less
(b) Atomic number one unit less
(c) Mass number one unit more
(d) Atomic number one unit more
14. If the amount of radioactive substance is increased three times, the number of atoms disintegrated per unit time would [MP PMT 1994]
- (a) Be double (b) Be triple
(c) Remain one third (d) Not change
15. β -particles are emitted from the atom
- (a) Due to disintegration of neutron
(b) Due to disintegration of proton
(c) Due to removal of electron from K shell
(d) Due to removal of electron from outermost orbit
16. Nd ($Z = 60$) is a member of group -3 in periodic table. An isotope of it is β -active. The daughter nuclei will be a member of
- (a) Group -3 (b) Group -4
(c) Group -1 (d) Group -2
17. Number of neutrons in a parent nucleus X , which gives ${}_{7}\text{N}^{14}$ nucleus after two successive β emissions would be [CBSE PMT 1998; MP PMT 2003]
- (a) 9 (b) 8
(c) 7 (d) 6
18. The disintegration of an isotope of sodium, ${}_{11}\text{Na}^{24} \rightarrow {}_{12}\text{Mg}^{24} + {}_{-1}\text{e}^0$ shown is due to [AMU (Engg.) 2000]
- (a) The emission of β -radiation
(b) The formation of a stable nuclide
(c) The fall in the neutron : proton ratio
(d) None of these
19. After losing a number of α and β -particles, ${}_{92}\text{U}^{238}$ is changed to ${}_{82}\text{Pb}^{206}$. The total number of α -particles lost in this process is [UPSEAT 1999, 2000]
- (a) 10 (b) 5
(c) 8 (d) 32
20. Which element is the end product of each natural radioactive series [MP PMT 1993; UPSEAT 2001]
- (a) Sn (b) Bi
(c) Pb (d) C
21. ${}_{13}^{27}\text{Al}$ is a stable isotope. ${}_{13}^{29}\text{Al}$ is expected to disintegrate by [IIT 1996; UPSEAT 2001]
- (a) α -emission (b) β -emission
(c) Positron emission (d) Proton emission
22. An isotope ${}_Y\text{A}^X$ undergoes a series of m alpha and n beta disintegration to form a stable isotope ${}_{Y-10}\text{B}^{X-32}$. The values of m and n are respectively [MP PET 1995]
- (a) 6 and 8 (b) 8 and 10
(c) 5 and 8 (d) 8 and 6
23. During a β -decay the mass of the atomic nucleus [MP PET 1996]
- (a) Decreases by one unit (b) Increases by one unit
(c) Decreases by two units (d) Remains unaffected
24. Which one of the following notations shows the product incorrectly [MP PET/2000]
- (a) ${}_{96}^{242}\text{Cm}(\alpha, 2n){}_{97}^{243}\text{Bk}$ (b) ${}_{5}^{10}\text{B}(\alpha, n){}_{7}^{13}\text{N}$
(c) ${}_{7}^{14}\text{N}(n, p){}_{6}^{14}\text{C}$ (d) ${}_{14}^{28}\text{Si}(d, n){}_{15}^{29}\text{P}$
25. An atom has mass number 232 and atomic number 90. How many α -particles should it emit after emission of two β -particles, so that the new element's atom has mass number 212 and atomic number 82
- (a) 4 (b) 5
(c) 6 (d) 3
26. After the emission of one α -particle followed by one β -particle from the atom of ${}_{92}\text{X}^{238}$, the number of neutrons in the atom will be [CBSE PMT 1995]
- (a) 142 (b) 146
(c) 144 (d) 143
27. A nuclide of an alkaline earth metal undergoes radioactive decay by emission of the α -particles in succession. The group of the periodic table to which the resulting daughter element would belong is [CBSE PMT 2005]
- (a) Gr.14 (b) Gr.16
(c) Gr.4 (d) Gr.6
28. Which one of the following is not correct [MP PMT 1997]
- (a) ${}_3\text{Li}^7 + {}_1\text{H}^1 \rightarrow {}_4\text{Be}^7 + {}_0\text{n}^1$
(b) ${}_{21}\text{Sc}^{45} + {}_0\text{n}^1 \rightarrow {}_{20}\text{Ca}^{45} + {}_0\text{n}^1$
(c) ${}_{33}\text{As}^{75} + {}_2\text{He}^4 \rightarrow {}_{35}\text{Br}^{78} + {}_0\text{n}^1$
(d) ${}_{83}\text{Bi}^{209} + {}_1\text{H}^2 \rightarrow {}_{84}\text{Po}^{210} + {}_0\text{n}^1$
29. The end product of $(4n+2)$ radioactive disintegration series is [MP PET 1997; Pb. PMT 1998; BHU 2000]
- (a) ${}_{82}\text{Pb}^{208}$ (b) ${}_{82}\text{Pb}^{206}$
(c) ${}_{82}\text{Pb}^{207}$ (d) ${}_{83}\text{Bi}^{210}$

30. The element ${}_{90}\text{Th}$ belongs to thorium series. Which of the following will act as the end product of the series [BHU 2005]
- (a) ${}_{82}\text{Pb}$ (b) ${}_{84}\text{Po}$
(c) ${}_{86}\text{Rn}$ (d) ${}_{88}\text{Ra}$
31. On bombarding ${}_{8}\text{O}^{16}$ with deuterons, the nuclei of the product formed will be [NCERT 1978]
- (a) ${}_{9}\text{F}^{18}$ (b) ${}_{9}\text{F}^{17}$
(c) ${}_{8}\text{O}^{17}$ (d) ${}_{7}\text{N}^{14}$
32. An element with atomic number 84 and mass number 218 loses one α -particle and two β -particles in three successive stages, the resulting element will have [NCERT 1979; CPMT 1990]
- (a) At. no. 84 and mass number 214
(b) At. no. 82 and mass number 214
(c) At. no. 84 and mass number 218
(d) At. no. 82 and mass number 218
33. Group displacement law was given by [DPMT 1984]
- (a) Becquerel (b) Rutherford
(c) Soddy and Fajan (d) Madam Curie
34. How many alpha particles are emitted in the nuclear transformation ${}_{84}\text{Po}^{215} \longrightarrow {}_{82}\text{Pb}^{211}$ [CPMT 1993]
- (a) 0 (b) 1
(c) 2 (d) 3
35. If uranium (mass no. 238 and atomic no. 92) emits α -particle, the product has mass number and atomic number [CPMT 1984, 90, 93, 94; MNR 1991; IIT 1981]
- (a) 234, 90 (b) 236, 92
(c) 238, 90 (d) 236, 90
36. Initial mass of a radioactive element is 40 g. How many grams of it would be left after 24 years, if its half-life period is 8 years [MP PMT 1985]
- (a) 2 (b) 5
(c) 10 (d) 20
37. What is the symbol for the nucleus remaining after ${}_{20}\text{Ca}^{42}$ undergoes β -emission [MNR 1987; UPSEAT 2000, 02]
- (a) ${}_{21}\text{Ca}^{42}$ (b) ${}_{20}\text{Sc}^{42}$
(c) ${}_{21}\text{Sc}^{42}$ (d) ${}_{21}\text{Sc}^{41}$
38. When a radioactive nucleus emits an α -particle, the mass of the atom [NCERT 1973, 82]
- (a) Increases and its at. number decreases
(b) Decreases and its at. number decreases
(c) Decreases and its at. number increases
(d) Remains same and its at. number decreases
39. A photon of hard gamma radiation knocks a proton out of ${}_{12}^{24}\text{Mg}$ nucleus to form [AIEEE 2005]
- (a) The isotope of parent nucleus
(b) The isobar of parent nucleus
(c) The nuclide ${}_{11}^{23}\text{Na}$
(d) The isobar of ${}_{11}^{23}\text{Na}$
40. ${}_{84}\text{Pb}^{210} \longrightarrow {}_{82}\text{Pb}^{206} + {}_2\text{He}^4$. From the above equation, deduce the position of polonium in the periodic table (lead belongs to group IV A) [AIIMS 1980]
- (a) II A (b) IV B
(c) VI B (d) VI A
41. Whenever the parent nucleus emits a β -particle, the daughter element is shifted in the periodic table [NCERT 1984]
- (a) One place to the right
(b) One place to the left
(c) Two places to the right
(d) Two places to the left
42. In the nuclear reaction ${}_{92}\text{U}^{238} \longrightarrow {}_{82}\text{Pb}^{206}$, the number of alpha and beta particles decayed are [DPMT 1983; MNR 1985; Roorkee Qualifying 1998]
- (a) $4\alpha, 3\beta$ (b) $8\alpha, 6\beta$
(c) $6\alpha, 4\beta$ (d) $7\alpha, 5\beta$
43. Atomic number after a β -emission from a nucleus having atomic number 40, will be [BHU 1981]
- (a) 36 (b) 39
(c) 41 (d) 44
44. A certain nuclide has a half-life period of 30 minutes. If a sample containing 600 atoms is allowed to decay for 90 minutes, how many atoms will remain [NCERT 1978]
- (a) 200 atoms (b) 450 atoms
(c) 75 atoms (d) 500 atoms
45. The reaction which disintegrates neutron is or neutron is emitted (which completes first) [IIT 1988; MP PMT 1991; KCET 2005]
- (a) ${}_{96}\text{Am}^{240} + {}_2\text{He}^4 \longrightarrow {}_{97}\text{Bk}^{244} + {}_{+1}\text{e}^0$
(b) ${}_{15}\text{P}^{30} \longrightarrow {}_{14}\text{Si}^{30} + {}_1\text{e}^0$
(c) ${}_6\text{C}^{12} + {}_1\text{H}^1 \longrightarrow {}_7\text{N}^{13}$
(d) ${}_{13}\text{Al}^{27} + {}_2\text{He}^4 \longrightarrow {}_{15}\text{P}^{30}$
- If ${}_{92}\text{U}^{236}$ nucleus emits one α -particle, the remaining nucleus will have [MP PMT 1976, 80; BHU 1985; CPMT 1980]
- (a) 119 neutrons and 119 protons
(b) 142 neutrons and 90 protons
(c) 144 neutrons and 92 protons
(d) 146 neutrons and 90 protons
47. α -rays have high ionization power because they possess [CPMT 1982]
- (a) Lesser kinetic energy
(b) Higher kinetic energy
(c) Lesser penetrating power
(d) Higher penetrating power
48. When radium atom which is placed in II group, loses an α -particle, a new element is formed which should be placed in group [CPMT 1979]
- (a) Second (b) First
(c) Fourth (d) Zero
49. Starting from radium, the radioactive disintegration process terminates when the following is obtained [CPMT 1979]
- (a) Lead (b) Radon
(c) Radium A (d) Radium B
50. The appreciable radioactivity of uranium minerals is mainly due to [NCERT 1980]
- (a) An uranium isotope of mass number 235
(b) A thorium isotope of mass number 232
(c) Actinium
(d) Radium
51. After losing a number of α and β -particles, ${}_{92}\text{U}^{238}$ changed to ${}_{82}\text{Pb}^{206}$. The total number of particles lost in this process is [MNR 1985]

- (a) 14 (b) 5
(c) 8 (d) 32
52. When an radioactive element emits an alpha particle, the daughter element is placed in the periodic table [MP PET 1991; MADT Bihar 1981]
(a) Two positions to the left of the parent element
(b) Two positions to the right of the parent element
(c) One position to the right of the parent element
(d) In the same position as the parent element
53. If the quantity of a radioactive element is doubled, then its rate of disintegration per unit time will be [NCERT 1972, 92; MP PET 1989]
(a) Unchanged
(b) Reduced to half
(c) Increased by $\sqrt{2}$ times
(d) Doubled
54. The number of α and β -particles emitted during the transformation of ${}_{90}\text{Th}^{232}$ to ${}_{82}\text{Pb}^{208}$ are respectively [MNR 1978; NCERT 1984; CPMT 1989; RPET 1999; MP PMT 2001; KCET 2003]
(a) 4, 2 (b) 2, 2
(c) 8, 6 (d) 6, 4
55. The atomic number of a radioactive element increases by one unit in
(a) Alpha emission (b) Beta emission
(c) Gamma emission (d) Electron capture
56. The end product of $(4n + 1)$ radioactive disintegration series is [MP PMT 1999]
(a) ${}_{83}\text{Bi}^{209}$ (b) ${}_{84}\text{Po}^{210}$
(c) ${}_{82}\text{Pb}^{208}$ (d) ${}_{82}\text{Pb}^{207}$
57. When a β -particle emits from the atom of an element, then [MP PET 1990]
(a) Atomic number increases by two units
(b) Atomic number increases by three units
(c) Atomic number decreases by one unit
(d) Atomic number increases by one unit
58. The number of β -particles emitted in radioactive change ${}_{92}\text{U}^{238} \rightarrow {}_{82}\text{Pb}^{206} + {}_2\text{He}^4$ is [KCET 2000]
(a) 2 (b) 4
(c) 6 (d) 10
59. If half-life of a certain radioactive nucleus is 1000 s, the disintegration constant is [MP PET 2001]
(a) $6.93 \times 10^{-2} \text{ s}^{-1}$ (b) $6.93 \times 10^{-4} \text{ s}^{-1}$
(c) $6.93 \times 10^{-4} \text{ s}^{-1}$ (d) $6.93 \times 10^{-3} \text{ s}^{-1}$
60. Radioactivity of naptunium stops when it is converted to [JIPMER 2001]
(a) Bi (b) Rn
(c) Th (d) Pb
61. The highest binding energy per nucleon will be for [AIIMS 2001]
(a) Fe (b) H_2
(c) O_2 (d) U
62. In the Thorium series, ${}_{90}\text{Th}^{232}$ loses total of 6 α -particles and 4 β -particles in ten stages. The final isotope produced in the series is [MP PET 2001]
(a) ${}_{82}\text{Pb}^{209}$ (b) ${}_{83}\text{Bi}^{209}$
(c) ${}_{82}\text{Pb}^{208}$ (d) ${}_{82}\text{Pb}^{206}$
63. All the nuclei from the initial element to the final element constitute a series which is called [Kerala (Med.) 2002]
(a) g-series (b) b-series
(c) b-g series (d) Disintegration series
64. The number of neutrons in the parent nucleus which gives N^{14} on β -emission is [Pb.CET 2004]
(a) 7 (b) 14
(c) 6 (d) 8
65. The nuclear binding energy for Ar (39.962384 amu) is: (given mass of proton and neutron are 1.007825 amu and 1.008665 amu respectively) [Pb.CET 2002]
(a) 343.81 MeV (b) 0.369096 MeV
(c) 931 MeV (d) None of these
66. The number α - and β - particles emitted respectively during the transformation of ${}_{90}^{232}\text{Th}$ to ${}_{82}^{208}\text{Pb}$ is [Kerala PMT 2004]
(a) 3, 6 (b) 6, 3
(c) 4, 6 [AMCET 1997] (d) 6, 4
(e) 6, 8
67. Consider the following nuclear reactions, ${}_{92}^{238}\text{M} \rightarrow {}_x^y\text{N} + 2 {}_2^4\text{He}$
 ${}_x^y\text{N} \rightarrow {}_B^A\text{L} + 2\beta^+$
The number of neutrons in the element L is [AIEEE 2004]
(a) 140 (b) 144
(c) 142 (d) 146
68. The number of α - and β - particles emitted when a radioactive element ${}_{90}\text{E}^{232}$ changes into ${}_{86}\text{G}^{220}$ will be [MP PET 2004]
(a) 5 and 4 (b) 2 and 3
(c) 3 and 2 (d) 4 and 1
69. The disintegration constant of radium with half-life 1600 years is [MHCET 2004]
(a) $2.12 \times 10^{-4} \text{ year}^{-1}$ (b) $4.33 \times 10^{-4} \text{ year}^{-1}$
(c) $3.26 \times 10^{-3} \text{ year}^{-1}$ (d) $4.33 \times 10^{-12} \text{ year}^{-1}$
70. The number of α and β - particles emitted in the nuclear reaction ${}_{92}\text{U}^{238} \rightarrow {}_{90}\text{Th}^{234} \rightarrow {}_{91}\text{Pa}^{234}$ are respectively [Pb.CET 2001]
(a) 1 and 1 (b) 1 and 2
(c) 2 and 1 (d) 2 and 2
71. In which radiation mass number and atomic number will not change [JEE Orissa]
(a) α (b) β
(c) γ (d) α and 2β
72. Disintegration constant for a radioactive substance is 0.58 hr^{-1} . Its half-life period [BHU 2004]
(a) 8.2 hr (b) 5.2 hr
(c) 1.2 hr (d) 2.4 hr
73. A radioactive nucleus will not emit [DPMT 2005]
(a) Alpha and beta rays simultaneously
(b) Beta and gamma rays simultaneously
(c) Gamma and alpha rays

- (d) Gamma rays only
74. ${}_{72}^{180}\text{X} \xrightarrow{2\alpha} \xrightarrow{\beta} \xrightarrow{\gamma} {}_Z^AX'$. Z and A are [DPMPT 2005]
 (a) 69, 172 (b) 172, 69
 (c) 180, 70 (d) 182, 68
75. Loss of a beta particle is equivalent to [J & K 2005]
 (a) Increase of one neutron only
 (b) Decrease of one neutron only
 (c) Both (a) and (b)
 (d) None of these

Rate of decay and Half-life

1. The half-life period of a radioactive substance is 8 years. After 16 years, the mass of the substance will reduce from starting 16.0 g to [MP PMT 1999]
 (a) 8.0 g (b) 6.0 g
 (c) 4.0 g (d) 2.0 g
2. The atomic mass of an element is 12.00710 amu. If there are 6 neutrons in the nucleus of the atom of the element, the binding energy per nucleon of the nucleus will be [MP PMT 1999]
 (a) 7.64 MeV (b) 76.4 MeV
 (c) 764 MeV (d) 0.764 MeV
 ($e^- = 0.00055$ amu, $p = 1.00814$ amu, $n = 1.00893$ amu)
3. Half-life period of a metal is 20 days. What fraction of metal does remain after 80 days [BHU 1996]
 (a) 1 (b) 1/16
 (c) 1/4 (d) 1/8
4. In the radioactive decay ${}_{92}\text{X}^{232} \rightarrow {}_{89}\text{Y}^{220}$, how many α and β - particles are ejected from X to form Y [CBSE 1999]
 (a) 3α and 3β (b) 5α and 3β
 (c) 3α and 5β (d) 5α and 5β
5. Which of the following does not take place by α - decay [MP PMT 1996]
 (a) ${}_{92}\text{U}^{238} \longrightarrow {}_{90}\text{Th}^{234}$ (b) ${}_{90}\text{Th}^{232} \longrightarrow {}_{88}\text{Ra}^{228}$
 (c) ${}_{88}\text{Ra}^{226} \longrightarrow {}_{86}\text{Rn}^{222}$ (d) ${}_{83}\text{Bi}^{213} \longrightarrow {}_{84}\text{Po}^{213}$
6. 1.0 g of a radioactive isotope was found to reduce to 125 mg after 24 hours. The half-life of the isotope is [MP PET 1996]
 (a) 8 hours (b) 24 hours
 (c) 6 hours (d) 4 hours
7. A radioactive element decays at such a rate that after 15 minutes only 1/10 of the original amount is left. How many more minutes will be needed when only 1/100 of the original amount will be left
 (a) 1.5 minutes (b) 15.0 minutes
 (c) 16.5 minutes (d) 30 minutes
8. The radioactive decay of ${}_{35}\text{X}^{88}$ by a beta emission produces an unstable nucleus which spontaneously emits a neutron. The final product is [MNR 1995; CBSE 2001]
 (a) ${}_{37}\text{X}^{88}$ (b) ${}_{35}\text{Y}^{89}$
 (c) ${}_{34}\text{Z}^{88}$ (d) ${}_{36}\text{W}^{87}$
9. What is the half-life of a radioactive substance if 75% of a given amount of the substance disintegrates in 30 minutes
 (a) 7.5 minutes (b) 25 minutes
- (c) 20 minutes (d) 15 minutes
10. In radioactive decay which one of the following moves the fastest [MP PET/PMT 1999]
 (a) α -particle (b) β -particle
 (c) γ -rays (d) Positron
11. The half-life of a radionuclide is 69.3 minutes. What is its average life (in minutes)
 (a) 100 (b) 10^{-2}
 (c) $(69.3)^{-1}$ (d) 0.693×69.3
12. 10 gm of a radioactive substance is reduced to 1.25 gm after 15 days. Its 1 kg mass will reduce (in how many days) to 500 gm in
 (a) 500 days (b) 125 days
 (c) 25 days (d) 5 days
13. A radioactive isotope having a half-life of 3 days was received after 12 days. It was found that there were 3 gm of the isotope in the container. The initial weight of the isotope when packed was [NCERT 1980; CPMT 1999; KCET 2000; Pb.CET 2001]
 (a) 12 gm (b) 24 gm
 (c) 36 gm (d) 48 gm
14. C^{14} is radioactive. The activity and the disintegration product are
 (a) β -active, ${}_{7}\text{N}^{14}$ (b) α - active, ${}_{7}\text{Be}^{10}$
 (c) Positron active, ${}_{5}\text{B}^{14}$ (d) γ - active, C^{14}
15. Radioactivity of a radioactive element remains $\frac{1}{10}$ of the original radioactivity after 2.303 seconds. The half-life period is
 (a) 2.303 (b) 0.2303
 (c) 0.693 (d) 0.0693
16. A radioactive substance has $t_{1/2}$ 60 minutes. After 3 hrs, what percentage of radioactive substance will remain [BHU 1995]
 (a) 50% (b) 75%
 (c) 25% (d) 12.5%
17. A freshly prepared radioactive source of half-life 2 hours emits radiations of intensity which is 64 times the permissible safe level. The minimum time after which it would be possible to work safely with this source is [IIT 1988]
 (a) 6 hours (b) 12 hours
 (c) 24 hours (d) 128 hours
18. During a negative β -decay [MNR 1990; IIT 1985]
 (a) An atomic electron is ejected
 (b) An electron which is already present within the nucleus is ejected
 (c) A neutron in the nucleus decays emitting an electron
 (d) A part of the binding energy of the nucleus is converted into an electron
19. The decay constant of a radioactive sample is ' λ '. The half-life and mean life of the sample are respectively [MNR 1990; IIT 1989]
 (a) $\frac{1}{\lambda}, \frac{\ln 2}{\lambda}$ (b) $\frac{\ln 2}{\lambda}, \frac{1}{\lambda}$
 (c) $\lambda \ln 2, \frac{1}{\lambda}$ (d) $\frac{\lambda}{\ln 2}, \frac{1}{\lambda}$
20. The half-life of a radio isotope is 20 hours. After 60 hours, how much amount will be left behind [MP PMT 1991]

- (a) $1/8$ (b) $1/4$
(c) $1/3$ (d) $1/2$
21. Half-life period of a zero order reaction is [AMU (Engg.) 1999]
(a) Inversely proportional to the concentration
(b) Independent of the concentration
(c) Directly proportional to the initial concentration
(d) Directly proportional to the final concentration
22. If 12 g of sample is taken, and 6 g of a sample decays in 1 hr. The amount of sample showing decay in next hour is [AMU (Engg.) 1999]
(a) 3 g (b) 1 g
(c) 2 g (d) 6 g
23. What will be half-life period of a nucleus if at the end of 4.2 days, $N = 0.798 N_0$ [MP PET 2000]
(a) 15 days (b) 10 days
(c) 12.83 days (d) 20 days
24. If 2.0 g of a radioactive substance has half-life of 7 days. The half-life of 1 g sample is [MP PET 2000]
(a) 7 days (b) 14 days
(c) 28 days (d) 35 days
25. The half-life of $^{90}_{38}\text{Sr}$ is 20 years. If its sample having initial activity of 8000 dis/min is taken, what would be its activity after 80 years
(a) 500 dis/min (b) 800 dis/min
(c) 1000 dis/min (d) 1600 dis/min
26. $^{24}_{11}\text{Na}$ half-life is 15 hours. On heating it will
(a) Reduce (b) Remain unchanged
(c) Depend on temperature (d) Become double
27. In a radioactive decay, an emitted electron comes from [CBSE 1994; Pb. PET 1999]
(a) Nucleus of the atom
(b) Inner orbital of the atom
(c) Outermost orbit of the atom
(d) Orbit having principal quantum number one
28. What is the value of decay constant of a compound having half-life time $T_{1/2} = 2.95$ days [AFMC 1997]
(a) $2.7 \times 10^{-5} \text{ s}^{-1}$ (b) $2.7 \times 10^6 \text{ s}^{-1}$
(c) $2.7 \times 10^{-6} \text{ s}^{-1}$ (d) $3 \times 10^5 \text{ s}^{-1}$
29. What kind of radioactive decay does not lead to the formation of a daughter nucleus that is an isobar of the parent nucleus
(a) α -rays (b) β -rays
(c) Positron (d) Electron capture
30. The half-life of $^{14}_6\text{C}$ if its K or λ is 2.31×10^{-4} is [BHU 1999]
(a) $2 \times 10^2 \text{ yrs}$ (b) $3 \times 10^3 \text{ yrs}$
(c) $3.5 \times 10^4 \text{ yrs}$ (d) $4 \times 10^3 \text{ yrs}$
31. A radioactive isotope has a half-life of 10 days. If today 125 mg is left over, what was its original weight 40 days earlier [KCET 2005]
(a) 2g (b) 600 mg
(c) 1 g (d) 1.5 g
32. The binding energy of $^{16}_8\text{O}$ is 127 MeV. Its binding energy per neutron is [MH CET 1999]
(a) 0.794 MeV (b) 1.5875 MeV
(c) 7.94 MeV (d) 15.875 MeV
33. If the half-life period of a first order reaction is 138.6 minutes, then the value of decay constant for the reaction will be [MH CET 1999]
(a) 5 minute (b) 0.5 minute
(c) 0.05 minute (d) 0.005 minute
34. Half-life of 10 gm of radioactive substance is 10 days. The half-life of 20 gm is [CPMT 1996]
(a) 10 days (b) 20 days
(c) 25 days (d) Infinite
35. 8 gm of the radioactive isotope, cesium-137 were collected on February 1 and kept in a sealed tube. On July 1, it was found that only 0.25 gm of it remained. So the half-life period of the isotope is [KCET 1989]
(a) 37.5 days (b) 30 days
(c) 25 days (d) 50 days
36. The half-life of radium (226) is 1620 years. The time taken to convert 10 grams of radium to 1.25 grams is [MP PET 1994; UPSEAT 2001]
(a) 810 years (b) 1620 years
(c) 3240 years (d) 4860 years
37. Half-life of a radioactive substance is 120 days. After 480 days, 4 gm will be reduced to [EAMCET 1993]
(a) 2 (b) 1
(c) 0.5 (d) 0.25
38. The half-life of $^{60}_{27}\text{Co}$ is 7 years. If one gm of it decays, the amount of the substance remaining after 28 years is [EAMCET 1992]
(a) 0.25 gm (b) 0.125 gm
(c) 0.0625 gm (d) 0.50 gm
39. A radioactive isotope decays at such a rate that after 96 minutes only $\frac{1}{8}$ th of the original amount remains. The half-life of this nuclide in minutes is [KCET 1992]
(a) 12 (b) 24
(c) 32 (d) 48
40. $\text{C}-14$ has a half-life of 5760 years. 100mg of a sample containing $\text{C}-14$ is reduced to 25mg in [Bihar CEE 1992; AMU 2002; MHCET 1999]
(a) 11520 years (b) 2880 years
(c) 1440 years (d) 17280 years
41. Half-life of a radioactive element is 100 yrs. The time in which it disintegrates to 50% of its mass, will be [MP PET 1999]
(a) 50 yrs (b) 200 yrs
(c) 100 yrs (d) 25 yrs
42. The average life period of a radioactive element is the reciprocal of its [MP PET 1995]
(a) Half-life period
(b) Disintegration constant
(c) Number of atoms present at any time
(d) Number of neutrons
43. The half-life period of a radioactive element is 30 minutes. One sixteenth of the original quantity of the element will remain unchanged after [CPMT 1983; MP PMT 1994]
(a) 60 minutes (b) 120 minutes
(c) 70 minutes (d) 75 minutes
44. For a radioactive substance with half-life period 50 years, the time for complete decay of 100 milligram of it would be [MADT Bihar 1984]
(a) 1000 years (b) 100×500 years

- (c) 500 years (d) Infinite time
45. A substance of which one gram is taken, after half-life time what fraction of it is left ? [MADT Bihar 1983]
- (a) $\frac{1}{4}$ (b) $\frac{1}{8}$
- (c) $\frac{1}{2}$ (d) $\frac{1}{32}$
46. The half-life of the radio element ${}_{83}\text{Bi}^{210}$ is 5 days. Starting with 20 g of this isotope, the amount remaining after 15 days is
- (a) 10 g (b) 5 g
- (c) 2.5 g (d) 6.66 g
47. In radioactive decay of X into Y below, ${}_Z Y^m$ is
- $${}_6 X^{14} \xrightarrow{-3\beta} {}_Z Y^m$$
- (a) ${}_6 Y^{15}$ (b) ${}_7 Y^{17}$
- (c) ${}_9 Y^{14}$ (d) ${}_8 Y^{14}$
48. 75% of the first order reaction was completed in 32 minutes. When was 50% of the reaction completed [MNR 1983; MP PET 1997; EAMCET 1998]
- (a) 24 minutes (b) 16 minutes
- (c) 8 minutes (d) 4 minutes
49. If 2.0 g of a radioactive isotope has a half-life of 20 hr, the half-life of 0.5 g of the same substance is [MP PMT 1990; MNR 1992]
- (a) 20 hr (b) 80 hr
- (c) 5 hr (d) 10 hr
50. Radioactive lead ${}_{82}\text{Pb}^{201}$ has a half-life of 8 hours. Starting from one milligram of this isotope, how much will remain after 24 hours [MP PMT 1990]
- (a) $1/2$ mg (b) $1/3$ mg
- (c) $1/8$ mg (d) $1/4$ mg
51. The half-life of ${}_{92}\text{U}^{238}$ is 4.5×10^9 years. After how many years, the amount of ${}_{92}\text{U}^{238}$ will be reduced to half of its present amount [CPMT 1990; MP PET 1999]
- (a) 9.0×10^9 years (b) 13.5×10^9 years
- (c) 4.5×10^9 years (d) $4.5 \times 10^{4.5}$ years
52. Radium has atomic weight 226 and a half-life of 1600 years. The number of disintegrations produced per second from 1 gm are [BHU 1990]
- (a) 4.8×10^{10} (b) 9.2×10^6
- (c) 3.7×10^{10} (d) Zero
53. The half-life of a radioactive element is 6 months. The time taken to reduce its original concentration to its $1/16$ value is [MP PET 1991]
- (a) 1 year (b) 16 years
- (c) 2 years (d) 8 years
54. In the case of a radio isotope the value of $T_{1/2}$ and λ are identical in magnitude. The value is [KCET 2002]
- (a) 0.693 (b) $(0.693)^{1/2}$
- (c) $1/0.693$ (d) $(0.693)^2$
55. A radioactive element has half-life of one day. After three days, the amount of the element left will be [MNR 1985; UPSEAT 2000, 01; MH CET 2002]
- (a) $1/2$ of the original amount
- (b) $1/4$ of the original amount
- (c) $1/8$ of the original amount
- (d) $1/16$ of the original amount
56. The radioactivity due to C^{14} isotope (half-life 6000 years) of a sample of wood from an ancient tomb was found to be nearly half that of fresh wood, the tomb is therefore about [NCERT 1980, 81; MP PET 1989]
- (a) 3000 years old (b) 6000 years old
- (c) 9000 years old (d) 1200 years old
57. The decay of a radioactive element follows first order kinetics, as a result [NCERT 1982]
- (a) Half-life period = constant / k , where k is the decay constant
- (b) Rate of decay is independent of temperature
- (c) Rate can be changed by changing chemical conditions
- (d) The element will be completely transformed into a new element after expiry of two half-life period
58. Half-life of a radioactive substance which disintegrates by 75 % in 60 minutes, will be [MP PMT 2002]
- (a) 120 min (b) 30 min
- (c) 45 min (d) 20 min
59. 87.5% decomposition of a radioactive substance complete in 3 hours. What is the half-life of that substance [MP PMT 2003]
- (a) 2 hours (b) 3 hours
- (c) 90 minutes (d) 1 hours
60. Tritium undergoes radioactive decay giving [CPMT 1976; NCERT 1978]
- (a) α -particles (b) β -particles
- (c) Neutrons (d) None of these
61. Given that a radioactive species decays according to exponential law $N = N_0 e^{-\lambda t}$. The half-life of the species is [Kerala (Med.) 2003]
- (a) λ (b) No
- (c) $\lambda/\ln 2$ (d) $\ln 2/\lambda$
62. Half-life of a radioactive disintegration ($A \rightarrow B$) having rate constant 231 sec^{-1} is [CPMT 1988]
- (a) $3.0 \times 10^{-2} \text{ sec}$ (b) $3.0 \times 10^{-3} \text{ sec}$
- (c) $3.3 \times 10^{-2} \text{ sec}$ (d) $3.3 \times 10^{-3} \text{ sec}$
63. The amount of ${}_{53}\text{I}^{128}$ ($t_{1/2} = 25$ minutes) left after 50 minutes will be [AIIMS 1982; DPMT 1982, 83]
- (a) One – half (b) One – third
- (c) One – fourth (d) Nothing
64. If $3/4$ quantity of a radioactive element disintegrates in two hours, its half-life would be [MP PMT 1989; CPMT 1984]
- (a) 1 hour (b) 45 minutes
- (c) 30 minutes (d) 15 minutes
65. Radioactive decay is a [MP PMT 1989, 97]
- (a) Second order reaction (b) First order reaction
- (c) Zero order reaction (d) Third order reaction
66. The half-life of a radioactive element depends upon [EAMCET 1980]
- (a) The amount of the element
- (b) The temperature
- (c) The pressure
- (d) None of these
67. The activity of radio isotope changes with [MNR 1986]
- (a) Temperature (b) Pressure
- (c) Chemical environment (d) None of these
68. A certain nuclide has a half-life of 25 minutes. If one starts with 100 g of it, how much of it will remain at the end of 100 minutes

- (a) 1.0 g (b) 4.0 g
(c) 6.25 g (d) 12.50 g
69. If U^{235} is bombarded with neutrons, atom will split into [CPMT 1981]
(a) $Sr + Pb$ (b) $Cs + Rb$
(c) $Kr + Cd$ (d) $Ba + Kr$
70. If 8.0 g of a radioactive isotope has a half-life of 10 hrs. The half-life of 2.0 g of the same substance is [UPSEAT 2001]
(a) 2.5 hrs. (b) 5 hrs.
(c) 10 hrs. (d) 40 hrs.
71. If the disintegration constant is 6.93×10^{-6} , then half-life of ${}_6C^{14}$ will be [KCET 2001]
(a) 10^2 yrs (b) 10^3 yrs
(c) 10^4 yrs (d) 10^5 yrs
72. The decay constant of Ra^{226} is $1.37 \times 10^{-11} \text{ sec}^{-1}$. A sample of Ra^{226} having an activity of 1.5 millicurie will contain atoms
(a) 4.1×10^{18} (b) 3.7×10^{17}
(c) 2.05×10^{15} (d) 4.7×10^{10}
73. Amount of ${}_{53}I^{128}$ ($t_{1/2} = 25 \text{ min}$) left after 75 minutes is [DCE 2002]
(a) 1/6 (b) 1/4
(c) 1/8 (d) 1/9
74. The half-life of a radioisotope is four hours. If the initial mass of the isotope was 200 g, the mass remaining after 24 hours undecayed is
(a) 3.125 g (b) 2.084 g
(c) 1.042 g (d) 4.167 g
75. An artificial radioactive isotope gave ${}_{7}^{14}N$ after two successive β -particle emissions. The number of neutrons in the parent nucleus must be [KCET 2004]
(a) 9 (b) 14
(c) 5 (d) 7
76. If the half-life of an isotope X is 10 years, its decay constant is [DCE 2004]
(a) 6.932 yr^{-1} (b) 0.6932 yr^{-1}
(c) 0.06932 yr^{-1} (d) 0.006932 yr^{-1}
77. A radioactive isotope decays at such a rate that after 192 minutes only 1/16 of the original amount remains. The half-life of the radioactive isotope is [Kerala CET 2004]
(a) 32 min (b) 48 min
(c) 12 min (d) 24 min
78. In the given reaction, ${}_{92}U^{235} \xrightarrow{-\alpha} (A) \xrightarrow{-\beta} (B) \xrightarrow{-\beta} (C)$ isotope are [Pb. CET 2000]
(a) A and C (b) ${}_{92}U^{235}$ and C
(c) A and B (d) A , B and C
79. Rate constant for a reaction is λ . Average life is representative by
(a) $1/\lambda$ (b) $\ln 2/\lambda$
(c) $\frac{\lambda}{\sqrt{2}}$ (d) $\frac{0.693}{\lambda}$
80. For a reaction, the rate constant is 2.34 sec^{-1} . The half-life period for the reaction is
(a) 0.30 sec (b) 0.60 sec
(c) 3.3 sec (d) Data is insufficient
81. $T_{1/2}$ of C^{14} isotope is 5770 years. time after which 72% of isotope left is [Orissa JEE 2005]
(a) 2740 years (b) 274 years
(c) 2780 years (d) 278 years
82. A radioactive substance takes 20 min to decay 25%. How much time will be taken to decay 75% [Orissa JEE 2005]
(a) 96.4 min (b) 68 min
(c) 964 min (d) 680 min
83. A radioactive sample is emitting 64 times radiations than non hazardous limit. if its half life is 2 hours, after what time it becomes non-hazardous [DPMT 2005]
(a) 16 hr (b) 12 hr
(c) 8 hr (d) 4 hr
84. If 8.0 g of a radioactive substance has a half-life of 10 hrs., the half life of 2.0 g of the same substance is [J & K 2005]
(a) 2.6 hr (b) 5 hr
(c) 10 hr (d) 40 hr

Artificial transmutation

1. The age of most ancient geological formation is estimated by [NCERT 1981; MP PET/PMT 1988; CBSE 1989; MP PET 1997; MP PMT 2002]
(a) Potassium – Argon method
(b) Carbon – 14 dating method
(c) Radioactive carbon method
(d) Uranium – Lead method
2. The equation ${}_3Li^6 + {}_1H^2 \longrightarrow {}_2He^4 + \text{energy}$ represents
(a) Synthesis of helium (b) Transmutation of element
(c) Fusion reaction (d) Nuclear fission
3. The phenomenon of radioactivity arises from the [Kerala (Med.) 2002]
(a) Binary fission
(b) Nuclear fusion
(c) Stable nuclei
(d) Decay of unstable nuclei
4. The first artificial disintegration of an atomic nucleus was achieved by [Kerala (Engg.) 2002]
(a) Geiger (b) Wilson
(c) Madame curie (d) Rutherford
(e) Soddy
5. Artificial elements have been prepared by bombardment reactions in high energy accelerators. What is the mass number of the element X produced in the following nuclear reaction
 ${}_{95}^{249}Cf + {}_{7}^{15}N \rightarrow {}_{105}X + {}_{0}^1n$ [AMU (Engg.) 2002]
(a) 261 (b) 264
(c) 260 (d) 257
6. Radioactive carbon dating was discovered by [MP PET 2001]
(a) W.F. Libby (b) G.N. Lewis
(c) J. Willard Gibbs (d) W. Nernst
7. The nuclear reaction
 ${}_{29}^{63}Cu + {}_2^4He \rightarrow {}_{17}^{37}Cl + 14{}_1^1H + 16{}_0^1n$ is referred to as [MP PET 2002]
(a) Spallation reaction (b) Fusion reaction
(c) Fission reaction (d) Chain reaction

8. The carbon dating is based on [MP PMT 2001]
 (a) ${}^{15}_6\text{C}$ (b) ${}^{14}_6\text{C}$
 (c) ${}^{13}_6\text{C}$ (d) ${}^{11}_6\text{C}$
9. A possible material for use in the nuclear reactors as a fuel is [DPMT 1986]
 (a) Thorium (b) Zirconium
 (c) Beryllium (d) Plutonium
10. Heavy water freezes at [UPSEAT 2001]
 (a) 0°C (b) 3.8°C
 (c) 38°C (d) -0.38°C
11. To determine the masses of the isotopes of an element which of the following techniques is useful [NCERT 1978; MNR 1979]
 (a) The acceleration of charged atoms by an electric field and their subsequent deflection by a variable magnetic field
 (b) The spectroscopic examination of the light emitted by vaporised elements subjected to electric discharge
 (c) The photographing of the diffraction patterns which arise when X-rays are passed through crystals
 (d) The bombardment of metal foil with alpha particles
12. The radioisotope, tritium (${}^3_1\text{H}$) has a half-life of 12.3 years. If the initial amount of tritium is 32 mg. How many milligrams of it would remain after 49.2 years [CBSE 2003]
 (a) 8 mg (b) 1 mg
 (c) 2 mg (d) 4 mg
13. Neutron is used as a [CPMT 1988]
 (a) Reducing agent (b) Moderator
 (c) Tracer (d) In biological programme
14. Hydrogen bomb is based on the phenomenon of [EAMCET 1980; CPMT 1984, 96; MP PMT 1993, 95, 2002; RPET 1999]
 (a) Nuclear fission (b) Nuclear fusion
 (c) Nuclear explosion (d) Disintegration
15. In the nuclear reactors the speed of the neutrons is slowed down by
 (a) Heavy water (b) Ordinary water
 (c) Zinc rods (d) Molten caustic soda
16. By which law, energy produced in nuclear reaction is given [MP PET 2000]
 (a) Graham's law (b) Charle's law
 (c) Gas Lussac's Law (d) Einstein's law
17. If two light nuclei are fused together in nuclear reaction the average energy per nucleon [Pb. PMT 2001]
 (a) Increases (b) Cannot be determined
 (c) Remains same (d) Decreases
18. A wood piece is 11460 years old. What is the fraction of ${}^{14}_6\text{C}$ activity left in the piece? (Half-life period of ${}^{14}_6\text{C}$ is 5730 years) [MP PMT 2000]
 (a) 0.12 (b) 0.25
 (c) 0.50 (d) 0.75
19. When nuclear energy is intended to be harnessed for generation of electricity, potentially destructive neutron released in a nuclear reactor are absorbed by [MH CET 2001]
 (a) Long rods of Cd (b) Heavy water
 (c) Cubical blocks of steel (d) Both (a) and (c)
20. The proper rays for radiocarbon dating are [MP PET 2002]
 (a) UV-rays (b) IR-rays
 (c) Cosmic rays (d) X-rays
21. ${}_1\text{H}^2 + {}_1\text{H}^2 \rightarrow {}_2\text{He}^3 + {}_0\text{n}^1$. Above nuclear reaction is called [UPSEAT 2001]
 (a) Nuclear fission
 (b) Nuclear fusion
 (c) Artificial transmutation
 (d) Spontaneous disintegration
22. Which of the following is used as a moderator in a nuclear reactor
 (a) D_2O (b) N_2O
 (c) H_2O (d) NaOH
23. The fuel of atomic pile is [NCERT 1973; AFMC 1989]
 (a) Thorium (b) Sodium
 (c) Uranium (d) Petroleum
24. Atom bomb is based on the principle of [CPMT 1982; BHU 1985]
 (a) Nuclear fusion
 (b) Nuclear fission
 (c) Radioactivity
 (d) Fusion and fission both
25. Who observed that when the nucleus of uranium atom was bombarded with fast moving neutrons, it becomes so very unstable that it is immediately broken into two nuclei of nearly equal mass besides other fragments
 (a) J.J. Thomson (b) Chadwick
 (c) Einstein (d) Hahn and Strassmann
26. When a radioactive substance is subjected to vacuum, the rate of disintegration per second [DPMT 1985; NCERT 1972]
 (a) Increases considerably
 (b) Increases only if the products are gaseous
 (c) Is not affected
 (d) Suffers a slight decrease
27. A radio isotope will not emit [KCET 2002]
 (a) Gamma and alpha rays simultaneously
 (b) Gamma rays only
 (c) Alpha and beta rays simultaneously
 (d) Beta and gamma rays simultaneously
28. What is the packing fraction of ${}^{56}_{26}\text{Fe}$ [JIPMER 2002]
 (Isotopic mass = 55.92066)
 (a) -14.167 (b) 173.90
 (c) -14.187 (d) -73.90
29. The energy released in an atom bomb explosion is mainly due to
 (a) Release of neutrons
 (b) Release of electrons
 (c) Greater mass of products than initial material
 (d) Lesser mass of products than initial material
30. C^{14} is [KCET 2002]
 (a) A natural radioactive isotope
 (b) A natural non-radioactive isotope
 (c) An artificial radioactive isotope
 (d) An artificial non-radioactive isotope
31. A radioactive isotope has a half-life of 10 years. What percentage of the original amount of it remain after 20 years [KCET 2001]
 (a) 0 (b) 12.5
 (c) 8 (d) 25
32. In a chain reaction, uranium atom gets fissioned forming two different materials. The total weight of these put together is
 (a) More than the weight of parent uranium atom

- (b) Less than the weight of parent uranium atom
(c) More or less depends upon experimental conditions
(d) Neither more nor less
33. A substance used as a moderator in nuclear reactors is [MP PET 1995]
(a) Cadmium (b) Uranium-235
(c) Lead (d) Heavy water
34. Equation ${}_{17}\text{Cl}^{37} + {}_1\text{H}^2 \rightarrow {}_{18}\text{Ar}^{38} + {}_0\text{n}^1$ is [MP PMT 1989]
(a) Nuclear fission
(b) Nuclear fusion
(c) Transformation of chlorine
(d) Synthesis of argon
35. 1.0 gm radioactive sodium on decay becomes 0.25 gm in 16 hours. How much time 48 gm of same radioactive sodium will need to become 3.0 gm
(a) 48 hours (b) 32 hours
(c) 20 hours (d) 16 hours
36. Large energy released in an atomic bomb explosion is mainly due to
(a) Products having a lesser mass than initial substance
(b) Conversion of heavier to lighter atoms
(c) Release of neutrons
(d) Release of electrons
37. The reaction ${}_1\text{H}^2 + {}_1\text{H}^3 \rightarrow {}_2\text{He}^4 + {}_0\text{n}^1 + \text{energy}$ represents [MP PMT 1990; CPMT 1990; KCET 1992]
(a) Nuclear fission
(b) Nuclear fusion
(c) Artificial disintegration
(d) Transmutation of element
38. Carbon-14 dating method is based on the fact that [CBSE 1997]
(a) Carbon-14 fraction is the same in all objects
(b) Carbon-14 is highly insoluble
(c) Ratio of carbon-14 and carbon-12 is constant
(d) All of these
39. Half-life period of a radioactive element is 10.6 yrs. How much time will it take in its 99% decomposition [RPET 1999]
(a) 7046 yrs (b) 7.046 yrs
(c) 704.6 yrs (d) 70.4 yrs
40. Deuterium resembles hydrogen in chemical properties but reacts
(a) More vigorously than hydrogen
(b) Faster than hydrogen
(c) Slower than hydrogen
(d) Just as hydrogen
41. Which of the following is heavy water [AFMC 1997]
(a) H_2O_{18} (b) H_2O_{16}
(c) H_2O_3 (d) D_2O
42. D_2O is used in [CPMT 1997]
(a) Industry (b) Nuclear reactor
(c) Medicine (d) Insecticide
43. India conducted an underground nuclear test at [KCET 1998]
(a) Tarapur (b) Narora
(c) Pokhran (d) Pushkar
44. Energy required to separate neutron and proton from the nucleus is called [RPMT 1999]
(a) Bond energy (b) Nuclear energy
(c) Chemical energy (d) Radiation energy
45. Liquid sodium finds use in nuclear reactors. Its function is
(a) To collect the reaction products
(b) To act as a heat exchanger or coolant
(c) To absorb the neutrons in order to control the chain reaction
(d) To act as a moderator which slows down the neutrons
46. Which is least effective for artificial transmutation [DPMT 2000]
(a) Deuterons (b) Neutrons
(c) α -particles (d) Protons
47. A piece of wood was found to have $\text{C}^{14}/\text{C}^{12}$ ratio 0.7 times that in a living plant. The time period when the plant died is (Half-life of $\text{C}^{14} = 5760 \text{ yrs}$) [Pb. PMT 1999]
(a) 2770 yrs (b) 2966 yrs
(c) 2980 yrs (d) 3070 yrs
48. When a slow neutron goes sufficiently close to a U^{235} nucleus, then the process which takes place is [AFMC 2000]
(a) Fusion of U^{235} (b) Fission of U^{235}
(c) Emission of α -particle (d) First (a) then (b)
49. ${}_{13}\text{Al}^{28}$ when radiated by suitable projectile gives ${}_{15}\text{P}^{31}$ and neutron. The projectile used is [MP PMT/PET 1988; CPMT 1985, 82]
(a) Proton (b) Neutron
(c) Alpha particle (d) Deuteron
50. Which of the following statements about radioactivity of an element is incorrect
(a) It is a nuclear property
(b) It does not involve any rearrangement of electrons
(c) Its rate is affected by change in temperature and/or pressure
(d) It remains unaffected by the presence of other element or elements chemically combined with it
51. Radioactive iodine is being used to diagnose the disease of [MP PET 1996]
(a) Bones (b) Kidneys
(c) Blood cancer (d) Thyroid
52. C-14 is used in carbon dating of dead objects because [DPMT 1996]
(a) Its half-life is 10^3 years
(b) Its half-life is 10^4 years
(c) It is found in nature abundantly and in definite ratio
(d) It is found in dead animals abundantly
53. A radioactive element resembling iodine in properties is [Kurukshetra CEE 1998]
(a) Astatine (b) Lead
(c) Radium (d) Thorium
54. For artificial transmutation of nuclei, the most effective one is [MP PMT 1996]
(a) Proton (b) Deuteron
(c) Helium nuclei (d) Neutron
55. Which of the following cannot be accelerated [KCET 2005]
(a) α -particle (b) β -particle
(c) Protons (d) Neutrons
56. For the fission reaction
 ${}_{92}\text{U}^{235} + {}_0\text{n}^1 \rightarrow {}_{56}\text{Ba}^{140} + {}_y\text{E}^x + 2{}_0\text{n}^1$
The value of x and y will be
(a) $x = 93$ and $y = 34$ (b) $x = 92$ and $y = 35$
(c) $x = 89$ and $y = 44$ (d) $x = 94$ and $y = 36$
57. Heavy water is used as

- [Bihar MEE 1996; UPSEAT 1999, 2000, 02]
- (a) Control rods (b) Moderator
(c) Fuel (d) Coolant
(e) None of these
58. Unit for radioactive constant is [MP PET 1990]
(a) Time^{-1} (b) Time
(c) $\text{Mole} - \text{time}^{-1}$ (d) $\text{Time} - \text{mole}^{-1}$
59. Which of the following is used in dating archeological findings or In a method of absolute dating of fossils a radioactive element is used. It is [CPMT 1983, 85; NCERT 1978; BHU 1981; MP PMT 1993; AFMC 1997]
(a) ${}_{92}\text{U}^{235}$ (b) ${}_6\text{C}^{14}$
(c) ${}_6\text{C}^{12}$ (d) ${}_{20}\text{Ca}^{40}$
60. A radioactive isotope has a half-life of 20 days. If 100 gm of the substance is taken, the weight of the isotope remaining after 40 days is [NCERT 1979]
(a) 25 gm (b) 2.5 gm
(c) 60 gm (d) 40 gm
61. In a fission reaction the nucleus of an element [NCERT 1977]
(a) Loses only some elementary nuclear particles from another nucleus
(b) Captures some elementary nuclear particles from another nucleus
(c) Breaks up into several smaller nuclei
(d) Breaks up into two smaller nuclei with the loss of same elementary nuclear particles
62. The huge amount of energy which is released during atomic fission is due to [CPMT 1990]
(a) Loss of mass (b) Loss of electrons
(c) Loss of protons (d) Loss of α -particles
63. The measure of binding energy of a nucleus is the [CPMT 1982; Kurukshetra CEE 1998]
(a) Mass defect (b) Energy of protons
(c) Energy of neutrons (d) Total energy of nucleons
64. The first controlled artificial disintegration of an atomic nucleus was achieved by [BHU 1987]
(a) Geiger (b) Wilson
(c) Cockcroft (d) Rutherford
65. Artificial radioactivity was first discovered by [CPMT 1972; BHU 1984; KCET 1999]
(a) Seaberg (b) Rutherford
(c) Einstein (d) Irene Curie & Juliot
66. The half-life period of a radioactive element is 140 days. After 560 days, one gram of the element will reduce to [CPMT 1989; IIT 1986; EAMCET 1992; MP PET 1997; UPSEAT 1999]
(a) $1/2$ g (b) $1/4$ g
(c) $1/8$ g (d) $1/16$ g
67. A device used for the measurement of radioactivity is [BHU 1979]
(a) Mass spectrometer (b) Cyclotron
(c) Nuclear reactor (d) G.M. counter
68. In a nuclear reactor, chain reaction is controlled by introducing
(a) Iron rod (b) Cadmium rod
(c) Graphite rod (d) Platinum rod
69. In atomic reactors, graphite is used as a [NCERT 1980; MP PET 1989]
(a) Lubricant
(b) Moderator to slow down neutrons
(c) Fuel
(d) Liner of the reactor
70. The modern basis of atomic weight is [MP PET 1989; CPMT 1993]
(a) Isotope $H^1 = 1.000$ (b) Oxygen = 16.000
(c) Isotope $O^{16} = 16.000$ (d) Isotope $C^{12} = 12.000$
71. Which radioactive carbon has been helpful in understanding the mechanism of photosynthesis in plants
(a) ${}_6\text{C}^{14}$ (b) ${}_6\text{C}^{13}$
(c) ${}_6\text{C}^{12}$ (d) ${}_6\text{C}^{15}$
72. Artificial transmutation was discovered by [Pb.CET 2003]
(a) Pauli (b) Rutherford
(c) Soddy (d) Curie
73. Which of the following is an example of nuclear fusion [MP PMT 1989; DCE 2004]
(a) ${}_1\text{H}^2 + {}_1\text{H}^2 \rightarrow {}_2\text{He}^4 + \text{energy}$
(b) ${}_{92}\text{U}^{235} + {}_0n^1 \rightarrow {}_{56}\text{Ba}^{141} + {}_{36}\text{Kr}^{92} + {}_{30}\text{n}^1 + \text{energy}$
(c) ${}_{13}\text{Al}^{27} + {}_1\text{H}^1 \rightarrow {}_{12}\text{Mg}^{24} + {}_2\text{He}^4$
(d) None of these
74. The radioactivity isotope ${}_{27}^{60}\text{Co}$ which is used in the treatment of cancer can be made by (n, p) reaction. For this reaction the target nucleus is [CBSE PMT 2004]
(a) ${}_{28}^{60}\text{Ni}$ (b) ${}_{27}^{60}\text{Co}$
(c) ${}_{28}^{59}\text{Ni}$ (d) ${}_{27}^{59}\text{Co}$
75. Fusion bomb involves [AFMC 2004]
(a) Combination of lighter nuclei into bigger nucleus
(b) Destruction of heavy nucleus into smaller nuclei
(c) Combustion of oxygen
(d) Explosion of TNT
76. The element used for dating the ancient remains is [AFMC 2004]
(a) Ni (b) C-14
(c) C-12 (d) Rd
77. If radium and chlorine combine to form radium chloride the compound is [Kerala PMT 2004]
(a) No longer radioactive
(b) Twice as radioactive as radium
(c) Half as radioactive as radium
(d) As radioactive as radium
(e) Thrice as radioactive as radium
78. Which of the following is an example of nuclear fission [Pb. CET 2002]
(a) ${}_1\text{H}^2 + {}_1\text{H}^2 \rightarrow {}_2\text{He}^4 + \gamma$
(b) $A + B \rightarrow C + \text{energy}$ [EAMCET 1984]
(c) ${}_{92}\text{U}^{235} + {}_0n^1 \rightarrow {}_{56}\text{Ba}^{141} + {}_{36}\text{Kr}^{92} + 3 {}_0n^1 + \text{energy}$
(d) ${}_{13}\text{Al}^{27} + {}_2\text{He}^4 \rightarrow {}_{15}\text{P}^{30} + {}_0n^1$

79. The C^{14} to C^{12} ratio in a wooden article is 13% that of the fresh wood. Calculate the age of the wooden article. Given that the half-life of C^{14} is 5770 years [Pb.CET 2004]
- (a) 16989 years (b) 16858 years
(c) 15675 years (d) 17700 years
80. Hydrogen bomb is based on the principle of [AIEEE 2005]
- (a) Nuclear fission (b) Natural radioactivity
(c) Nuclear fusion (d) Artificial radioactivity
81. Match List –I and List-II and choose right one by using code given in list [Kerala CET 2005]
- | List – I | List –II |
|-----------------|----------------|
| Nuclear reactor | Used substance |
| Component | |
| 1. Moderator | (A) Uranium |
| 2. Control rods | (B) Graphite |
| 3. Fuel rods | (C) Boron |
| 4. Coolant | (D) Lead |
| | (E) Sodium |
- Code :
- | | 1 | 2 | 3 | 4 |
|-----|---|---|---|---|
| (a) | B | A | C | E |
| (b) | B | C | A | E |
| (c) | C | B | A | E |
| (d) | C | D | A | B |
| (e) | D | C | B | A |

Isotopes-Isotones and Nuclear isomers

1. Substances which have identical chemical properties but differ in atomic weights are called [EAMCET 1980, 83; DPMT 1985; MNR 1982]
- (a) Isothermals (b) Isotopes
(c) Isentropus (d) Elementary particles
2. Tritium is an isotope of [DPMT 1985]
- (a) Hydrogen (b) Titanium
(c) Tantalum (d) Tellurium
3. O – 18 isotope of oxygen will have [CPMT 1972, 79]
- (a) 18 protons
(b) 9 protons and 9 neutrons
(c) 8 neutrons and 10 protons
(d) 10 neutrons and 8 protons
4. Which of the following is an isobaric pair [CPMT 1987, 93]
- (a) ${}_6C^{13}, {}_7N^{13}$ (b) ${}_6C^{13}, {}_7N^{14}$
(c) ${}_7N^{14}, {}_8O^{15}$ (d) ${}_7N^{13}, {}_8O^{15}$
5. Isotopes are atoms having the same [EAMCET 1978, 79; MP PMT 1980; CPMT 1973; BHU 2001; AFMC 2003]
- (a) Atomic mass (b) Mass number
(c) Atomic number (d) Number of neutrons
6. Successive emission of an α -particle and two β -particles by an atom of an element results in the formation of its [MP PMT/PET 1988; BHU 1979]
- (a) Isobar (b) Isomorph
(c) Isotope (d) Isomer
7. In treatment of cancer, which of the following isotope is used [DPMT 1985; BHU 1995; KCET 1999; AMU 1999; Pb.CET 2001; MP PET 2002; Kurukshetra CET 2002]
- (a) ${}_{53}I^{131}$ (b) ${}_{15}P^{32}$
(c) ${}_{27}Co^{60}$ (d) ${}_1H^2$
8. Elements having different nuclear charge but the same mass number are called [NCERT 1974; MP PMT 1991; CBSE PMT 1991; CPMT 1989; EAMCET 1992]
- (a) Isotopes (b) Isobars
(c) Isomers (d) Isotones
9. Which isotope on bombardment with α -particles will give ${}_8O^{17}$ and ${}_1H^1$ [NCERT 1983]
- (a) ${}_8O^{16}$ (b) ${}_7N^{14}$
(c) ${}_7N^{15}$ (d) ${}_6C^{14}$
10. Emission of β -particle by an atom of an element results in the formation of its [BHU 1979; DPMT 1985; KCET 1999]
- (a) Isotope (b) Isomer
(c) Isomorph (d) Isobar
11. Radioactive isotopes that have an excessive neutron/proton ratio generally exhibit
- (a) e^- emission (b) ${}_2He^4$ emission
(c) e^+ emission (d) K -electron capture
12. Atomic weights of carbon, nitrogen and oxygen are 12, 14 and 16 respectively. An atom of atomic weight 14 and nuclear charge + 6 is an isotope of
- (a) Oxygen (b) Carbon
(c) Nitrogen (d) None of these
13. Isotopes of an element have [MNR 1985]
- (a) Similar chemical properties but different physical properties
(b) Similar chemical and physical properties
(c) Similar physical properties but different chemical properties
(d) Different chemical and physical properties
14. Whose number is common in isotopes [AIIMS 1988]
- (a) Proton (b) Neutron
(c) Proton and neutron (d) Nucleon
15. In the following radioactive transformation $R \xrightarrow{\alpha} X \xrightarrow{\beta} Y \xrightarrow{\beta} Z$; the nuclei R and Z are [BHU 1987]
- (a) Isotopes (b) Isobars
(c) Isomers (d) Isotones
16. Which one of the following pairs represents isobars [CPMT 1988]
- (a) ${}_2He^3$ and ${}_2He^4$
(b) ${}_{12}Mg^{24}$ and ${}_{12}Mg^{25}$
(c) ${}_{19}K^{40}$ and ${}_{20}Ca^{40}$
(d) ${}_{19}K^{39}$ and ${}_{19}K^{40}$
17. Nuclei of isotopes differ in [CPMT 1986, 90; MP PMT 1987]
- (a) The number of protons
(b) The number of neutrons
(c) The number of protons and neutrons both
(d) None of these
18. An isotope of 'parent' is produced, when its nucleus loses [CPMT 1987; MP PET 1991]
- (a) One α -particle
(b) One β -particle

- (c) One α and two β -particles
(d) One β and two α - particles
19. Which of the following isotopes is likely to be most stable [EAMCET 1982]
(a) ${}_{30}\text{Zn}^{71}$ (b) ${}_{30}\text{Zn}^{66}$
(c) ${}_{30}\text{Zn}^{64}$ (d) None of these
20. Which of the following statement is false [Manipal MEE 1995]
(a) In chlorine gas, the ratio of Cl^{35} and Cl^{37} is 1 : 3
(b) The hydrogen bomb is based on the principle of nuclear fusion
(c) The atom bomb is based on the principle of nuclear fission
(d) The penetrating power of a proton is less than that of an electron
21. Isotones are elements having [Bihar MEE 1996; Bihar CEE 1995]
(a) Same mass number but different neutrons
(b) Same atomic number but different neutrons
(c) Same atomic number, mass number and neutrons
(d) Different atomic and mass number but same neutrons
22. Isobaric atoms may contain
(a) Same number of p^+ and different number of n^0
(b) Same number of n^0 and different number of p^+
(c) Same number of both p^+ and n^0
(d) Different numbers of both p^+ and n^0
23. ${}_{20}\text{X}^{40}$ and ${}_{21}\text{X}^{40}$ are [CPMT 1996]
(a) Isobars (b) Isotopes
(c) Isotones (d) Isostereomers
24. Which property is different for neutral atoms of the two isotopes of the same element [JIPMER 2001]
(a) Number of protons (b) Atomic number
(c) Number of neutrons (d) None of these
25. Which of the following species is isotonic with ${}_{37}\text{Rb}^{86}$ [BHU 2001]
(a) ${}_{36}\text{Kr}^{84}$ (b) ${}_{37}\text{Rb}^{85}$
(c) ${}_{38}\text{Sr}^{87}$ (d) ${}_{39}\text{Y}^{89}$
26. The maximum sum of the number of neutrons and protons in an isotope of hydrogen is [Pb. PMT 2001]
(a) 4 (b) 5
(c) 6 (d) 3
27. Difference in ${}_{17}\text{Cl}^{35}$ and ${}_{17}\text{Cl}^{37}$ is of [AFMC 2000]
(a) Atomic number (b) Number of protons
(c) Number of neutrons (d) Number of electrons
28. Which of the following is an isotonic pair [AMU (Engg.) 2000]
(a) ${}_{19}^{40}\text{K}$, ${}_{20}^{40}\text{Ca}$ (b) ${}_{19}^{39}\text{K}$, ${}_{20}^{40}\text{Ca}$
(c) ${}_{18}^{33}\text{Ar}$, ${}_{18}^{40}\text{Ar}$ (d) ${}_{18}^{40}\text{Ar}$, ${}_{20}^{40}\text{Ca}$
29. ${}_{6}\text{C}^{11}$ and ${}_{5}\text{B}^{11}$ are referred as [NCERT 1978]
(a) Nuclear isomers (b) Isobars
(c) Isotopes (d) Fission products
30. The atomic number of bromine is 35 and its atomic weight is 79. Two isotopes of bromine are present in equal amounts. Which of the following statements represents the correct number of neutrons [NCERT 1983]
(a) 44 (b) 45
(c) 46 (d) 47
31. Isotopes are those which contain [RPMT 1997]
(a) Same number of neutrons
(b) Same physical properties
(c) Same chemical properties
(d) Different atomic mass
32. An element 'A' emits an α -particle and forms 'B'. 'A' and 'B' are [DPMT 1990]
(a) Isotopes (b) Isobars
(c) Isotones (d) Isodiasphere
33. Which of the following properties are different for neutral atoms of isotopes of the same element [EAMCET 1987; NCERT 1971; CPMT 1976; MP PET 1994]
(a) Mass
(b) Atomic number
(c) General chemical reactions
(d) Number of electrons
34. The isotope ${}_{92}\text{U}^{235}$ decays in a number of steps to an isotope of lead ${}_{82}\text{Pb}^{207}$. The groups of particles emitted in this process will be [MP PMT 1987]
(a) $4\alpha, 7\beta$ (b) $6\alpha, 4\beta$
(c) $7\alpha, 4\beta$ (d) $10\alpha, 8\beta$
35. Addition of two neutrons in an atom A would [AMU 1984]
(a) Change the chemical nature of A
(b) Produce an isobar of A
(c) Produce an isotope of A
(d) Produce another element
36. Atomic weight of the isotope of hydrogen which contains 2 neutrons in the nucleus would be [CPMT 1980]
(a) 2 (b) 3
(c) 1 (d) 4
37. If a radioactive isotope with atomic number A and mass number M emits an α -particle, the atomic number and mass number of that new isotope will become [NCERT 1980]
(a) $A - 2$, $M - 4$ (b) $A - 2$, M
(c) A, $M - 2$ (d) $A - 4$, $M - 2$
38. Which character is different of the two isotopes of an element [NCERT 1971; EAMCET 1987]
(a) Atomic mass (b) Atomic number
(c) Number of electrons (d) Number of protons
39. The symbol of an isotope is ${}_{32}\text{X}^{65}$, this reveals that [MP PET 1991]
(a) Its atomic number is 32 and atomic weight is 65
(b) Its atomic number is 65
(c) It has 65 electrons
(d) It has 32 neutrons
40. Two atoms have the same atomic mass but different atomic numbers. Such atoms are called as [NCERT 1971, 76; IIT 1983]
(a) Isotopes (b) Isobars
(c) Isomer (d) Isoelectronic
- ${}_{18}\text{Ar}^{40}$, ${}_{20}\text{Ca}^{40}$ and ${}_{19}\text{K}^{40}$ are

[MNR 1983; DPMT 1991; EAMCET 1992;
RPMT 1997; Pb.CET 2000]

- (a) Isomers (b) Isotopes
(c) Isobars (d) Isotones
42. Atoms in hydrogen gas have preponderance of [CPMT 1972]
(a) ${}_1H^1$ atoms
(b) Deuteron atoms
(c) Tritium atoms
(d) All the three (a), (b) and (c) are in equal proportion
43. Positron emission results from the transformation of one nuclear proton into neutron. The isotope thus produced possesses
(a) Same mass number (b) Higher nuclear charge
(c) Intense radioactivity (d) No radioactivity
44. An isotope of oxygen has mass number 18. Other isotopes of oxygen will have the same [MP PMT 1985; MADT Bihar 1981]
(a) Mass number (b) Atomic weight
(c) Number of neutrons (d) Number of protons
45. Two nuclei which are not identical but have the same number of nucleons represent
(a) Isotopes (b) Isobars
(c) Isotones (d) None of the three
46. The β -decay of ${}_{11}Na^{24}$ produces an isotope of [NCERT 1978]
(a) *Mg* (b) *Na*
(c) *Al* (d) *Ne*
47. Isotopes differ in [NCERT 1973]
(a) Number of protons (b) Valency
(c) Chemical reactivity (d) Number of neutrons
48. The isobars are atoms with the same number of [DPMT 1982; CPMT 1994]
(a) Protons (b) Neutrons
(c) Protons and neutrons (d) Nucleons
49. Radioactive isotope of hydrogen is [MP PMT 2001; MPPET 2003]
(a) Tritium (b) Deuterium
(c) Para hydrogen (d) Ortho hydrogen
50. Isotopes of same elements have the same number of [BHU 1984; DPMT 1983; CPMT 1972, 78; AFMC 2000, 01]
(a) Protons (b) Neutrons
(c) Deutrons (d) None
51. In chlorine gas, ratio of Cl^{35} and Cl^{37} is [BHU 1984; CPMT 1977, 80]
(a) 1 : 3 (b) 3 : 1
(c) 1 : 1 (d) 1 : 4
52. An ordinary oxygen contains [NCERT 1977]
(a) Only $O-16$ isotopes
(b) Only $O-17$ isotopes
(c) A mixture of $O-16$ and $O-18$ isotopes
(d) A mixture of $O-16$, $O-17$ and $O-18$ isotopes
53. Isotopes were discovered by [AMU 1983; AFMC 1995]
(a) Aston (b) Soddy
(c) Thomson (d) Millikan
54. Which of the following are iso-electronic [CBSE 2002]
(a) CO_2 and NO (b) SO_2 and CO_2

(c) CN and CO (d) NO_2 and CO_2

55. Which of the following are pairs of isotopes [Bihar CEE 1982]
(a) ${}_1^2H^+$ and ${}_1^3H$ (b) ${}_1^3H$ and ${}_2^4H^-$
(c) ${}_2^3He$ and ${}_2^4He$ (d) ${}_6^{12}C$ and ${}_7^{14}N^+$
56. Which among the following isotope is not found in natural uranium
(a) ${}_{92}U^{234}$ (b) ${}_{92}U^{235}$
(c) ${}_{92}U^{238}$ (d) ${}_{92}U^{239}$
57. An isotone of ${}_{32}^{76}Ge$ is (one or more are correct) [MP PMT 1990] [IIT 1984; MADT Bihar 1995; MP PMT 1995]
(a) ${}_{32}^{77}Ge$ (b) ${}_{33}^{77}As$
(c) ${}_{34}^{77}Se$ (d) ${}_{34}^{78}Se$



Critical Thinking

Objective Questions

1. ${}_{11}^{23}Na$ is the more stable isotope of *Na*. Find out the process by which ${}_{11}^{24}Na$ can undergo radioactive decay [IIT Screening 2003]
(a) β^- emission (b) α emission
(c) β^+ emission (d) K electron capture
2. Oxygen contains 90% of O^{16} and 10% of O^{18} . Its atomic mass is
(a) 17.4 (b) 16.2
(c) 16.5 (d) 17
3. The missing particle in the reaction, ${}_{92}^{235}U + {}_0^1n \rightarrow {}_{56}Ba^{146} + \dots + 3{}_0^1n$ is [DPMT 2001]
(a) ${}_{32}^{87}Ge$ (b) ${}_{35}^{89}Br$
(c) ${}_{36}^{87}Kr$ (d) ${}_{35}^{86}Br$
4. Sulphur-35 (34.96903 *amu*) emits a β -particle but no γ -rays, the product is chlorine-35 (34.96885 *amu*). The maximum energy emitted by the β -particle is [DPMT 2004]
(a) 0.016767 *MeV* (b) 1.6758 *MeV*
(c) 0.16758 *MeV* (d) 16.758 *MeV*
5. A radioactive substance has a constant activity of 2000 disintegration/minute. The material is separated into two fractions, one of which has an initial activity of 1000 disintegrations per second while the other fraction decays with $t_{1/2} = 24$ hours. The total activity in both samples after 48 hours of separation is
(a) 2000 (b) 1250
(c) 1000 (d) 1500
6. How many alpha particles are emitted per second by 1 microgram of radium
(a) 3.62×10^4 / sec (b) 0.362×10^4 / sec
(c) 362×10^4 / sec (d) 36.2×10^4 / sec
7. If 1 microgram of radium has disintegrated for 500 years, how many alpha particles will be emitted per second
(a) 2.92×10^4 / sec (b) 292×10^4 / sec

[EAMCET 1991]

- (c) $0.292 \times 10^4 / \text{sec}$ (d) $29.2 \times 10^4 / \text{sec}$
8. A radioactive nuclide X decays at the rate of 1.00×10^5 disintegration $s^{-1} g^{-1}$. Radium decays at the rate of 3.70×10^{10} disintegration $s^{-1} g^{-1}$. The activity of X in millicuries g^{-1} ($mci g^{-1}$) is [MP PET 2001]
- (a) 0.027 (b) 0.270×10^{-5}
(c) 0.00270 (d) 0.000270
9. If ${}_{92}U^{235}$ nucleus absorbs a neutron and disintegrates in ${}_{54}Xe^{139}$, ${}_{38}Sr^{94}$ and X , then what will be the product X [CBSE 2002]
- (a) α -particle (b) β -particle
(c) 2-neutrons (d) 3-neutrons
10. The half-life of a radioactive isotope is 3 hours. Value of its disintegration constant is [BHU 2002]
- (a) 0.231 per hr (b) 2.31 per hr
(c) 0.2079 per hr (d) 2.079 per hr
11. The activity of carbon-14 in a piece of an ancient wood is only 12.5%. If the half-life period of carbon-14 is 5760 years, the age of the piece of wood will be ($\log 2 = 0.3010$) [MP PMT 1999]
- (a) 17.281×10^2 years (b) 172.81×10^2 years
(c) 1.7281×10^2 years (d) 1728.1×10^2 years
12. The radium and uranium atoms in a sample of uranium mineral are in the ratio of $1 : 2.8 \times 10^6$. If half-life period of radium is 1620 years, the half-life period of uranium will be [MP PMT 1999]
- (a) 45.3×10^9 years (b) 45.3×10^{10} years
(c) 4.53×10^9 years (d) 4.53×10^{10} years
13. Half-life of radium is 1580 yrs. Its average life will be [AIIMS 1999; AFMC 1999; CPMT 1999]
- (a) 2.5×10^3 yrs (b) 1.832×10^3 yrs
(c) 2.275×10^3 yrs (d) 8.825×10^2 yrs
14. 8 gms of a radioactive substance is reduced to 0.5 g after 1 hour. The $t_{1/2}$ of the radioactive substance is [DCE 2000]
- (a) 15 min (b) 30 min
(c) 45 min (d) 10 min
15. A first order nuclear reaction is half completed in 45 minutes. How long does it need 99.9% of the reaction to be completed [KCET 2001]
- (a) 5 hours (b) 7.5 hours
(c) 10 hours (d) 20 hours
16. Number of α -particles emitted per second by a radioactive element falls to $1/32$ of its original value in 50 days. The half-life-period of this elements is [AMU 2001]
- (a) 5 days (b) 15 days
(c) 10 days (d) 20 days
17. What is the half-life of a radioactive substance if 87.5% of any given amount of the substance disintegrates in 40 minutes [Kerala CET 1996]
- (a) 160 min (b) 10 min
(c) 20 min (d) 13 min 20 sec
18. A radioactive isotope has a $t_{1/2}$ of 10 days. If today 125 gm of it is left, what was its weight 40 days earlier
- (a) 600 gm (b) 1000 gm
(c) 1250 gm (d) 2000 gm
19. The half-life of ${}_6C^{14}$, if its decay constant is 6.31×10^{-4} is [CBSE PMT 2001]
- (a) 1098 yrs (b) 109.8 yrs
(c) 10.98 yrs (d) 1.098 yrs
20. A radioactive sample has a half-life of 1500 years. A sealed tube containing 1 gm of the sample will contain after 3000 years [MNR 1994; UPSEAT 1994]
- (a) 1 gm of the sample
(b) 0.5 gm of the sample
(c) 0.25 gm of the sample
(d) 0.00 gm of the sample
21. The half-life of a radioactive isotope is three hours. If the initial mass of the isotope were 256 g, the mass of it remaining undecayed after 18 hours would be [AIEEE 2003]
- (a) 4.0 g (b) 8.0 g
(c) 12.0 g (d) 16.0 g
22. $\frac{15}{16}$ th of a radioactive sample decays in 40 days half-life of the sample is [DCE 2001]
- (a) 100 days (b) 10 days
(c) 1 day (d) $\log_e 2$ days
23. A radioactive element with half-life 6.5 hrs has 48×10^{19} atoms. Number of atoms left after 26 hrs [BHU 2003]
- (a) 24×10^{19} (b) 12×10^{19}
(c) 3×10^{19} (d) 6×10^{19}
24. The half-life of 1 gm of radioactive sample is 9 hours. The radioactive decay obeys first order kinetics. The time required for the original sample to reduce to 0.2 gm is [AMU (Engg.) 2002]
- (a) 15.6 hours (b) 156 hours
(c) 20.9 hours (d) 2.09 hours
25. The half-life period of a radioactive substance is 140 days. After how much time 15 g will decay from 16 g sample of it [AFMC 2002]
- (a) 140 days (b) 560 days
(c) 280 days (d) 420 days
26. Percentage of a radioactive element decayed after 20 sec when half-life is 4 sec [BHU 2003]
- (a) 92.25 (b) 96.87
(c) 50 (d) 75
27. Consider an α -particle just in contact with a ${}_{92}U^{238}$ nucleus. Calculate the coulombic repulsion energy (i.e. the height of the coulombic barrier between U_{238} and alpha particle) assuming that the distance between them is equal to the sum of their radii [UPSEAT 2001]
- (a) $23.8517 \times 10^4 eV$
(b) $26.147738 \times 10^4 eV$

- (c) $25.3522 \times 10^4 \text{ eV}$
(d) $20.2254 \times 10^4 \text{ eV}$
28. The half-life period of Pb^{210} is 22 years. If 2 gm of Pb^{210} is taken, then after 11 years how much of Pb^{210} will be left
[KCET 2001]
- (a) 1.414 gm (b) 2.428 gm
(c) 3.442 gm (d) 4.456 gm
29. A wood specimen from an archeological centre shows a $^{14}_6C$ activity of 5.0 counts/min/gm of carbon. What is the age of the specimen ($t_{1/2}$ for $^{14}_6C$ is 5000 years) and a freshly cut wood gives 15 counts/min/gm of carbon
[AMU (Engg.) 2002]
- (a) $5.78 \times 10^4 \text{ years}$ (b) $9.85 \times 10^4 \text{ years}$
(c) $7.85 \times 10^3 \text{ years}$ (d) $0.85 \times 10^4 \text{ years}$
30. $_{92}U^{235} + n \rightarrow \text{fission product} + \text{neutron} + 3.20 \times 10^{-11} \text{ J}$. The energy released when 1g of $_{92}U^{235}$ undergoes fission is
[CBSE PMT 1997]
- (a) $12.75 \times 10^8 \text{ kJ}$ (b) $18.60 \times 10^9 \text{ kJ}$
(c) $8.21 \times 10^7 \text{ kJ}$ (d) $6.55 \times 10^6 \text{ kJ}$
31. The triad of nuclei that is isotonic is
[IIT 1988; DCE 2000; MP PMT 2004]
- (a) ${}_6C^{14}, {}_7N^{15}, {}_9F^{17}$
(b) ${}_6C^{12}, {}_7N^{14}, {}_9F^{19}$
(c) ${}_6C^{14}, {}_7N^{14}, {}_9F^{17}$
(d) ${}_6C^{14}, {}_7N^{14}, {}_9F^{19}$
32. The relative abundance of two isotopes of atomic weight 85 and 87 is 75% and 25% respectively. The average atomic weight of element is
[DCE 2003]
- (a) 75.5 (b) 85.5
(c) 40.0 (d) 86.0

Assertion & Reason

For AIIMS Aspirants

Read the assertion and reason carefully to mark the correct option out of the options given below :

- (a) If both assertion and reason are true and the reason is the correct explanation of the assertion.
(b) If both assertion and reason are true but reason is not the correct explanation of the assertion.
(c) If assertion is true but reason is false.
(d) If the assertion and reason both are false.
(e) If assertion is false but reason is true.

1. Assertion : Mass number of an atom is equal to total number of nucleons present in the nucleus.
Reason : Mass number defines the identity of an atom.
2. Assertion : ${}_1H^1, {}_1H^2$ and ${}_1H^3$ are isotopes of hydrogen.

- Reason : Nuclides of the same element of different mass numbers are called isotopes of that element.
3. Assertion : The activity of 1 g pure uranium-235 will be greater than the same amount present as U_3O_8 .
Reason : In the combined state, the activity of the radioactive element decreases.
4. Assertion : Nuclear forces are called short range forces.
Reason : Nuclear forces operate over very small distance i.e., 10^{-15} m or 1 fermi.
5. Assertion : An example of K -capture is $^{133}_{56}Ba + e^- \rightarrow ^{133}_{55}Cs + X - \text{ray}$.
Reason : The atomic number decreases by one unit as result of K -capture.
6. Assertion : Radioactive heavy nuclei decay by a series of α - and β -emission, to form a stable isotope of lead.
Reason : Radioactivity is a physical phenomenon.
7. Assertion : Actinium series is so called because it starts with an isotope of actinium.
Reason : Actinium is formed in the nature as such and is not formed from the disintegration of any other radioisotope.
8. Assertion : For maximum stability N/P ratio must be equal to 1.
Reason : Loss of α - and β -particles has no role in N/P ratio.
9. Assertion : The neutrons are better initiators of nuclear reactions, than the protons, deuterons or α -particles of the same energy.
Reason : Neutrons are uncharged particles and hence, they are not repelled by positively charged nucleus.
10. Assertion : Breeder reactor produces fissile $_{94}Pu^{239}$ from non-fissile uranium.
Reason : A breeder reactor is one that produces more fissionable nuclei than it consumes.
11. Assertion : The activation energies for fusion reactions are very low.
Reason : They require very low temperature to overcome electrostatic repulsion between the nuclei.
12. Assertion : The archeological studies are based on the radioactive decay of carbon-14 isotope.
Reason : The ratio of $C-14$ to $C-12$ in the animals and plants is same as that in the atmosphere.
13. Assertion : Photochemical smog is produced by nitrogen oxides.
Reason : Vehicular pollution is a major source of nitrogen oxides.
14. Assertion : A nuclear binding energy per nucleon is in the order $^9_4Be > ^7_3Li > ^4_2He$.

- Reason : Binding energy per nuclear increases linearly with difference in number of neutrons and protons.
15. Assertion : Nuclear fission is always accompanied by release of energy.
- Reason : Nuclear fission is a chain process.
- [AIIMS 1994]
16. Assertion : Protons are more effective than neutrons of equal energy in causing artificial disintegration of atoms.
- Reason : Neutrons are neutral they penetrate the nucleus. [AIIMS]
17. Assertion : A beam of electrons deflects more than a beam of α -particles in an electric field.
- Reason : Electrons possess negative charge while α -particles possess positive charge.
- [AIIMS 2002]
18. Assertion : ${}^{22}_{11}\text{Na}$ emits a positron giving ${}^{22}_{12}\text{Mg}$.
- Reason : In β^+ emission neutron is transformed into proton.
- [AIIMS 1994]

Answers

Nucleus (Stability and Reaction)

1	b	2	b	3	a	4	a	5	b
6	d	7	b	8	c	9	c	10	d
11	b	12	c	13	c	14	c	15	c
16	c	17	d	18	c	19	a	20	b
21	b	22	a	23	b	24	d	25	c
26	c	27	b	28	c	29	b	30	a
31	c	32	d	33	b	34	d	35	c
36	b	37	b	38	a	39	a	40	a
41	b	42	b	43	a	44	d	45	b
46	d	47	b	48	a	49	b	50	d
51	d	52	a	53	b	54	c	55	d
56	b	57	b	58	d	59	b	60	c
61	b	62	d	63	d	64	d	65	a
66	b	67	a	68	a				

Radioactivity and α , β and γ rays

1	c	2	d	3	a	4	b	5	c
6	b	7	c	8	c	9	b	10	b
11	a	12	a	13	a	14	b	15	c
16	c	17	a	18	a	19	c	20	b

21	a	22	c	23	d	24	b	25	c
26	d	27	d	28	c	29	c	30	a
31	d	32	c	33	a	34	b	35	a
36	c	37	a	38	acd	39	a		

Causes of radioactivity and Group displacement law

1	b	2	d	3	d	4	d	5	c
6	c	7	c	8	a	9	b	10	a
11	a	12	c	13	d	14	b	15	a
16	a	17	a	18	a,b,c	19	c	20	c
21	b	22	d	23	d	24	a	25	b
26	d	27	b	28	b	29	b	30	a
31	a	32	a	33	c	34	b	35	a
36	b	37	c	38	b	39	c	40	d
41	a	42	b	43	c	44	c	45	d
46	b	47	b	48	d	49	a	50	d
51	a	52	a	53	d	54	d	55	b
56	a	57	d	58	c	59	c	60	a
61	a	62	c	63	d	64	d	65	a
66	d	67	b	68	c	69	b	70	a
71	c	72	c	73	d	74	a	75	b

Rate of decay and Half-life

1	c	2	a	3	b	4	a	5	d
6	a	7	d	8	d	9	d	10	c
11	a	12	d	13	d	14	a	15	c
16	d	17	b	18	c	19	b	20	a
21	b	22	a	23	c	24	a	25	a
26	b	27	a	28	c	29	a	30	b
31	a	32	c	33	d	34	a	35	b
36	d	37	d	38	c	39	c	40	a
41	c	42	b	43	b	44	d	45	c
46	c	47	c	48	b	49	a	50	c
51	c	52	c	53	c	54	b	55	c
56	b	57	a	58	b	59	d	60	b
61	d	62	b	63	c	64	a	65	b
66	d	67	d	68	c	69	d	70	c
71	d	72	a	73	c	74	a	75	a
76	c	77	b	78	b	79	a	80	a
81	a	82	a	83	b	84	c		

Artificial transmutation

1	b	2	c	3	d	4	d	5	c
6	a	7	a	8	b	9	d	10	b
11	a	12	c	13	c	14	b	15	a
16	d	17	d	18	b	19	a	20	c
21	b	22	a	23	c	24	b	25	d
26	c	27	b	28	a	29	d	30	a
31	d	32	b	33	d	34	c	35	b
36	a	37	b	38	c	39	d	40	c
41	d	42	b	43	c	44	b	45	b
46	c	47	b	48	b	49	c	50	c
51	d	52	c	53	a	54	d	55	d
56	d	57	b,d	58	a	59	b	60	a
61	d	62	a	63	a	64	d	65	d
66	d	67	d	68	b	69	b	70	d
71	a	72	b	73	a	74	a	75	a
76	b	77	d	78	c	79	a	80	c
81	b								

Isotopes-Isotones and Nuclear isomers

1	b	2	a	3	d	4	a	5	c
6	c	7	c	8	b	9	b	10	d
11	a	12	b	13	a	14	a	15	a
16	c	17	b	18	c	19	c	20	a
21	d	22	d	23	a	24	c	25	c
26	d	27	c	28	b	29	b	30	b
31	cd	32	d	33	a	34	c	35	c
36	b	37	a	38	a	39	a	40	b
41	c	42	a	43	a	44	d	45	b
46	a	47	d	48	d	49	a	50	a
51	b	52	d	53	b	54	c	55	ac
56	d	57	bd						

Critical Thinking Questions

1	a	2	b	3	c	4	a	5	a
6	a	7	a	8	b	9	d	10	a
11	b	12	c	13	c	14	a	15	b
16	c	17	d	18	d	19	a	20	c
21	a	22	b	23	c	24	c	25	b
26	b	27	b	28	a	29	c	30	c
31	a	32	b						

Assertion & Reason

1	c	2	a	3	d	4	a	5	b
---	---	---	---	---	---	---	---	---	---

6	c	7	d	8	c	9	a	10	a
11	d	12	a	13	b	14	d	15	b
16	e	17	b	18	d				

AS

Answers and Solutions

Nucleus (Stability and Reaction)

- (b) Protons + Neutrons = Nucleons
- (b) A deuteron (${}_1H^2$) contains a neutron and a proton
- (a) Low binding energy causes radioactivity.
- (a) ${}_7N^{14} + {}_2He^4 \rightarrow {}_8O^{17} + {}_1H^1$
- (b) Follow Einstein mass-energy relation.
- (d) Mass (weight) of positron and electron is $9.11 \times 10^{-31} \text{ kg}$.