

10.  $H_2S$  reacts with halogens, the halogens [JIPMER 2000]  
 (a) Form sulphur halides (b) Are oxidised  
 (c) Are reduced (d) None of these
11.  $H_2O_2$  reduces  $K_4Fe(CN)_6$  [MP PMT 1985]  
 (a) In neutral solution (b) In acidic solution  
 (c) In non-polar solvent (d) In alkaline solution
12. Max. number of moles of electrons taken up by one mole of  $NO_3^-$  when it is reduced to [DPMT 2002]  
 (a)  $NH_3$  (b)  $NH_2OH$   
 (c)  $NO$  (d)  $NO_2$
13. In the reaction  $3Mg + N_2 \rightarrow Mg_3N_2$  [MP PMT 1999]  
 (a) Magnesium is reduced (b) Magnesium is oxidized  
 (c) Nitrogen is oxidized (d) None of these
14. When sodium metal is dissolved in liquid ammonia, blue colour solution is formed. The blue colour is due to [NCERT 1981]  
 (a) Solvated  $Na^+$  ions (b) Solvated electrons  
 (c) Solvated  $NH_2^-$  ions (d) Solvated protons
15. Following reaction describes the rusting of iron  
 $4Fe + 3O_2 \rightarrow 4Fe^{3+} + 6O^{2-}$   
 Which one of the following statement is incorrect [NCERT 1981; MNR 1991; AIIMS 1998]  
 (a) This is an example of a redox reaction  
 (b) Metallic iron is reduced to  $Fe^{3+}$   
 (c)  $Fe^{3+}$  is an oxidising agent  
 (d) Metallic iron is a reducing agent
16.  $SnCl_2$  gives a precipitate with a solution of  $HgCl_2$ . In this process  $HgCl_2$  is [CPMT 1983]  
 (a) Reduced  
 (b) Oxidised  
 (c) Converted into a complex compound containing both  $Sn$  and  $Hg$   
 (d) Converted into a chloro complex of  $Hg$
17. Oxidation involves [NCERT 1971, 81; CPMT 1980, 82, 83; MP PMT 1983]  
 (a) Loss of electrons  
 (b) Gain of electrons  
 (c) Increase in the valency of negative part  
 (d) Decrease in the valency of positive part
18. Incorrect statement regarding rusting is [MP PET 2000]  
 (a) Metallic iron is oxidised to  $Fe^{3+}$  ions  
 (b) Metallic iron is reduced to  $Fe^{2+}$  ions  
 (c) Oxygen gas is reduced to oxide ion  
 (d) Yellowish – brown product is formed
19. When copper turnings are added to silver nitrate solution, a blue coloured solution is formed after some time. It is because, copper [CPMT 1974, 79; DPMT 2000]  
 (a) Displaces silver from the solution  
 (b) Forms a blue coloured complex with  $AgNO_3$   
 (c) Is oxidised to  $Cu^{2+}$   
 (d) Is reduced to  $Cu^{2+}$
20. Solution of sodium metal in liquid ammonia is strongly reducing due to the presence of the following in the solution [NCERT 1977; KCET (Med.) 2000]  
 (a) Sodium atoms (b) Solvated electrons  
 (c) Sodium hydride (d) Sodium amide
21. When  $Sn^{2+}$  changes to  $Sn^{4+}$  in a reaction [CPMT 1981]  
 (a) It loses two electrons (b) It gains two electrons  
 (c) It loses two protons (d) It gains two protons
22. Oxidation of thiosulphate ( $S_2O_3^{2-}$ ) ion by iodine gives [NCERT 1976]  
 (a)  $SO_3^{2-}$  (b)  $SO_4^{2-}$   
 (c)  $S_4O_6^{2-}$  (d)  $S_2O_6^{2-}$
23.  $Zn^{2+}(aq) + 2e^- \rightarrow Zn(s)$ . This is [CPMT 1985]  
 (a) Oxidation (b) Reduction  
 (c) Redox reaction (d) None of these
24. One gas bleaches the colour of flowers by reduction while the other by oxidation [EAMCET 1980]  
 (a)  $CO$  and  $Cl_2$  (b)  $SO_2$  and  $Cl_2$   
 (c)  $H_2S$  and  $Br_2$  (d)  $NH_3$  and  $SO_2$
25. Reduction involves NCERT 1972  
 (a) Loss of electrons  
 (b) Gain of electrons  
 (c) Increase in the valency of positive part  
 (d) Decrease in the valency of negative part
26. In a reaction between zinc and iodine, in which zinc iodide is formed, what is being oxidised [NCERT 1975]  
 (a) Zinc ions (b) Iodide ions  
 (c) Zinc atom (d) Iodine
27. Which one of the following reactions does not involve either oxidation or reduction [EAMCET 1982]  
 (a)  $VO_2^+ \rightarrow V_2O_3$  (b)  $Na \rightarrow Na^+$   
 (c)  $CrO_4^{2-} \rightarrow Cr_2O_7^{2-}$  (d)  $Zn^{2+} \rightarrow Zn$
28. In the following reaction,  
 $3Br_2 + 6CO_3^{2-} + 3H_2O = 5Br^- + BrO_3^- + 6HCO_3^-$   
 [MP PMT 1994, 95]  
 (a) Bromine is oxidised and carbonate is reduced  
 (b) Bromine is reduced and water is oxidised  
 (c) Bromine is neither reduced nor oxidised  
 (d) Bromine is both reduced and oxidised
29. In the following reaction,  
 $4P + 3KOH + 3H_2O \rightarrow 3KH_2PO_2 + PH_3$  [Pb. PMT 2002]  
 (a)  $P$  is oxidized as well as reduced  
 (b)  $P$  is reduced only  
 (c)  $P$  is oxidised only  
 (d) None of these
30. In the following reaction  
 $Cr_2O_7^{2-} + 14H^+ + 6I^- \rightarrow 2Cr^{3+} + 3H_2O + 3I_2$   
 Which element is reduced [CPMT 1976]  
 (a)  $Cr$  (b)  $H$   
 (c)  $I$  (d)  $I$
31. The conversion of sugar  $C_{12}H_{22}O_{11} \rightarrow CO_2$  is  
 (a) Oxidation  
 (b) Reduction  
 (c) Neither oxidation nor reduction  
 (d) Both oxidation and reduction
32. Which halide is not oxidised by  $MnO_2$  [MNR 1985; JIPMER 2000]  
 (a)  $F$  (b)  $Cl$

- (c)  $Br$  (d)  $I$
33. When  $Fe^{2+}$  changes to  $Fe^{3+}$  in a reaction  
(a) It loses an electron (b) It gains an electron  
(c) It loses a proton (d) It gains a proton
34. In acid solution, the reaction  $MnO_4^- \rightarrow Mn^{2+}$  involves [MP PMT 1989]  
(a) Oxidation by 3 electrons  
(b) Reduction by 3 electrons  
(c) Oxidation by 5 electrons  
(d) Reduction by 5 electrons
35. When iron or zinc is added to  $CuSO_4$  solution, copper is precipitated. It is due to [CPMT 1974, 79]  
(a) Oxidation of  $Cu^{+2}$  (b) Reduction of  $Cu^{+2}$   
(c) Hydrolysis of  $CuSO_4$  (d) Ionization of  $CuSO_4$
36. In the reaction,  $4Fe + 3O_2 \rightarrow 4Fe^{3+} + 6O^{2-}$  which of the following statement is incorrect [UPSEAT 2001, 02]  
(a) A Redox reaction  
(b) Metallic iron is a reducing agent  
(c)  $Fe^{3+}$  is an oxidising agent  
(d) Metallic iron is reduced to  $Fe^{3+}$
37. Which of the following is redox reaction [CBSE PMT 1997]  
(a)  $H_2SO_4$  with  $NaOH$   
(b) In atmosphere,  $O_3$  from  $O_2$  by lightning  
(c) Evaporation of  $H_2O$   
(d) Nitrogen oxides form nitrogen and oxygen by lightning
- (c)  $HClO_2$  (d)  $HOCl$
6. Identify the correct statement about  $H_2O_2$  [AIIMS 1996]  
(a) It acts as reducing agent only  
(b) It acts as both oxidising and reducing agent  
(c) It is neither an oxidiser nor reducer  
(d) It acts as oxidising agent only
7. Several blocks of magnesium are fixed to the bottom of a ship to [AIEEE 2003]  
(a) Keep away the sharks  
(b) Make the ship lighter  
(c) Prevent action of water and salt  
(d) Prevent puncturing by under-sea rocks
8. Which of the following behaves as both oxidising and reducing agents [AFMC 1995]  
(a)  $H_2SO_4$  (b)  $SO_2$   
(c)  $H_2S$  (d)  $HNO_3$
9. The reaction  $H_2S + H_2O_2 \rightarrow 2H_2O + S$  shows [JIPMER 2001]  
(a) Oxidizing action of  $H_2O_2$   
(b) Reducing action of  $H_2O_2$   
(c) Alkaline nature of  $H_2O_2$   
(d) Acidic nature of  $H_2O_2$
10. Which of the following is not a reducing agent [EAMCET 1987]  
(a)  $NaNO_2$  (b)  $NaNO_3$   
(c)  $HI$  (d)  $SnCl_2$
11. Which of the following cannot work as oxidising agent [CPMT 1996]  
(a)  $O_2$  (b)  $KMnO_4$   
(c)  $I_2$  (d) None of these
12.  $H_2O_2$  is used as [CPMT 1994]  
(a) An oxidant only  
(b) A reductant only  
(c) An acid only  
(d) An oxidant, a reductant and an acid
13. In  $C + H_2O \rightarrow CO + H_2$ ,  $H_2O$  acts as [AFMC 1988]  
(a) Oxidising agent (b) Reducing agent  
(c) (a) and (b) both (d) None of these
14. Strongest reducing agent is [CPMT 1977; BHU 1984, 96; MP PET 1990; AMU 1999]  
(a)  $F^-$  (b)  $Cl^-$   
(c)  $Br^-$  (d)  $I^-$
15. A solution of sulphur dioxide in water reacts with  $H_2S$  precipitating sulphur. Here sulphur dioxide acts as [NCERT 1980]  
(a) As oxidising agent (b) A reducing agent  
(c) An acid (d) A catalyst
16. Which of these substances is a good reducing agent [NCERT 1979; CPMT 1988]  
(a)  $NaOCl$  (b)  $HI$   
(c)  $FeCl_3$  (d)  $KBr$
17. The strongest reducing agent is [MNR 1982]  
(a)  $HNO_2$  (b)  $H_2S$

### Oxidizing and Reducing agent

1. Equation  $H_2S + H_2O_2 \rightarrow S + 2H_2O$  represents [UPSEAT 2001]  
(a) Acidic nature of  $H_2O_2$   
(b) Basic nature of  $H_2O_2$   
(c) Oxidising nature of  $H_2O_2$   
(d) Reducing nature of  $H_2O_2$
2. In the reaction  
 $C_2O_4^{2-} + MnO_4^- + H^+ \rightarrow Mn^{2+} + CO_2 + H_2O$   
the reductant is [EAMCET 1991]  
(a)  $C_2O_4^{2-}$  (b)  $MnO_4^-$   
(c)  $Mn^{2+}$  (d)  $H^+$
3. A reducing agent is a substance which can [CPMT 1971, 74, 76, 78, 80; NCERT 1976]  
(a) Accept electron (b) Donate electrons  
(c) Accept protons (d) Donate protons
4. Which of the following is the most powerful oxidizing agent [MNR 1990; CPMT 2003]  
(a)  $F_2$  (b)  $Cl_2$   
(c)  $Br_2$  (d)  $I_2$
5. Of the four oxyacids of chlorine the strongest oxidising agent in dilute aqueous solution is [MP PET 2000]  
(a)  $HClO_4$  (b)  $HClO_3$

[JEE Orissa 2004]

- (c)  $H_2SO_3$  (d)  $SnCl_2$
18. Which one is an oxidising agent [DPMT 1996]  
 (a)  $FeSO_4$   
 (b)  $HNO_3$   
 (c)  $FeSO_4 \cdot (NH_4)_2SO_4 \cdot 6H_2O$   
 (d)  $H_2SO_4$
19. In which of the following reactions  $H_2O_2$  is a reducing agent [CPMT 1981; NCERT 1981; BHU 1999]  
 (a)  $2FeCl_2 + 2HCl + H_2O_2 \rightarrow 2FeCl_3 + 2H_2O$   
 (b)  $Cl_2 + H_2O_2 \rightarrow 2HCl + O_2$   
 (c)  $2HI + H_2O_2 \rightarrow 2H_2O + I_2$   
 (d)  $H_2SO_3 + H_2O_2 \rightarrow H_2SO_4 + H_2O$
20. When  $NaCl$  is dissolved in water the sodium ion becomes [NCERT 1976]  
 (a) Oxidised (b) Reduced  
 (c) Hydrolysed (d) Hydrated
21. Strongest reducing agent is [MNR 1984, 89]  
 (a)  $K$  (b)  $Mg$   
 (c)  $Al$  (d)  $Br$   
 (e)  $Na$
22. Which substance is serving as a reducing agent in the following reaction  
 $14H^+ + Cr_2O_7^{2-} + 3Ni \rightarrow 2Cr^{3+} + 7H_2O + 3Ni^{2+}$   
 [CBSE PMT 1994; AFMC 2000; DPMT 2001]  
 (a)  $H_2O$  (b)  $Ni$   
 (c)  $H^+$  (d)  $Cr_2O_7^{2-}$
23. Which of the following acid possesses oxidising, reducing and complex forming properties [MNR 1985]  
 (a)  $HNO_3$  (b)  $H_2SO_4$   
 (c)  $HCl$  (d)  $HNO_2$
24. Which one is oxidising substance [CPMT 1997]  
 (a)  $C_2H_2O_2$  (b)  $CO$   
 (c)  $H_2S$  (d)  $CO_2$
25. The compound that can work both as oxidising and reducing agent is [CPMT 1986; MP PET 2000]  
 (a)  $KMnO_4$  (b)  $H_2O_2$   
 (c)  $BaO_2$  (d)  $K_2Cr_2O_7$
26. Which one is oxidising agent in the reaction below  
 $2CrO_4^{2-} + 2H^+ \rightarrow Cr_2O_7^{2-} + H_2O$  [CPMT 1997]  
 (a)  $H^+$  (b)  $Cr_2O_4^-$   
 (c)  $Cr^{++}$  (d) None of these
27. Which is the best description of the behaviour of bromine in the reaction given below  
 $H_2O + Br_2 \rightarrow HOBr + HBr$  [CBSE PMT 2004]  
 (a) Oxidised only  
 (b) Reduced only  
 (c) Proton acceptor only  
 (d) Both oxidised and reduced
28. What is the oxidising agent in chlorine water  
 (a)  $HCl$  (b)  $HClO_2$   
 (c)  $HOCl$  (d) None of these
29. In the reaction  
 $Ag_2O + H_2O_2 \rightarrow 2Ag + H_2O + O_2$ , the  $H_2O_2$  acts as [BHU 2004]  
 (a) Reducing agent (b) Oxidising agent  
 (c) Bleaching agent (d) None of the above
30. In the reaction  
 $HAsO_2 + Sn^{2+} \rightarrow As + Sn^{4+} + H_2O$  oxidising agent is [BVP 2004]  
 (a)  $Sn^{2+}$  (b)  $Sn^{4+}$   
 (c)  $As$  (d)  $HAsO_2$
31. Which of the following substances acts as an oxidising as well as a reducing agent [UPSEAT 2004; DCE 2004]  
 (a)  $Na_2O$  (b)  $SnCl_2$   
 (c)  $Na_2O_2$  (d)  $NaNO_2$
32. In the reaction  
 $P + NaOH \rightarrow PH_3 + NaH_2PO_2$  [MP PET 2004]  
 (a)  $P$  is oxidised only  
 (b)  $P$  is reduced only  
 (c)  $P$  is oxidized as well as reduced  
 (d)  $Na$  is reduced

## Oxidation number and Oxidation state

1. The oxidation number of  $C$  in  $CO_2$  is [MP PET 2001]  
 (a)  $-2$  (b)  $+2$   
 (c)  $-4$  (d)  $+4$
2. The oxidation number of  $As$  is [RPMT 1997]  
 (a)  $+2$  and  $+3$  (b)  $+3$  and  $+5$   
 (c)  $+3$  and  $+4$  (d) None of these
3. The oxidation number of  $Ba$  in barium peroxide is [Pb. PMT 2002]  
 (a)  $+6$  (b)  $+2$   
 (c)  $1$  (d)  $+4$
4.  $HNO_2$  acts both as reductant and oxidant, while  $HNO_3$  acts only as oxidant. It is due to their [AIIMS 2000]  
 (a) Solubility ability  
 (b) Maximum oxidation number  
 (c) Minimum oxidation number  
 (d) Minimum number of valence electrons
5. Chlorine is in  $+1$  oxidation state in [MP PMT 1981; NCERT 1974; CPMT 1971, 78]  
 (a)  $HCl$  (b)  $HClO_4$   
 (c)  $ICl$  (d)  $Cl_2O$
6. The valency of  $Cr$  in the complex  $[Cr(H_2O)_4Cl_2]^+$  [MP PMT 2000]  
 (a)  $1$  (b)  $3$   
 (c)  $5$  (d)  $6$
7. In the conversion  $Br_2 \rightarrow BrO_3^-$ , the oxidation state of bromine changes from [EAMCET 1990; AMU 1999; RPMT 2002]  
 (a)  $-1$  to  $-1$  (b)  $0$  to  $-1$   
 (c)  $0$  to  $+5$  (d)  $0$  to  $-5$

8. In the chemical reaction  $Cl_2 + H_2S \rightarrow 2HCl + S$ , the oxidation number of sulphur changes from [MP PMT 1999]  
(a) 0 to 2 (b) 2 to 0  
(c) -2 to 0 (d) -2 to -1
9. Oxidation number of cobalt in  $K[Co(CO)_4]$  is [KCET 1996]  
(a) +1 (b) +3  
(c) -1 (d) -3
10. When  $K_2Cr_2O_7$  is converted to  $K_2CrO_4$ , the change in the oxidation state of chromium is [NCERT 1981]  
(a) 0 (b) 6  
(c) 4 (d) 3
11. The oxidation number of chlorine in  $HOCl$   
(a) -1 (b) 0  
(c) +1 (d) +2
12. Oxidation number of S in  $S^{2-}$  is [CPMT 1979]  
(a) -2 (b) 0  
(c) -6 (d) +2
13. Oxidation number of N in  $(NH_4)_2SO_4$  is [CPMT 1996]  
(a) -1/3 (b) -1  
(c) +1 (d) -3
14. In which compound, oxidation state of nitrogen is 1 [MP PMT 1989]  
(a) NO (b)  $N_2O$   
(c)  $NH_2OH$  (d)  $N_2H_4$
15. Oxidation number of nickel in  $Ni(CO)_4$  [AIIMS 1984; MNR 1985; CPMT 1997; MP PET/PMT 1998; AMU 2000; 01]  
(a) 0 (b) +4  
(c) -4 (d) +2
16. The oxidation number of sulphur in  $H_2SO_4$  is [CPMT 1979Pb. CET 2002]  
(a) -2 (b) +2  
(c) +4 (d) +6
17. Oxidation state of chlorine in perchloric acid is [EAMCET 1989]  
(a) -1 (b) 0  
(c) -7 (d) +7
18. Oxidation number of N in  $HNO_3$  is [BHU 1997]  
(a) -3.5 (b) +3.5  
(c) -3, +5 (d) +5
19. The oxidation number of Mn in  $MnO_4^{-1}$  is  
(a) +7 (b) -5  
(c) +6 (d) +5
20.  $Sn^{++}$  loses two electrons in a reaction. What will be the oxidation number of tin after the reaction  
(a) +2 (b) Zero  
(c) +4 (d) -2
21. The oxidation state of Mn in  $K_2MnO_4$  [CPMT 1982, 83, 84; DPMT 1982; NCERT 1973; AMU 2000]  
(a) +2 (b) +7  
(c) -2 (d) +6
22. Oxidation number of oxygen in  $O_2$  molecule is [CPMT 1984]  
(a) +1 (b) 0  
(c) +2 (d) -2
23. Maximum oxidation state of Cr is [RPMT 2002]  
(a) 3 (b) 4  
(c) 6 (d) 7
24. In which of the following compound transition metal has zero oxidation state [CBSE PMT 1999; BHU 2000]  
(a)  $CrO_5$  (b)  $NH_2.NH_2$   
(c)  $NOClO_4$  (d)  $[Fe(CO)_5]$
25. Carbon is in the lowest oxidation state in [NCERT 1979; MH CET 1999]  
(a)  $CH_4$  (b)  $CCl_4$   
(c)  $CF_4$  (d)  $CO_2$
26. Oxidation number of carbon in  $H_2C_2O_4$  is [CPMT 1982]  
(a) +4 (b) +3  
(c) +2 (d) -2
27. The oxidation number of Pt in  $[Pt(C_2H_4)Cl_3]^-$  is [MNR 1993]  
(a) +1 (b) +2  
(c) +3 (d) +4
28. The oxidation number of carbon in  $CH_2Cl_2$  is [CPMT 1976; Pb. PET 1999; AFMC 2004]  
(a) 0 (b) +2  
(c) -2 (d) +4
29. The oxidation states of phosphorus vary from [CPMT 1976]  
(a) -3 to +5 (b) -1 to +1  
(c) -3 to +3 (d) -5 to +1
30. The process in which oxidation number increases is known as [CPMT 1976]  
(a) Oxidation (b) Reduction  
(c) Auto-oxidation (d) None of the above
31. The oxidation number of S in  $H_2S_2O_8$  is [MP PET 2002]  
(a) +2 (b) +4  
(c) +6 (d) +7
32. The oxidation state of nitrogen in  $N_3H$  is [NCERT 1977, 81]  
(a)  $+\frac{1}{3}$  (b) +3  
(c) -1 (d)  $-\frac{1}{3}$
33. Which of the following statements is correct [AFMC 1997]  
(a) Hydrogen has oxidation number -1 and +1  
(b) Hydrogen has same electronegativity as halogens  
(c) Hydrogen will not be liberated at anode  
(d) Hydrogen has same ionization potential as alkali metals
34. The oxidation state of Cr in  $[Cr(NH_3)_4Cl_2]^+$  is [AIEEE 2005]  
(a) +3 (b) +2  
(c) +1 (d) 0
35. Sulphur has highest oxidation state in [EAMCET 1991]  
(a)  $SO_2$  (b)  $H_2SO_4$   
(c)  $Na_2S_2O_3$  (d)  $Na_2S_4O_6$
36. The oxidation number of Fe and S in iron pyrites are

- (a) 4, -2 (b) 2, -1 (c) 3, -1.5 (d) 3, -1 [RPMT 1997]
37. The oxidation number of nitrogen in  $NO_3^-$  is [CPMT 1982]
- (a) -1 (b) +2 (c) +3 (d) +5
38. Oxidation state of elemental carbon is [MNR 1983]
- (a) 0 (b) 1 (c) 2 (d) 3
39. The sum of the oxidation numbers of all the carbons in  $C_6H_5CHO$  is [EAMCET 1986]
- (a) +2 (b) 0 (c) +4 (d) -4
40. Which one of the following has the highest oxidation number of iodine [CPMT 1982]
- (a)  $KI_3$  (b)  $KI$  (c)  $IF_5$  (d)  $KIO_4$
41. The oxidation number of N in  $N_2H_5^+$  [Pb. PMT 2001]
- (a) -3 (b) -2 (c) -1 (d) +2
42. In which of the following compounds the oxidation number of carbon is maximum
- (a)  $HCHO$  (b)  $CHCl_3$  (c)  $CH_3OH$  (d)  $C_{12}H_{22}O_{11}$
43. The oxidation state of chlorine in  $KClO_4$  is [CPMT 1985]
- (a) -1 (b) +1 (c) +7 (d) -7
44. The oxidation state of I in  $H_4IO_6^-$  is [CBSE PMT 1994]
- (a) +7 (b) +5 (c) +1 (d) -1
45. An element which never has a positive oxidation number in any of its compounds [AIIMS 1981]
- (a) Boron (b) Oxygen (c) Chlorine (d) Fluorine
46. In an oxidation process, oxidation number [CPMT 1976]
- (a) Decreases (b) Increases (c) Does not change (d) First increases then decreases
47. If  $HNO_3$  changes into  $N_2O$ , the oxidation number is changed by [BHU 1987; AFMC 2001]
- (a) +2 (b) -1 (c) 0 (d) +4
48. The characteristic oxidation number of atoms in free metals is [NCERT 1975]
- (a) Minus one (b) Any number (c) One (d) Zero
49. In which one of the following changes there are transfer of five electrons [NCERT 1982]
- (a)  $MnO_4^- \rightarrow Mn^{2+}$  (b)  $CrO_4^{2-} \rightarrow Cr^{3+}$  (c)  $MnO_4^{2-} \rightarrow MnO_2$  (d)  $Cr_2O_7^{2-} \rightarrow 2Cr^{3+}$
50. Oxidation number of C in  $C_6H_{12}O_6$  is [KCET 1992]
- (a) +6 (b) -6 (c) 0 (d) +4
51. In which of the following compounds iron has lowest oxidation state [MNR 1984]
- (a)  $FeSO_4 \cdot (NH_4)_2SO_4 \cdot 6H_2O$  (b)  $K_4Fe(CN)_6$  (c)  $Fe(CO)_5$  (d)  $Fe_2O$  (e)  $K_2FeO_4$
52. The oxidation number of hydrogen in  $MH_2$  is [CPMT 1976]
- (a) +1 (b) -1 (c) +2 (d) -2
53. Oxidation number of iodine varies from [CPMT 1982]
- (a) -1 to +1 (b) -1 to +7 (c) +3 to +5 (d) -1 to +5
54. When  $SO_2$  is passed through acidic solution of potassium dichromate, then chromium sulphate is formed. Change in valency of chromium is [CPMT 1979]
- (a) +4 to +2 (b) +5 to +3 (c) +6 to +3 (d) +7 to +2
55. The oxidation states of the most electronegative element in the products of the reaction of  $BaO_2$  with dilute  $H_2SO_4$  are [IIT 1991; CBSE PMT 1992; BHU 2000]
- (a) 0 and -1 (b) -1 and -2 (c) -2 and 0 (d) -2 and +1
56. The highest oxidation state of Mn is shown by [MNR 1983; RPMT 1999]
- (a)  $K_2MnO_4$  (b)  $KMnO_4$  (c)  $MnO_2$  (d)  $Mn_2O_2$  (e)  $MnO$
57. The oxidation number of carbon in  $CH_2O$  is [IIT 1982; EAMCET 1985; MNR 1990; UPSEAT 2001 CPMT 1997, 2004]
- (a) -2 (b) +2 (c) 0 (d) +4
58. Oxidation state of oxygen in hydrogen peroxide is [DPMT 1984; 91; CPMT 1988; MNR 1994; UPSEAT 2001; RPMT 2002; JEE Orissa 2004]
- (a) -1 (b) +1 (c) 0 (d) -2
59. The oxidation number of Cr in  $K_2Cr_2O_7$  is [CPMT 1981, 85, 90, 93, 99; KCET 1992; BHU 1988, 98; AFMC 1991, 99; EAMCET 1986; MP PMT 1996, 99, 2002; MP PET/PMT 1998; Bihar CEE 1995; RPET 2000]
- (a) +6 (b) -7 (c) +2 (d) -2
60. In which of the following compounds transition metal is in oxidation state zero [NCERT 1982]
- (a)  $[Co(NH_3)_6]Cl_2$  (b)  $[Fe(H_2O)_6]SO_4$  (c)  $[Ni(CO)_4]$  (d)  $[Fe(H_2O)_3](OH)_2$
61. Oxidation number of osmium (Os) in  $OsO_4$  is [AIIMS 1999]
- (a) +4 (b) +6 (c) +7 (d) +8
62. The atomic number of an element which shows the oxidation state of +3 is [CPMT 1989, 94]
- (a) 13 (b) 32 (c) 33 (d) 17
63. The oxidation number of iron in the compound  $K_4[Fe(CN)_6]$  is [NCERT 1976; MNR 1986; AIIMS 2000]
- (a) +6 (b) +4 (c) +3 (d) +2

64. The brown ring complex compound is formulated as  $[Fe(H_2O)_5NO]SO_4$ . The oxidation state of iron is  
[EAMCET 1987; IIT 1987; MP PMT 1994; AIIMS 1997; DCE 2000]  
(a) 1 (b) 2  
(c) 3 (d) 0
65. Oxidation state of oxygen in  $F_2O$  is  
[BHU 1982; UPSEAT 2001; MH CET 2002]  
(a) +1 (b) +2  
(c) -1 (d) -2
66. Phosphorus has the oxidation state of +3 in  
[NCERT 1982; RPMT 1999]  
(a) Orthophosphoric acid (b) Phosphorus acid  
(c) Metaphosphoric acid (d) Pyrophosphoric acid
67. Oxidation number of P in  $Mg_2P_2O_7$  is  
[CPMT 1989; MP PMT 1995]  
(a) +3 (b) +2  
(c) +5 (d) -3
68. The oxidation state of nitrogen is highest in  
[MP PMT 2001; BHU 2002]  
(a)  $N_3H$  (b)  $NH_2OH$   
(c)  $N_2H_4$  (d)  $NH_3$
69. Oxidation number of P in  $KH_2PO_2$  is  
[CPMT 1987; MH CET 1999]  
(a) +1 (b) +3  
(c) +5 (d) -4
70. The most common oxidation state of an element is -2. The number of electrons present in its outermost shell is  
[BHU 1983; NCERT 1974; CPMT 1977]  
(a) 4 (b) 2  
(c) 6 (d) 8
71. Sulphur has lowest oxidation number in  
[EAMCET 1993]  
(a)  $H_2SO_3$  (b)  $SO_2$   
(c)  $H_2SO_4$  (d)  $H_2S$
72. The oxidation number and covalency of sulphur in the sulphur molecule ( $S_8$ ) are respectively  
[NCERT 1977]  
(a) 0 and 2 (b) 6 and 8  
(c) 0 and 8 (d) 6 and 2
73. In ferrous ammonium sulphate oxidation number of Fe is  
[CPMT 1988]  
(a) +3 (b) +2  
(c) +1 (d) -2
74. The oxidation number of nitrogen in  $NH_2OH$  is  
[NCERT 1981]  
(a) +1 (b) -1  
(c) -3 (d) -2
75. The oxidation number of phosphorus in  $Ba(H_2PO_2)_2$  is  
[Kurukshetra CEE 1998; DCE 2004]  
(a) -1 (b) +1  
(c) +2 (d) +3
76. A compound is in its low oxidation state. Then it will be  
[DCE 2001]  
(a) Highly acidic  
(b) Highly basic  
(c) Highest oxidising property  
(d) Half acidic, half basic
77. The oxidation number and the electronic configuration of sulphur in  $H_2SO_4$  is  
[KCET 2002]  
(a) +4;  $1s^2 2s^2 2p^6 3s^2$   
(b) +2;  $1s^2 2s^2 2p^6 3s^2 3p^2$   
(c) +3;  $1s^2 2s^2 2p^6 3s^2 3p^1$   
(d) +6;  $1s^2 2s^2 2p^6$
78. The oxidation number of Mn in  $KMnO_4$  is  
[CPMT 1982, 83; EAMCET 1992, 93; RPET 1999]  
(a) +7 (b) -7  
(c) +1 (d) -1
79. Oxidation number of As atoms in  $H_3AsO_4$  is  
[DPMT 2001]  
(a) -3 (b) +4  
(c) +6 (d) +5
80. In  $XeO_3$  and  $XeF_6$  the oxidation state of Xe is  
[MP PET 2003]  
(a) +4 (b) +6  
(c) +1 (d) +3
81. Oxidation number of carbon in  $CH_3 - Cl$  is  
[MP PET 2000]  
(a) -3 (b) -2  
(c) -1 (d) 0
82. The oxidation state of Cr in  $Cr_2O_7^{2-}$  is  
[BHU 2000; CPMT 2000]  
(a) 4 (b) -6  
(c) 6 (d) -2
83. Oxidation state of S in  $H_2SO_3$   
[RPET 2003]  
(a) +3 (b) +6  
(c) +4 (d) +2
84. Oxidation numbers of two Cl atoms in bleaching powder,  $CaOCl_2$  are  
(a) -1, -1 (b) +1, -1  
(c) +1, +1 (d) 0, -1
85. Select the compound in which chlorine is assigned the oxidation number +5  
[NCERT 1984, 94]  
(a)  $HClO_4$  (b)  $HClO_2$   
(c)  $HClO_3$  (d)  $HCl$
86. When  $KMnO_4$  is reduced with oxalic acid in acidic solution, the oxidation number of Mn changes from  
[MNR 1987; MP PET 2000; CBSE PMT 2000; UPSEAT 2000, 02; BHU 2003; AMU 2002]  
(a) 7 to 4 (b) 6 to 4  
(c) 7 to 2 (d) 4 to 2
87. Oxygen has oxidation states of +2 in the  
[NCERT 1973; DPMT 1983; MP PET 2000]  
(a)  $H_2O_2$  (b)  $CO_2$   
(c)  $H_2O$  (d)  $OF_2$
88. The element exhibiting most stable +2 oxidation state among the following is  
[IIT 1995]  
(a) Ag (b) Fe  
(c) Sn (d) Pb
89. Oxidation number of sulphur in  $S_2O_3^{2-}$  is  
[CPMT 1979]  
(a) -2 (b) +2  
(c) +6 (d) 0
90. Carbon has zero oxidation number in  
[Kurukshetra CEE 2002]

- (a)  $CO$  (b)  $CH_4$   
(c)  $CH_2Cl_2$  (d)  $CH_3Cl$
91. Oxidation state of oxygen atom in potassium superoxide is [MNR 1988; NCERT 1980]  
(a) 0 (b)  $-1$   
(c)  $-\frac{1}{2}$  (d)  $-2$
92. Oxidation number of  $S$  in  $S_2Cl_2$  is  
(a)  $+1$  (b)  $-1$   
(c)  $+6$  (d) 0
93. What is the oxidation number of sulphur in  $Na_2S_4O_6$  [AIIMS 1998; DCE 1999]  
(a)  $\frac{2}{3}$  (b)  $\frac{3}{2}$   
(c)  $\frac{3}{5}$  (d)  $\frac{5}{2}$
94. When  $CuSO_4$  reacts with  $KI$ , the oxidation number of  $Cu$  changes by [BHU 1997]  
(a) 0 (b)  $-1$   
(c) 1 (d) 2
95. The oxidation number of  $N$  in  $NH_4Cl$  is  
(a)  $+5$  (b)  $+3$   
(c)  $-5$  (d)  $-3$
96. In which reaction there is a change in valency [NCERT 1971; CPMT 1971]  
(a)  $2NO_2 \rightarrow N_2O_4$   
(b)  $2NO_2 + H_2O \rightarrow HNO_2 + HNO_3$   
(c)  $NH_4OH \rightarrow NH_4^+ + OH^-$   
(d)  $CaCO_3 \rightarrow CaO + CO_2$
97. Oxidation state of  $Fe$  in  $Fe_3O_4$  is [CBSE PMT 1999; AIIMS 2002]  
(a)  $\frac{3}{2}$  (b)  $\frac{4}{5}$   
(c)  $\frac{5}{4}$  (d)  $\frac{8}{3}$
98. Nitrogen show different oxidation states in the range [Kerala (Med.) 2003]  
(a) 0 to  $+5$  (b)  $-3$  to  $+5$   
(c)  $-5$  to  $+3$  (d)  $-3$  to  $+3$
99. Oxidation number of  $Mn$  in  $K_2MnO_4$  and  $MnSO_4$  are respectively [CPMT 1997]  
(a)  $+7, +2$  (b)  $+6, +2$   
(c)  $+5, +2$  (d)  $+2, +6$
100. Identify the element which can have highest oxidation numbers [AIIMS 1996]  
(a)  $N$  (b)  $O$   
(c)  $Cl$  (d)  $C$
101. What is the oxidation number of  $Co$  in  $[Co(NH_3)_4ClNO_2]$  [BHU 1999]  
(a)  $+2$  (b)  $+3$   
(c)  $+4$  (d)  $+5$
102. The oxidation number of nickel in  $K_4[Ni(CN)_4]$  is [JIPMER 1999]  
(a)  $-2$  (b)  $-1$   
(c)  $+2$  (d) 0
103. The oxidation number of fluorine in  $F_2O$  is [CPMT 1982; BHU 1982; EAMCET 1986]  
(a)  $-1$  (b)  $+1$   
(c)  $+2$  (d)  $-2$
104. Oxidation number of  $Fe$  in  $K_3[Fe(CN)_6]$  is [AMU 1988]  
(a)  $+2$  (b)  $+3$   
(c)  $+1$  (d)  $+4$
105. Oxidation number of  $N$  in  $NH_3$  is [CPMT 1979; Pb CET 2004]  
(a)  $-3$  (b)  $+3$   
(c) 0 (d)  $+5$
106. What is the net charge on ferrous ion [AFMC 2004]  
(a)  $+2$  (b)  $+3$   
(c)  $+4$  (d)  $+5$
107. Which of the following elements never show positive oxidation number [CPMT 2004]  
(a)  $O$  (b)  $Fe$   
(c)  $Ga$  (d)  $F$
108. The oxidation state shown by silicon when it combines with strongly electropositive metals is [MH CET 2004]  
(a)  $-2$  (b)  $-4$   
(c)  $+4$  (d)  $-2$
109. The oxidation number of sulphur in  $H_2S$  is [Pb. CET 2002]  
(a)  $-2$  (b)  $+3$   
(c)  $+2$  (d)  $-3$
110. Oxidation number of nitrogen in  $NaNO_2$  is [Pb. CET 2000]  
(a)  $+2$  (b)  $+3$   
(c)  $+4$  (d)  $-3$
111. Oxidation number of  $S$  in  $SO_4^{2-}$  [BCECE 2005]  
(a)  $+6$  (b)  $+3$   
(c)  $+2$  (d)  $-2$
112. The oxidation state of chromium in the final product formed by the reaction between  $KI$  and acidified potassium dichromate solution is [AIEEE 2005]  
(a)  $+4$  (b)  $+6$   
(c)  $+2$  (d)  $+3$
113. The oxidation state of  $I$  in  $IPO_4$  is [Orissa JEE 2005]  
(a)  $+1$  (b)  $+3$   
(c)  $+5$  (d)  $+7$

### Redox reaction and Method for balancing Redox reaction

1. The value of  $x$  in the partial redox equation  $MnO_4^- + 8H^+ + xe \rightleftharpoons Mn^{2+} + 4H_2O$  is  
(a) 5 (b) 3  
(c) 1 (d) 0
2.  $C_2H_6(g) + nO_2 \rightarrow CO_2(g) + H_2O(l)$   
In this equation, the ratio of the coefficients of  $CO_2$  and  $H_2O$  is [KCET 1992]  
(a) 1 : 1 (b) 2 : 3  
(c) 3 : 2 (d) 1 : 3
3. The number of electrons involved in the reduction of  $Cr_2O_7^{2-}$  in acidic solution to  $Cr^{3+}$  is [EAMCET 1983]  
(a) 0 (b) 2  
(c) 3 (d) 5
4. When  $KMnO_4$  acts as an oxidising agent and ultimately forms  $[MnO_4]^{-2}$ ,  $MnO_2$ ,  $Mn_2O_3$ ,  $Mn^{+2}$  then the number of electrons transferred in each case respectively is [AIEEE 2002]

- (a) 4, 3, 1, 5 (b) 1, 5, 3, 7  
(c) 1, 3, 4, 5 (d) 3, 5, 7, 1
5. Starch paper is used to test for the presence of [NCERT 1979]  
(a) Iodine (b) Oxidising agent  
(c) Iodide ion (d) Reducing agent
6. How many moles of  $K_2Cr_2O_7$  can be reduced by 1 mole of  $Sn^{2+}$  [MP PMT 2003]  
(a)  $1/3$  (b)  $1/6$   
(c)  $2/3$  (d) 1
7.  $2MnO_4^- + 5H_2O_2 + 6H^+ \rightarrow 2Z + 5O_2 + 8H_2O$ . In this reaction Z is [RPMT 2002]  
(a)  $Mn^{+2}$  (b)  $Mn^{+4}$   
(c)  $MnO_2$  (d)  $Mn$
8. What is 'A' in the following reaction  
 $2Fe^{3+}_{(aq)} + Sn^{2+}_{(aq)} \rightarrow 2Fe^{2+}_{(aq)} + A$  [MP PET 2003]  
(a)  $Sn^{3+}_{(aq)}$  (b)  $Sn^{4+}_{(aq)}$   
(c)  $Sn^{2+}_{(aq)}$  (d)  $Sn$
9. For the redox reaction  
 $MnO_4^- + C_2O_4^{2-} + H^+ \rightarrow Mn^{2+} + CO_2 + H_2O$   
the correct coefficients of the reactants for the balanced reaction are [IIT 1988, 92; BHU 1995; CPMT 1997; RPMT 1999; DCE 2000; MP PET 2003]
- | $MnO_4^-$ | $C_2O_4^{2-}$ | $H^+$ |
|-----------|---------------|-------|
| (a) 2     | 5             | 16    |
| (b) 16    | 5             | 2     |
| (c) 5     | 16            | 2     |
| (d) 2     | 16            | 5     |
10. Which of the following is a redox reaction [AIEEE 2002]  
(a)  $NaCl + KNO_3 \rightarrow NaNO_3 + KCl$   
(b)  $CaC_2O_4 + 2HCl \rightarrow CaCl_2 + H_2C_2O_4$   
(c)  $Mg(OH)_2 + 2NH_4Cl \rightarrow MgCl_2 + 2NH_4OH$   
(d)  $Zn + 2AgCN \rightarrow 2Ag + Zn(CN)_2$
11. Which of the following reaction is a redox reaction [MP PMT 2003]  
(a)  $P_2O_5 + 2H_2O \rightarrow H_4P_2O_7$   
(b)  $2AgNO_3 + BaCl_2 \rightarrow 2AgCl + Ba(NO_3)_2$   
(c)  $BaCl_2 + H_2SO_4 \rightarrow BaSO_4 + 2HCl$   
(d)  $Cu + 2AgNO_3 \rightarrow 2Ag + Cu(NO_3)_2$
12. Which of the following reactions involves oxidation-reduction [NCERT 1972; AFMC 2000; Pb. CET 2004; CPMT 2004]  
(a)  $NaBr + HCl \rightarrow NaCl + HBr$   
(b)  $HBr + AgNO_3 \rightarrow AgBr + HNO_3$   
(c)  $H_2 + Br_2 \rightarrow 2HBr$   
(d)  $2NaOH + H_2SO_4 \rightarrow Na_2SO_4 + 2H_2O$
13. Which of the following is the strongest oxidising agent [Pb. CET 2000]  
(a)  $BrO_3^- / Br^{2+}, E^\circ = +1.50$   
(b)  $Fe^{3+} / Fe^{2+}, E^\circ = +0.76$   
(c)  $MnO_4^- / Mn^{2+}, E^\circ = +1.52$   
(d)  $Cr_2O_7^{2-} / Cr^{3+}, E^\circ = +1.33$
14. In the balanced chemical reaction,  
 $IO_3^- + aI^- + bH^+ \rightarrow cH_2O + dI_2$   
a, b, c and d respectively correspond to [AIIMS 2005]  
(a) 5, 6, 3, 3 (b) 5, 3, 6, 3  
(c) 3, 5, 3, 6 (d) 5, 6, 5, 5
15. The number of moles of  $KMnO_4$  reduced by one mole of  $KI$  in alkaline medium is: [CBSE PMT 2005]  
(a) One fifth (b) five  
(c) One (d) Two

## Auto oxidation and Disproportionation

1. In the equation  $H_2S + 2HNO_3 \rightarrow 2H_2O + 2NO_2 + S$   
The equivalent weight of hydrogen sulphide is [BVP 2003]  
(a) 16 (b) 68  
(c) 34 (d) 17
2. 1.2 g of metal displace 1.12 litre hydrogen at normal temperature and pressure, equivalent weight of metal would be [DPMT 2001]  
(a) 24 (b) 12  
(c)  $1.2 \div 11.2$  (d)  $1.2 \times 11.2$
3. Which one of the following nitrates will leave behind a metal on strong heating [AIEEE 2003]  
(a) Ferric nitrate (b) Copper nitrate  
(c) Manganese nitrate (d) Silver nitrate
4. To prevent rancidification of food material, which of the following is added [CPMT 1996]  
(a) Reducing agent (b) Anti-oxidant  
(c) Oxidising agent (d) None of these
5. Prevention of corrosion of iron by zinc coating is called [MP PMT 1993; CPMT 2002]  
(a) Galvanization (b) Cathodic protection  
(c) Electrolysis (d) Photo-electrolysis
6. The metal used in galvanizing of iron is [MP PET 1985, 96]  
(a) Pb (b) Zn  
(c) Al (d) Sn
7. In which of the following reactions there is no change in valency [NCERT 1974; CPMT 1995]  
(a)  $4KClO_3 \rightarrow 3KClO_4 + KCl$   
(b)  $SO_2 + 2H_2S \rightarrow 2H_2O + 3S$   
(c)  $BaO_2 + H_2SO_4 \rightarrow BaSO_4 + H_2O_2$   
(d)  $2BaO + O_2 \rightarrow 2BaO_2$
8. The equivalent weight of phosphoric acid ( $H_3PO_4$ ) in the reaction  
 $NaOH + H_3PO_4 \rightarrow NaH_2PO_4 + H_2O$  is [AIIMS 1999]  
(a) 25 (b) 49  
(c) 59 (d) 98
9. What is the equivalent mass of  $IO_4^-$  when it is converted into  $I_2$  in acid medium [Kerala PMT 2004]



- (a)  $M/6$  (b)  $M/7$   
(c)  $M/5$  (d)  $M/4$   
(e) None of these
10. For decolourization of 1 mole of  $KMnO_4$ , the moles of  $H_2O_2$  required is [AIIMS 2004]  
(a)  $1/2$  (b)  $3/2$   
(c)  $5/2$  (d)  $7/2$
11. In the reaction  $I_2 + 2S_2O_3^{2-} \rightarrow 2I^- + S_4O_6^{2-}$  equivalent weight of iodine will be equal to [MP PET 2004]  
(a)  $1/2$  of molecular weight  
(b) Molecular weight  
(c)  $1/4$  of molecular weight  
(d) None
12. The equivalent weight of  $KIO_3$  in the reaction  $2Cr(OH)_3 + 4OH + KIO_3 \rightarrow 2CrO_4^{2-} + 5H_2O + KI$  is [MP PMT 2004]  
(a) Mole wt. (b)  $\frac{\text{Mol. wt.}}{6}$   
(c)  $\frac{\text{Mol. wt.}}{2}$  (d)  $\frac{\text{Mol. wt.}}{3}$
13. The product of oxidation of  $I^-$  with  $MnO_4^-$  in alkaline medium is [IIT-JEE Screening 2004]  
(a)  $IO_3^-$  (b)  $I_2$   
(c)  $IO^-$  (d)  $IO_4^-$
14. In alkaline medium  $ClO_2$  oxidize  $H_2O_2$  in  $O_2$  and reduced itself in  $Cl^-$  then how many mole of  $H_2O_2$  will oxidize by one mole of  $ClO_2$  [Kerala CET 2005]  
(a) 1.0 (b) 1.5  
(c) 2.5 (d) 3.5  
(e) 5.0
4. Of all the three common mineral acids, only sulphuric acid is found to be suitable for making the solution acidic because [Kurukshetra CEE 2002]  
(a) It does not react with  $KMnO_4$  or the reducing agent  
(b) Hydrochloric acid reacts with  $KMnO_4$   
(c) Nitric acid is an oxidising agent which reacts with reducing agent  
(d) All of the above are correct
5. For  $H_3PO_3$  and  $H_3PO_4$  the correct choice is [IIT Screening 2003]  
(a)  $H_3PO_3$  is dibasic and reducing  
(b)  $H_3PO_3$  is dibasic and non-reducing  
(c)  $H_3PO_4$  is tribasic and reducing  
(d)  $H_3PO_3$  is tribasic and non-reducing
6. Match List I with List II and select the correct answer using the codes given below the lists  
List I (Compound) List II (Oxidation state of N)  
(A)  $NO_2$  (1) +5  
(B)  $HNO$  (2) -3  
(C)  $NH_3$  (3) +4  
(D)  $N_2O_5$  (4) +1  
Codes :  
(a) A B C D  
2 3 4 1  
(b) A B C D  
3 1 2 4  
(c) A B C D  
3 4 2 1  
(d) A B C D  
2 3 1 4
7.  $M^{+3}$  ion loses  $3e^-$ . Its oxidation number will be [CPMT 2002]  
(a) 0 (b) +3  
(c) +6 (d) -3
8. In the reaction  $Zn + 2H^+ + 2Cl^- \rightarrow Zn^{2+} + 2Cl^- + H_2$ , the spectator ion is [AIIMS 2001]  
(a)  $Cl^-$  (b)  $Zn^{2+}$   
(c)  $H^+$  (d) All of these
9. The oxidation number of sulphur in  $H_2S_2O_7$  and iron in  $K_4Fe(CN)_6$  is respectively [AIIMS 2000]  
(a) +6 and +2 (b) +2 and +2  
(c) +8 and +2 (d) +6 and +4
10. Oxidation number of oxygen in potassium super oxide ( $KO_2$ ) is [UPSEAT 1999]  
(a) -2 (b) -1  
(c) -1/2 (d) -1/4
11. One mole of  $N_2H_4$  loses 10 mol of electrons to form a new compound Y. Assuming that all nitrogen appear in the new compound, what is the oxidation state of N in Y? (There is no change in the oxidation state of hydrogen) [IIT 1981; Pb. PMT 1998]  
(a) +3 (b) -3  
(c) -1 (d) +5

## Critical Thinking

### Objective Questions

1. In which of the following acid, which acid has oxidation reduction and complex formation properties [UPSEAT 2001]  
(a)  $HNO_3$  (b)  $H_2SO_4$   
(c)  $HCl$  (d)  $HNO_2$
2. The compound which could not act both as oxidising as well as reducing agent is [IIT Screening 1991]  
(a)  $SO_2$  (b)  $MnO_2$   
(c)  $Al_2O_3$  (d)  $CrO$
3.  $H_2S$  acts only as a reducing agent while  $SO_2$  can act both as a reducing and oxidizing agent because [AMU 1999]  
(a) S in  $H_2S$  has -2 oxidation state  
(b) S in  $SO_2$  has oxidation state +4  
(c) Hydrogen in  $H_2S$  more +ve than oxygen  
(d) Oxygen is more -ve in  $SO_2$
4. Of all the three common mineral acids, only sulphuric acid is found to be suitable for making the solution acidic because [Kurukshetra CEE 2002]  
(a) It does not react with  $KMnO_4$  or the reducing agent  
(b) Hydrochloric acid reacts with  $KMnO_4$   
(c) Nitric acid is an oxidising agent which reacts with reducing agent  
(d) All of the above are correct
5. For  $H_3PO_3$  and  $H_3PO_4$  the correct choice is [IIT Screening 2003]  
(a)  $H_3PO_3$  is dibasic and reducing  
(b)  $H_3PO_3$  is dibasic and non-reducing  
(c)  $H_3PO_4$  is tribasic and reducing  
(d)  $H_3PO_3$  is tribasic and non-reducing
6. Match List I with List II and select the correct answer using the codes given below the lists  
List I (Compound) List II (Oxidation state of N)  
(A)  $NO_2$  (1) +5  
(B)  $HNO$  (2) -3  
(C)  $NH_3$  (3) +4  
(D)  $N_2O_5$  (4) +1  
Codes :  
(a) A B C D  
2 3 4 1  
(b) A B C D  
3 1 2 4  
(c) A B C D  
3 4 2 1  
(d) A B C D  
2 3 1 4
7.  $M^{+3}$  ion loses  $3e^-$ . Its oxidation number will be [CPMT 2002]  
(a) 0 (b) +3  
(c) +6 (d) -3
8. In the reaction  $Zn + 2H^+ + 2Cl^- \rightarrow Zn^{2+} + 2Cl^- + H_2$ , the spectator ion is [AIIMS 2001]  
(a)  $Cl^-$  (b)  $Zn^{2+}$   
(c)  $H^+$  (d) All of these
9. The oxidation number of sulphur in  $H_2S_2O_7$  and iron in  $K_4Fe(CN)_6$  is respectively [AIIMS 2000]  
(a) +6 and +2 (b) +2 and +2  
(c) +8 and +2 (d) +6 and +4
10. Oxidation number of oxygen in potassium super oxide ( $KO_2$ ) is [UPSEAT 1999]  
(a) -2 (b) -1  
(c) -1/2 (d) -1/4
11. One mole of  $N_2H_4$  loses 10 mol of electrons to form a new compound Y. Assuming that all nitrogen appear in the new compound, what is the oxidation state of N in Y? (There is no change in the oxidation state of hydrogen) [IIT 1981; Pb. PMT 1998]  
(a) +3 (b) -3  
(c) -1 (d) +5

12. Amongst the following identify the species with an atom in + 6 oxidation state [IIT Screening 2000]
- (a)  $MnO_4^-$  (b)  $Cr(CN)_6^{3-}$   
(c)  $NiF_6^{2-}$  (d)  $CrO_2Cl_2$
13. In which of the following compounds, is the oxidation number of iodine is fractional [BVP 2003]
- (a)  $IF_3$  (b)  $IF_2$   
(c)  $I_3^-$  (d)  $IF_7$
14. The compound  $YBa_2Cu_3O_7$  which shows superconductivity has copper in oxidation state ..... Assume that the rare earth element Yttrium is in its usual +3 oxidation state [IIT 1994]
- (a)  $3/7$  (b)  $7/3$   
(c)  $3$  (d)  $7$
15. The oxidation number of sulphur in  $S_8, S_2F_2, H_2S$  respectively, are [IIT 1999]
- (a) 0, +1 and -2 (b) +2, +1 and -2  
(c) 0, +1 and +2 (d) -2, +1 and -2
16. Which one of the following reactions is not an example of redox reaction [Kurukshetra CEE 1998]
- (a)  $Cl_2 + 2H_2O + SO_2 \rightarrow 4H^+ + SO_4^{2-} + 2Cl^-$   
(b)  $Cu^{++} + Zn \rightarrow Zn^{++} + Cu$   
(c)  $2H_2 + O_2 \rightarrow 2H_2O$   
(d)  $HCl + H_2O \rightarrow H_3O^+ + Cl^-$
17. For the reactions,  $C + O_2 \rightarrow CO_2; \Delta H = -393J$   
 $2Zn + O_2 \rightarrow 2ZnO; \Delta H = -412J$  [AIEEE 2002]
- (a) Carbon can oxidise Zn  
(b) Oxidation of carbon is not feasible  
(c) Oxidation of Zn is not feasible  
(d) Zn can oxidise carbon
18. In the reaction  $B_2H_6 + 2KOH + 2X \rightarrow 2Y + 6H_2$ , X and Y are respectively [EAMCET 2003]
- (a)  $H_2, H_3BO_3$  (b)  $HCl, KBO_3$   
(c)  $H_2O, KBO_3$  (d)  $H_2O, KBO_2$
19. In a balanced equation  $H_2SO_4 + xHI \rightarrow H_2S + yI_2 + zH_2O$ , the values of x, y, z are [EAMCET 2003]
- (a)  $x = 3, y = 5, z = 2$   
(b)  $x = 4, y = 8, z = 5$   
(c)  $x = 8, y = 4, z = 4$   
(d)  $x = 5, y = 3, z = 4$
20. Which of the following can act as an acid and as a base [AMU 1999]
- (a)  $HClO_3^-$  (b)  $H_2PO_4^-$   
(c)  $HS^-$  (d) All of these
21.  $MnO_4^{2-}$  (1 mole) in neutral aqueous medium is disproportionate to [AIIMS 2003]
- (a)  $2/3$  mole of  $MnO_4^-$  and  $1/3$  mole of  $MnO_2$   
(b)  $1/3$  mole of  $MnO_4^-$  and  $2/3$  mole of  $MnO_2$   
(c)  $1/3$  mole of  $Mn_2O_7$  and  $1/3$  mole of  $MnO_2$   
(d)  $2/3$  mole of  $Mn_2O_7$  and  $1/3$  mole of  $MnO_2$
22. The conductivity of a saturated solution of  $BaSO_4$  is  $3.06 \times 10^{-6} \text{ ohm}^{-1} \text{ cm}^{-1}$  and its equivalent conductance is  $1.53 \text{ ohm}^{-1} \text{ cm}^{-1} \text{ equivalent}^{-1}$ . The  $K_{sp}$  of the  $BaSO_4$  will be [KCET 1996]
- (a)  $4 \times 10^{-12}$  (b)  $2.5 \times 10^{-9}$   
(c)  $2.5 \times 10^{-13}$  (d)  $4 \times 10^{-6}$
23. When  $MnO_2$  is fused with  $KOH$ , a coloured compound is formed, the product and its colour is [IIT Screening 2003]
- (a)  $K_2MnO_4$ , purple green  
(b)  $KMnO_4$ , purple  
(c)  $Mn_2O_3$ , brown  
(d)  $Mn_3O_4$  black

## Assertion & Reason

For AIIMS Aspirants

Read the assertion and reason carefully to mark the correct option out of the options given below :

- (a) If both assertion and reason are true and the reason is the correct explanation of the assertion.  
(b) If both assertion and reason are true but reason is not the correct explanation of the assertion.  
(c) If assertion is true but reason is false.  
(d) If the assertion and reason both are false.  
(e) If assertion is false but reason is true.

1. Assertion :  $SO_2$  and  $Cl_2$  both are bleaching agents.  
Reason : Both are reducing agents. [AIIMS 1995]
2. Assertion : Fluorine exists only in -1 oxidation state.  
Reason : Fluorine has  $2s^2 2p^5$  configuration. [AIIMS 2001]
3. Assertion : Stannous chloride is a powerful oxidising agent which oxidises mercuric chloride to mercury.  
Reason : Stannous chloride gives grey precipitate with mercuric chloride, but stannic chloride does not do so. [AIIMS 2002]
4. Assertion :  $HClO_4$  is a stronger acid than  $HClO_3$ .  
Reason : Oxidation state of Cl in  $HClO_4$  is +VII and in  $HClO_3$  +V. [AIIMS 2004]
5. Assertion : In a reaction  $Zn(s) + CuSO_4(aq) \rightarrow ZnSO_4(aq) + Cu(s)$ , Zn is a reductant but itself get oxidized.

- Reason : In a redox reaction, oxidant is reduced by accepting electrons and reductant is oxidized by losing electrons.
6. Assertion : Oxidation number of carbon in  $CH_2O$  is zero.
- Reason :  $CH_2O$  formaldehyde, is a covalent compound.
7. Assertion : The oxidation numbers are artificial, they are useful as a 'book-keeping' device of electrons in reactions.
- Reason : The oxidation numbers do not usually represent real charges on atoms, they are simply conventions that indicate what the maximum charge could possibly be on an atom in a molecule.
8. Assertion :  $H_2SO_4$  cannot act as reducing agent.
- Reason : Sulphur cannot increase its oxidation number beyond + 6.
9. Assertion : Equivalent weight of  $NH_3$  in the reaction  $N_2 \rightarrow NH_3$  is 17/3 while that of  $N_2$  is 28/6.
- Reason : Equivalent weight  

$$= \frac{\text{Molecular weight}}{\text{number of } e^- \text{ lost or gained}}$$

26	b	27	b	28	a	29	a	30	a
31	c	32	d	33	a	34	a	35	b
36	a	37	d	38	a	39	d	40	d
41	b	42	b	43	c	44	a	45	d
46	b	47	d	48	d	49	a	50	c
51	c	52	b	53	b	54	c	55	b
56	b	57	c	58	a	59	a	60	c
61	d	62	a	63	d	64	b	65	b
66	b	67	c	68	a	69	a	70	c
71	d	72	a	73	b	74	b	75	b
76	c	77	d	78	a	79	d	80	b
81	b	82	c	83	c	84	b	85	c
86	c	87	d	88	d	89	b	90	c
91	c	92	a	93	d	94	c	95	d
96	b	97	d	98	b	99	b	100	c
101	a	102	d	103	a	104	b	105	a
106	a	107	d	108	b	109	a	110	b
111	a	112	d	113	b				

# Answers

## Oxidation, Reduction

1	b	2	b	3	c	4	c	5	c
6	a	7	b	8	b	9	a	10	c
11	b	12	a	13	b	14	b	15	b
16	a	17	a	18	b	19	c	20	b
21	a	22	c	23	b	24	b	25	b
26	c	27	c	28	d	29	a	30	a
31	a	32	a	33	a	34	d	35	b
36	d	37	d						

## Oxidizing and Reducing agent

1	c	2	a	3	b	4	a	5	d
6	b	7	c	8	b	9	a	10	b
11	c	12	d	13	a	14	d	15	a
16	b	17	b	18	bd	19	b	20	d
21	a	22	b	23	d	24	d	25	b
26	d	27	d	28	c	29	a	30	d
31	d	32	c						

## Oxidation number and Oxidation state

1	d	2	b	3	b	4	b	5	d
6	b	7	c	8	c	9	c	10	a
11	c	12	a	13	d	14	b	15	a
16	d	17	d	18	d	19	a	20	c
21	d	22	b	23	c	24	d	25	a

## Redox reaction and Method for balancing Redox reaction

1	a	2	b	3	c	4	c	5	a
6	a	7	a	8	b	9	a	10	d
11	d	12	c	13	c	14	a	15	d

## Auto oxidation and Disproportionation

1	d	2	a	3	d	4	b	5	a
6	b	7	c	8	d	9	b	10	c
11	a	12	d	13	a	14	c		

## Critical Thinking Questions




1	d	2	c	3	a,b	4	d	5	a
6	c	7	c	8	a	9	a	10	c
11	a	12	d	13	c	14	b	15	a
16	d	17	d	18	d	19	c	20	d
21	a	22	d	23	a				

## Assertion & Reason

1	c	2	b	3	e	4	b	5	a
6	b	7	a	8	a	9	a		

# Answers and Solutions

### Oxidation, Reduction



It shows oxidation and reduction (Redox) properties.

