

Electrochemistry

SET Self Evaluation Test -12

1. The mass of copper deposited from a solution of CuSO_4 by passage of 5 A current for 965 second is (Mol. wt. of Copper = 63.5) [AIIMS 2001]
- (a) 15.875 g (b) 1.5875 g
(c) 4825 g (d) 96500 g
2. The current in a given wire is 1.8 A. The number of coulombs that flow in 1.36 minutes will be [AIIMS 2001]
- (a) 100 C (b) 147 C
(c) 247 C (d) 347 C
3. A solution of a salt of a metal was electrolysed for 150 minutes with a current of 0.15 amperes. The weight of metal deposited was 0.783 gm. The equivalent weight of the metal is [AFMC 2001]
- (a) 55.97 gm (b) 65.97 gm
(c) 75.97 gm (d) 85.97 gm
4. The resistance of 0.01N NaCl solution at 25°C is 200 Ω . Cell constant of conductivity cell is 1 cm. The equivalent conductance is
- (a) $5 \times 10^2 \Omega^{-1} \text{cm}^2 \text{eq}^{-1}$ (b) $6 \times 10^3 \Omega^{-1} \text{cm}^2 \text{eq}^{-1}$
(c) $7 \times 10^4 \Omega^{-1} \text{cm}^2 \text{eq}^{-1}$ (d) $8 \times 10^5 \Omega^{-1} \text{cm}^2 \text{eq}^{-1}$
5. Which of the following reaction is possible at anode [AIEEE 2002]
- (a) $2\text{Cr}^{3+} + 7\text{H}_2\text{O} \rightarrow \text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+$
(b) $\text{F}_2 \rightarrow 2\text{F}^-$
(c) $\frac{1}{2}\text{O}_2 + 2\text{H}^+ \rightarrow \text{H}_2\text{O}$
(d) None of these
6. What is the standard cell potential for the cell
 $\text{Zn} / \text{Zn}^{2+} (1\text{M}) || \text{Cu}^{2+} (1\text{M}) / \text{Cu}$
 E° for $\text{Zn} / \text{Zn}^{2+} (1\text{M}) = -0.76 \text{ V}$ & $\text{Cu}^{2+} / \text{Cu} = +0.34 \text{ V}$ [AIIMS 1980]
- (a) $-0.76 + (-0.34) = -0.42 \text{ V}$
(b) $-0.34 + 0.76 = +0.42 \text{ V}$
(c) $0.34 - (-0.76) = 1.10 \text{ V}$
(d) $-0.76 - (+0.34) = -1.10 \text{ V}$
7. Normal aluminium electrode coupled with normal hydrogen electrode gives an emf of 1.66 volts. So the standard electrode potential of aluminium is [KCET 1987]
- (a) -1.66 V (b) $+1.66 \text{ V}$
(c) -0.83 V (d) $+0.83 \text{ V}$
8. Which one among the following is the strongest reducing agent
- $\text{Fe}^{2+} + 2\text{e}^- \rightarrow \text{Fe} (-0.44 \text{ V})$
 $\text{Ni}^{2+} + 2\text{e}^- \rightarrow \text{Ni} (-0.25 \text{ V})$
 $\text{Sn}^{2+} + 2\text{e}^- \rightarrow \text{Sn} (-0.14 \text{ V})$
 $\text{Fe}^{3+} + \text{e}^- \rightarrow \text{Fe}^{2+} (-0.77 \text{ V})$ [BHU 1998]
- (a) Fe (b) Fe^{2+}
(c) Ni (d) Sn
9. The [CBSE PMT 1999] cell reaction of the galvanic cell
 $\text{Cu}_{(\text{s})} | \text{Cu}^{2+}_{(\text{aq})} || \text{Hg}^{2+}_{(\text{aq})} | \text{Hg}_{(\text{l})}$ is [EAMCET 2003]
- (a) $\text{Hg} + \text{Cu}^{2+} \rightarrow \text{Hg}^{2+} + \text{Cu}$
(b) $\text{Hg} + \text{Cu}^{2+} \rightarrow \text{Cu}^+ + \text{Hg}^+$
(c) $\text{Cu} + \text{Hg} \rightarrow \text{CuHg}$
(d) $\text{Cu} + \text{Hg}^{2+} \rightarrow \text{Cu}^{2+} + \text{Hg}$
10. The specific conductivity of N/10 KCl solution at 20°C is $0.0212 \text{ ohm}^{-1} \text{cm}^{-1}$ and the resistance of cell containing this solution at 20°C is 55 ohm. The cell constant is [AIIMS 1999]
- (a) 1.166 cm^{-1}
(b) 2.173 cm^{-1}
(c) 3.324 cm^{-1}
(d) 4.616 cm^{-1}
11. The oxide which is not reduced by hydrogen is [JIPMER 1999]
- (a) Ag_2O (b) K_2O
(c) Fe_2O_3 (d) P_4O_{10}

1. (b) Current (I) = 5A and time (t) = 965 sec.
We know that equivalent weight of copper

$$= \frac{\text{Molecular weight}}{\text{Valency}} = \frac{63.5}{2}$$
 and quantity of electricity passed in coulomb = current \times time = $5 \times 965 = 4825 C$. Since 96500 coulombs will deposit $\frac{63.5}{2} g$ of copper therefore 4825 coulombs will deposit

$$= \frac{63.5 \times 4825}{96500 \times 2} = 1.5875 g.$$
2. (b) $Q = I \times t$; $1.8 \times 1.36 \times 60 = 147 C$.
3. (a) Time (t) = 150 min = 9000 sec
Current (I) = 0.15 A
Weight of metal (w) = 0.783 g.
We know $Q = I \times t = 0.15 \times 9000 = 1350 C$. Since 1350 C of electricity will deposited 0.783 g of metal, so, 96500 C of electricity will deposited $\frac{0.783 \times 96500}{1350} = 55.97 g$.
4. (a) $\lambda = k \times V = \frac{1}{R} \times \frac{l}{a} \times V = \frac{1}{200} \times 1 \times 10,000$

$$= 5 \times 10^2 \Omega^{-1} cm^2 eq.^{-1}$$
5. (a) Oxidation always occurs at anode.
6. (c) $E^o = E_{\text{cathode}} - E_{\text{anode}}$

$$E^o = 0.34 - (-0.76); E^o = 1.10 \text{ volt}.$$
7. (a) $E_{\text{cell}}^o = 1.66 = E_{H^+/H_2}^o - E_{Al^{3+}/Al}^o$

$$= 0 - E_{Al^{3+}/Al}^o \text{ or } E_{Al^{3+}/Al}^o = -1.66 V.$$
8. (a) The reduction potential of Fe is very high, so it is a strongest reducing agent.
9. (b) $Cu_{(s)} | Cu_{(Ag)}^{2+} || Hg_{(Ag)}^{2+} | Hg_{(l)}$
 anode oxidation cathode reduction
- $$\begin{array}{c}
 \text{Reduction} \\
 \text{Cu} + Hg^{2+} \rightarrow Cu^{2+} + Hg \\
 \text{Oxidation}
 \end{array}$$
10. (a) $K = \frac{1}{R} \times \text{cell constant}$

$$= K \times R = 0.0212 \times 55 = 1.166 cm^{-1}.$$
11. (b) On the basis of electrochemical series K_2O is not reduced by hydrogen.
