

# Tips & Tricks


Badecker reaction: This reaction involves the following chemical change.

 $Na_2[Fe(CN)_5NO] + Na_2SO_3 \rightarrow Na_4[Fe(CN)_5(NO.SO_3)]$ 

 $\angle$  Everitt's salt: It is  $K_2[Fe(CN)_6]$  obtained by reduction of prussian blue.

Masking: Masking is the process in which a substance without physical separation of it is so transformed that is does not enter into a particular reaction e.g., masking of  $Cu^{2+}$  by  $CN^-$  ion.

Macrocyclic effect: This term refers to the greater thermodynamic stability of a complex with a cyclic polydentate ligand when compared to the complex formed with a non-cyclic ligand. e.g., Zn(II) complex with ligand;



**E** Prussian blue and Turnbull's blue is pot. ferric ferrocyanide. However colour of Turnbull's blue is less intense than prussian blue. Decrease in colour is due to the presence in it of a white compound of the formula  $K_2\{Fe[Fe(CN)_6]\}$  named as potassium ferrous ferrocyanide.

Crystal field theory explains why certain geometries are more favoured than other by certain metals in terms of crystal field stablization energies.

The colour of complexes are explained in terms of electronic transitions between the various *d* orbitals of different energies.

 $m{\mathscr{L}}$  In octahedral complexes, the energy of the  $d_{r^2-v^2}$  and  $d_{r^2-v^2}$ orbitals increases much more than the energy of the  $d_{xy}, d_{yz}, d_{xz}$ orbitals. The potential energy increases as a result of repulsive interactions. The greater the repulsion, greater is the increases in energy.

Ligands with larger groups from unstable rings than the ligands with smaller groups. This is ascribable to steric hinderance produced by a larger group.

Men Ca or Mg forms complexes with EDTA, the pH of the solution decreases.

The copper sulphate solution e.g., turns deep blue when excess of ammonia is added. This is also due to the formation of [Cu(NH)]SO

Co-ordination compound with a general formula MA, MAB or MABCD do not show any geometric isomerism.

K Flexidentate character: polydentate ligand are said to have flexidentate character if they do not use all its donar atoms to get coordinated to the metal ion e.g., EDTA generally act as a hexadentate ligand but it can also act as a pentadentate and tetradentate ligand.

 Ø Octahedral complex of the type [M(ABCDEF)] have 15 different geometrical isomers with a pair of enantiomers. Although a few geometrical isomer have been prepared but none has been resolved e.g., [Pt(Py)(NH)(NO)(Cl)(Br)I].

# Ordinary Thinking

#### **Objective Questions**

#### **Basic Terms**

1.	ln	$K_{A}$	Fe	CN	)6
----	----	---------	----	----	----

- (a) (CN) are linked with primary valency
- (CN) are linked with secondary valency
- K are linked with secondary valency
- (d) K are linked with non-ionic valency
- The co-ordination number of copper in cuprammonium sulphate is (b) 6
  - (a) 2
    - (d) 4
- (c) 4 Which of the following acts as a bidentate ligand in complex 3. formation
  - (a) Acetate
- (b) Oxalate
- (c) Thiocyanate
- (d) **EDTA**
- The co-ordination number cobalt complex  $[Co(en), Br, ]Cl_2$  is
  - (a) 2

(b) 6

(c) 5

(d) 4

Which of the following ligands forms a chelate

[MP PET/PMT 1998]

- (a) Acetate
- (b) Oxalate
- (c) Cyanide
- (d) Ammonia
- According to Werner's theory 6.

[MP PMT 2000, 02]

- (a) Primary valency can be ionized
- (b) Secondary valency can be ionized
- (c) Primary and secondary valencies both cannot be ionized
- (d) Only primary valency cannot be ionized
- Which of the following is not true for ligand-metal complex

[MP PET 1993]

- (a) Larger the ligand, the more stable is the metal-ligand bond
- Highly charged ligand forms strong bond
- Larger the permanent dipole moment of ligand, the more stable is the bond
- Greater the ionization potential of central metal, the stronger is the bond
- What is the co-ordination number of the metal in  $|Co(en)_2 Cl_2|^{\dagger}$

7.

(b) 5

- (c) 6
- (d) 3
- Bidentate ligand is
  - (a)  $CN^-$
- (b) Ethylene diammine (en)
- (d) EDTA
- The coordination number of Pt in  $\left[Pt(NH_3)_4 Cl_2\right]^{++}$  ion is 10.

[MP PET 1995]

- (a) 2 (c) 6
- (b) 4 (d)
- Which is the example of hexadentate ligand
  - (a) 2, 2—dipyridyl
  - (b) Dimethyl glyoxime
  - (c) Aminodiacetate ion
  - (d) Ethylene diammine tetra acetate ion [EDTA]
- 12. The coordination number of a metal in coordination compounds is [MP PET 199
  - (a) Same as primary valency

IINIV	ERSAL		
	908 Co-ordination Chemistry		
	(b) Sum of primary and secondary valencies		(c) Double salt (d) Complex salt
	(c) Same as secondary valency	25.	A monodentate ligand has
13.	(d) None of these Ligand in a complex salt are [KCET 1992]		(a) One co-ordinate site (b) Two co-ordinate sites
13.	(a) Anions linked by coordinate bonds to a central metal atom or		(c) Any number of co-ordinate sites
	ion		(d) No capacity to co-ordinate
	(b) Cations linked by coordinate bonds to a central metal atom or	26.	EDTA has coordination number [AFMC 2004]
	(c) Molecules linked by coordinate bonds to a central metal atom		(a) 3 (b) 4
	(c) Molecules linked by coordinate bonds to a central metal atom or ion	~=	(c) 5 (d) 6
	(d) lons or molecules linked by coordinate bonds to a central metal atom or ion	27.	Among the properties (a) reducing (b) oxidising (c) complexing, the set of properties shown by $CN^-$ ion towards metal species is
14.	A group of atoms can function as a ligand only when		(a) c, a (b) b, c
	[KCET 1989; DCE 1999; MP PMT 2000]		(c) a, b (d) a, b, c
	<ul><li>(a) It is a small molecule</li><li>(b) It has an unshared electron pair</li></ul>	28.	That ion or molecule which forms a complex compound with transitional metal ion is called
	(c) It is a negatively charged ion		(a) Recipient (b) Ligand
	(d) It is a positively charged ion		(c) Coordinate ion (d) No special name
15.	Which of the following complexes show six coordination number [RPET	20009.	Coordination number of $Zn$ in $ZnS$ (zinc blende) is  [Orissa JEE 2004]
	(a) $[Zn(CN)_4]^{2-}$ (b) $[Cr(H_2O)_6]^{3+}$		(a) 6 (b) 4
	(c) $[Cu(CN)_4]^{2-}$ (d) $[Ni(NH_3)_4]^{2+}$		(c) 8 (d) 12
16.	The number of ions formed when cuprammonium sulphate is dissolved in water is [KCET 1993]	30.	Wilkinson's catalyst used as a homogeneous catalyst in the hydrogenation of alkenes contains
	(a) 1 (b) 2		(a) Iron (b) Aluminium
	(c) 4 (d) Zero		(c) Rhodium (d) Cobalt
17.	The coordination number of $\mathit{Cu}$ in complex $\left[\mathit{Cu}(H_2O)_4\right]^{\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!$	31. [	Given the molecular formula of the hexa coordinated complexes (A) $MP$ $COCl_3.5$ $NH_3$ (B) $CoCl_3.5$ $NH_3$ (C) $CoCl_3.4$ $NH_3$ . If the
	(a) 4 (b) 3		number of co-ordinated $NH_3$ molecules in $A$ , $B$ and $C$ respectively
	(c) 2 (d) 1		are 6, 5 and 4, the primary valency in (A), (B) and (C) are:[DCE 2003]
18.	The primary valence of the metal ion in the co-ordination compound $K_2[Ni(CN)_4]$ is		(a) 6, 5, 4 (b) 3, 2, 1 (c) 0, 1, 2 (d) 3, 3, 3
	(a) Four (b) Zero	32.	Generally, a group of atoms can function as a ligand if
	(c) Two (d) Six		[MP PET 1996]
19.	The metal which does not form a polynuclear carbonyl is		(a) They are positively charged ions
	(a) $Mn$ (b) $Co$		<ul><li>(b) They are free radicals</li><li>(c) They are either neutral molecules or negatively charged ions</li></ul>
	(c) $Cr$ (d) $Fe$		(d) None of these
20.	Which one of the following forms with an excess of $\ensuremath{\mathit{CN}^-}$ (Cyanide)	33.	The ligand in potassium ferricyanide is
	a complex having coordination number two		(a) $K^+$ (b) $CN^-$
	[AllMS 2004]		(c) $Fe^{3+}$ (d) $(CN)_6$
	(a) $Cu^+$ (b) $Ag^+$	34.	Co-ordination number of aluminum is [MHCET 2004]
	(c) $Ni^{2+}$ (d) $Fe^{2+}$	0	(a) 8 (b) 6
21.	According to Lewis the ligands are [MP PMT 2002]		(c) 12 (d) 4
	(a) Acidic in nature	35.	In $K_4 Fe(CN)_6$ , Fe is in the form of
	(b) Basic in nature (c) Neither acidic nor basic		(a) An atom (b) An ion
	(c) Neither acidic nor basic (d) Some are acidic and others are basic		(c) Cationic complex (d) Anionic complex
22.	The coordination number of a central metal atom in a complex is determined by  [AIEEE 2004]	36.	Which of the following ligands is expected to bidentate [CBSE PMT 1994]
	(a) The number of ligands around a metal ion bonded by sigma		(a) $Br$ (b) $C_2 O_4^{2-}$
	and pi-bonds both.		(c) $CH_3NH_2$ (d) $CH_3C \equiv N$
	<ul><li>(b) The number around a metal ion bonded by pi-bonds</li><li>(c) The number of ligands around a metal ion bonded by sigma bonds</li></ul>	37.	In the compound lithium tetrahydroaluminate, the ligand is [AlIMS 1997]
	(d) The number of only anionic ligands bonded to the metal ion		(a) $H^+$ (b) $H^-$

(c) H[MP PET 1989]

(a) Potassium ferrocyanide

(c) Potassium ferricyanide

(b) Ferrous ammonium sulphate

 $(d) \quad Tetrammine\ copper\ (11)\ sulphate$ 

Which of the following is the odd one out

38.

[AFMC 2000]

(d) None of these

[MP PET 1996]

In the extraction of which of the following, complex ion forms

(b) Ag

(d) Na

(b) Mixed salt

23.

24.

(a) *Cu* 

(c) Fe

(a) Normal salt

Potassium ferrocyanide is a



[UPSEAT 2000, 01]

39.	The basic ligand is		(d) Copper hydroxide is an amphoteric substance
	(a) $NH_3$ (b) $CN^-$	53	53. Zeigler—Natta catalyst is used for which type of reaction
			(a) Hydrogenation (b) Polymerization
40	(c) $F^-$ (d) All The negative ligand is		(c) Oxidation (d) Reduction
40.	(a) Aqua (b) Sulphato		<ol> <li>Which of the following is not considered as an organometallic compound. [AIIMS 2004]</li> </ol>
	(c) Carboxyl (d) Nitro soc		(a) Cis-platia (b) Ferrocene
41.	Which has yellow colour		(c) Zeise's salt (d) Gringard reagent
	(a) Potassium cobaltinitrite	55	55. Which one is organometallic compound[MP PMT 2004]
	(b) Potassium hexanitro cobaltate (III)		(a) Lithium methoxide (b) Lithium dimethyl amide
	(c) Fischer's salt		(c) Lithium acetate (d) Methyl lithium
40	(d) All the above	[MP PMT 2003]	<b>56.</b> An aqueous solution of potash alum gives [UPSEAT 2004]
42.	Ligands, in complex compounds	[MP PM1 2003]	(a) Two types of ions (b) Only one type of ion
	(a) Accept $e^-$ -pair		(c) Four types of ions (d) Three types of ions
	(b) Donate $e^-$ -pair	57	<b>67.</b> Carnallite in solution in $H_2O$ , shows the properties of
	(c) Neither accept $e^{-}$ -pair nor donate		[DCE 2003]
	(d) All of these happen		(a) $K^+, Mg^{2+}, Cl^-$ (b) $K^+, Cl^-, SO_4^{2-}, Br^-$
43.	Which of the following is a common donor atom		
	()	[BHU 2001]	(c) $K^+, Mg^{2+}, CO_3^{2-}$ (d) $K^+, Mg^{2+}, Cl^-, Br^-$
	(a) Arsenic (b) Nitrogen	50	<b>58.</b> What is the co-ordination number of cobalt in $Co(NH_3)_3Cl_3$
44	(c) Oxygen (d) Both (b) Trunbull's blue is a compound	and (c)	(a) 3 (b) 4
44.	Trumbun's blue is a compound	[KCET 1993]	(c) 5 (d) 6
	(a) Ferricyanide (b) Ferrous		59. The formula of alum is [Pb. CET 2002]
	(c) Ferrous cyanide (d) Ferriferro	•	(a) $K_2SO_4.Al_2(SO_4)_3.24H_2O$
45.	Tollen's reagent is	[KCET 1990]	2 1 2 1 1 2
	(a) $[Ag(NH_3)_2]^+$ (b) $Ag_2O$		(b) $K_4[Fe(CN)_6]$
	F ( ) 70		(c) $K_2SO_4.Al_2(SO_4)_3.6H_2O$
	(c) $[Cu(OH)_4]^{2-}$ (d) $Cu_2O$		(d) $Na_2CO_3.10H_2O$
46.	Finely divided iron combines with CO to give	60	<b>50.</b> Number of ions present in $K_4[Fe(CN)_6]$ [Pb. CET 2000]
		[UPSEAT 2002]	
	(a) $Fe(CO)_5$ (b) $Fe_2(CO)_5$	O) <sub>9</sub>	(a) 2 (b) 10 (c) 3 (d) 5
	(c) $Fe_2(CO)_{12}$ (d) $Fe(CO)_{12}$	)6	· · ·
47.	In a complex, the highest possible coordination i	61 number is	
••	(a) 6 (b) 12		[DCE 2002]
	(c) 4 (d) 8		(a) $Mg-I$ bond (b) $C-I$ bond
48.	The number of neutral molecules or negative gr	oups attached to the	(c) $C - Mg$ bond (d) $C - H$ bond
	central metal atom in a complex ion is called	62	<b>52.</b> What is the EAN of nickel in $Ni(CO)_4$ [BVP 2003]
		atomic number	
40	(c) Coordination number (d) Primary EDTA combines with cations to form	valency	(a) 34 (b) 35 (c) 32 (d) 36
49.	() - 1		(t) 32 (d) 30
	(a) lon-exchange resins (b) Chelates (c) Clathrates (d) Polymers		Namonalatura Ovidation atota and EAN number
50.	An example of a double salt is	[MP PET 2001]	Nomenclature, Oxidation state and EAN number
	(a) Bleaching powder (b) Hypo	-	
	(c) $K_4[Fe(CN)_6]$ (d) Potash a	1. Ium	. How many ions are produced in aqueous solution of $[Co(H_2O)_6]Cl_2$ [RPMT 2002]
51.	In complex compounds, metal ligand bond is		
J	(a) Coordinate bond (b) Hydroge	n bond	(a) 2 (b) 3
	(c) Ionic bond (d) Covalent		(c) 4 (d) 6
<b>5</b> 0	Ammonia forms the complex ion $[Cu(NH_3)_4]$	2	1UPAC name of $[Pt(NH_3)_3(Br)(NO_2)Cl]Cl$ is
52.	Ammonia forms the complex ion $[Cu(NH_3)_4]$ in alkaline solutions but not in acidic solution.		[CBSE PMT 1998]
	for it[AIEEE 2003]	windt is the reason	(a) Triamminechlorobromonitroplatinum (IV) chloride
	(a) In acidic solutions hydration protects coppe	er ions	(b) Triamminebromonitrochloroplatinum (IV) chloride
	(b) In acidic solutions protons coordinate with		(c) Triamminebromochloronitroplatinum (IV) chloride
	forming $NH_4^+$ ions and $NH_3$ molecules		(d) Triamminenitrochlorobromoplatinum (IV) chloride
	(c) In alkaline solutions insoluble $Cu(OH)_2$	is precipitated which	ondation state of incrogen is incorrectly given for

is soluble in excess of any alkali  $\,$ 



4.

5.

6.

7.

8.

9.

10.

11.

12.

# UNIVERSAL SELF SCORER 910 Co-ordination Chemistry

Compound Oxidation state		(d) Example of weak ligand field for $d^5$ configuration
(a) $[Co(NH_3)_5 Cl]Cl_2$ 0		$=\left[Mn^{II}F_{6}\right]^{-4}$
(b) $NH_2OH$ -1	13.	The oxidation number of chromium in sodium tetrafluoro
(c) $(N_2H_5)_2SO_4$ +2		oxochromate complex is (a) 11 (b) 1V
(d) $Mg_3N_2$ -3		(a) II (b) IV (c) VI (d) III
The formula of dichloro <i>bis</i> (urea) copper (11) is	14.	The IUPAC name of $K_4[Fe(CN)_6]$ is
[CBSE PMT 1997]		[CBSE PMT 1990; MP PET 1992;
(a) $\left[ Cu\{O = C(NH_2)_2\}Cl_2 \right]$		MP PMT 1995, 97; Kurukshetra CET 2002]
(b) $\left[ CuCl_2 \{ O = C(NH_2)_2 \}_2 \right]$		(a) Potassium hexacyanoferrate (II)
(c) $ [Cu\{O = C(NH_2)_2\}Cl]Cl $		(b) Potassium ferrocyanide (c) Tetrapotassium hexacyanoferrate (II)
(d) $[CuCl_2]{O = C(NH_2)_2H_2}$		(d) Tetrapotassium ferroushexacyanide (II)
The IUPAC name of the complex $[Pt(NH_3), Cl_2]$ is	15.	The IUPAC name of $\left[Ni(CO)_4\right]$ is $\left[ \text{RPET 1999} \right]$
(a) Platinum (II) diammino dichloride		(a) Tetra carbonyl nickel (II)
(b) Diammino dichloro platinate (IV)		(b) Tetra carbonyl nickel (0)
(c) Bis (ammino) dichloro platinum (IV)		(c) Tetra carbonyl nickelate (II)
(d) Dichloro diammine platinum (ll)	- 6	(d) Tetra carbonyl nickelate (0)
Correct formula of diammine silver (1) chloride is  (a) $Ag(NH_2)Cl$ (b) $Ag(NH_2)Cl$	16.	The correct nomenclature for $Fe_4[Fe(CN)_6]_3$ is
		[MP PMT 1994] (a) Ferroso-ferric cyanide
(c) $[Ag(NH_3)_2]Cl$ (d) $[Ag(NH_2)_2]Cl$		(b) Ferric-ferrous hexacyanate
The formula of sodium nitroprusside is [AIIMS 1992] (a) $Na_4[Fe(CN)_5NOS]$ (b) $Na_2[Fe(CN)_5NO]$		(c) Iron (III) hexacyanoferrate (II)
1		(d) Hexacynoferrate (III-II)
(c) $NaFe[Fe(CN)_6]$ (d) $Na_2[Fe(CN)_6NO_2]$	17.	The IUPAC name of compound $Na_3[Co(ONO)_6]$ will be
The correct name of $[Pt(NH_3)_4 Cl_2][PtCl_4]$ is		[MP PMT 2000]
[MP PET 2003]		(a) Hexanitritocobalt (III) sodium (b) Sodium cobalt nitrite
(a) Tetraammine dichloro platinum (iv) tetrachloro platinate (ii)		(c) Sodium hexanitrocobaltate (III)
(b) Dichloro tetra ammine platinium (iv) tetrachloro platinate (ii) (c) Tetrachloro platinum (ii) tetraammine platinate (iv)		(d) Sodium hexanitritocobaltate (III)
(d) Tetrachloro platinum (ii) dichloro tetraammine platinate (iv)	18.	In which of the following complexes oxidation state of metal is zero
Correct formula of potassium ferrocyanide is		(a) $\left[ Pt(NH_3)_2 Cl_2 \right]$ (b) $\left[ Cr(CO)_6 \right]$
[CBSE PMT 1988]		(c) $\left[ Cr(NH_3)_3 Cl_3 \right]$ (d) $\left[ Cr(en)_2 Cl_2 \right]$
(a) $K_4[Fe(CN)_6]$ (b) $K_2[Fe(CN)_6]H_2O$	19.	The oxidation number of $Cr$ in $[Cr(NH_3)_6]Cl_3$ is
(c) $K_3[Fe(CN)_6]$ (d) None of these		[CBSE PMT 2001]
The IUPAC name of $\left[Co(NH_3)_3(NO_2)_3\right]$ is		(a) 8 (b) 6
(a) Trinitrotriammine cobalt (III)		(c) 4 (d) 3
(b) Trinitrotriammine cobalt (II)	20.	In $\left[Ni(NH_3)_4\right]SO_4$ , the E.A.N. of $Ni$ is
(c) Trinitrotriammine cobalt (III) ion		(a) 34 (b) 35
(d) Trinitritriammine cobaltate (III)	01	(c) 36 (d) 37 IUPAC name of $[Co(ONO)(NH_3)_5 Cl_2]$ is
In $K_4[Fe(CN)_6]$ , the E.A.N. of $Fe$ is [DCE 2000]	21.	[AMU 2002]
(a) 33 (b) 35		(a) Pentaammine nitro cobalt (III) chloride
(c) 36 (d) 26		(b) Pentaammine nitrito cobalt (III) chloride
Which of the following pairs is not correctly matched [MP PET 1993]		(c) Pentaammine nitroso cobalt (III) chloride
•		(d) Pentaammine oxo-nitro cobalt (III) chloride
(a) Effective atomic number of $Pt$ in $[PtCl_6]^{2^-} = 84$	22.	The oxidation number of $Pt$ in $[Pt(C_2H_4)Cl_3]^-$ is
(b) Absorption peak for $\left[Cr^{III}(NH_3)_6\right]^{+3} = 21680cm^{-1}$		[MNR 1993]
(c) Crystal field stabilization energy of $d^2$ in weak ligand field		(a) +1 (b) +2 (c) +3 (d) +4
$= (-)0.8\Delta_0$	23.	The oxidation state of cobalt in the complex compound
•	-	$[Co(NH_3)_6]Cl_3$ is
		(a) + 3 (b) + 6

(b) Nitrochlorocyanotriammine cobalt (III)

 $(c) \quad Cyanonitrochlorotriammine\ cobalt\ (III)$ 

 $(d) \quad Triamminenitrochlorocyano \ cobalt \ (III)$ 

The oxidation number of Pt in  $[Pt(C_2H_4)Cl_3]^-$  is



	(c) + 5 (d) + 2		(a) Potassium tri oxalato iridium (III)
24.	The correct IUPAC name of potassium cuprochloride is		(b) Potassium tri oxalato iridate (III)
	(a) Potassium copper (1) tetrachloride		(c) Potassium tris (oxalato) iridium (III)
	(b) Potassium tetrachlorocuprate (1) (c) Tetrachloropotassium cuprate (1)		(d) Potassium tris (oxalato) iridate (III)
	(d) Tetrachlorocopper (l) potassiate	37.	The charge on $[Ag(CN)_2]^-$ complex is [AIIMS 2001]
25.	The effective atomic number of cobalt in the complex		(a) $-1$ (b) $+1$
	$[Co(NH_3)_6]^{3+}$ is [MP PET 2003]		(c) $+2$ (d) $+3$
	(a) 36 (b) 33	38.	The IUPAC name of $[Co(NH_3)_6]Cl_3$ is [IIT-JEE 1994]
	(c) 24 (d) 30		(a) Hexammine cobalt (III) chloride
26.	IUPAC name of $K_3 Fe(CN)_6$ is		(b) Hexammine cobalt (II) chloride
	[MP PMT 1993; MP PET 1997]		(c) Triammine cobalt (III) trichloride
	(a) Potassium ferrocyanide (II)		(d) None of these
	(b) Potassium hexaferrocyanate (III)	39.	IUPAC name of $\left[Co(NH_3)_3(H_2O)_2Cl\right]$ $Cl_2$ is
	(c) Potassium ferrohexacyanate (II)		[MP PET 1994]
	(d) Potassium hexacyanoferrate (III) The EAN of iron in potassium ferricyanide is		(a) Diaquachlorodiammine cobalt (III) chloride
27.	[Pb. CET 2000]		(b) Triamminediaquachloro cobalt (III) chloride
	(a) 18 (b) 54		(c) Chlorodiamminediaqua cobalt (III) chloride
	(c) 35 (d) 23		(d) Diamminediaquachloro cobalt (II) chloride
28.	In the coordination compound, $K_4[Ni(CN)_4]$ oxidation state of	40.	Dichloro diammine platinum (II) complex has the formula
	nickel is [AIEEE 2003]		[MP PMT 1997]
	(a) $-1$ (b) 0		(a) $Pt[Cl_2(NH_3)_2]$ (b) $Pt[R.(NH_2)_2]Cl_2$
	(c) $+1$ (d) $+2$		(c) $[PtCl_2(NH_3)_2]$ (d) $[Pt.R.(NH_2)_2]Cl_2$
29.	According to IUPAC nomenclature sodium nitroprussied is named is	41.	The formula of potassiumdicyano <i>bis</i> (oxalato) nickelate (II) is
	[CBSE PMT 2003] (a) Sodium pentacyanonitrosyl ferrate (III)	7"	(a) $K_4\left[Ni(CN)_2(Ox)_2\right]$ (b) $K_3\left[Ni_2(Ni_2(CN)_2(Ox)_2\right]$
	(a) Sodium pentacyanonitrosyl ferrate (III) (b) Sodium nitroferricyanide		
	(c) Sodium nitroferrocyanide		(c) $K[Ni(CN)(Ox)_2]$ (d) $K_2[Ni(CN)_2(Ox)_2]$
	(d) Sodium pentacyanonitrosyl ferrate (ll)	42.	The value of $x$ which appears in the complex $[Ni(CN)_4]^x$ is
30.	Pick out the complex compound in which the central metal atom	72.	
	obeys EAN rule strictly [KCET 2003]		(a) + 2 (b) - 2 (c) 0 (d) 4
	(a) $K_4[Fe(CN)_6]$ (b) $K_3[Fe(CN)_6]$		
	(c) $[Cr(H_2O)_6]Cl_3$ (d) $[Cu(NH_3)_4]SO_4$	43.	Pick the correct name of $[Co(NH_3)_5 Cl]Cl_2$ [AMU 2001]
31.	Which of the following is wrong statement [BHU 2003]		(a) Chloropentammine cobalt (III)
	(a) $Ni(CO)_4$ has oxidation number + 4 for $Ni$		(b) Pentammine cobalt (III) chloride
	(b) $Ni(CO)_4$ has zero oxidation number for $Ni$		(c) Chloropentammine cobalt (III) chloride
	(c) Ni is metal		(d) Chloropentammine cobalt (II) chloride
	(d) CO is gas	44.	The valency of cuprammonium ion is
32.	Oxidation state of $Fe$ in $K_3[Fe(CN)_6]$ [RPMT 2002]		(a) + 4 (b) + 2 (c) - 2 (d) - 4
	(a) 2 (b) 3	45.	In which of the following compounds transition metal has zero
	(c) 0 (d) None of these	40.	oxidation state [CBSE PMT 1999]
33.	Which complexes have zero oxidation state		(a) $CrO_5$ (b) $NH_2.NH_2$
	(a) Carbonyl (b) Ferrocyanide		(c) $NOClO_4$ (d) $[Fe(CO)_5]$
0.4	(c) Amine (d) Cyanide		
34.	The proper name for $K_2[PtCl]_6$ is [MH CET 2002]	46.	The complex chlorocompound diaquatriammine cobalt (III) chloride is represented as [CBSE PMT 2002]
	(a) Potassium platinum hexachloride		(a) $[Co(NH_3)_3(H_2O)_3]Cl_2$
	(b) Potassium hexachloro platinum IV (c) Potassium hexachloro platinate IV		( ) - ( ) 3.3 ( 2 ) 3- 2
	(d) Potassium hexachloro platinum		(b) $[Co(NH_2)_3 (H_2O)_2]Cl_2$
35.	IUPAC name of $K_3[Al(C_2O_4)_3]$ is called		(c) $[CoCl(NH_3)_3 (H_2O)_2]Cl_3$
55.	[MP PMT 1993, 02, 03]		(d) $[CoCl(NH_3)_3 (H_2O)_2]Cl_2$
	(a) Potassium alumino oxalato	47.	The complex compound $[Co(NH_3)_3 NO_2 ClCN]$ is named as
	(b) Potassium aluminium (III) trioxalate	.,.	(a) Chlorocyanonitrotriammine cobalt (III)
			(a) chiorocyanoma ou laminine cobait (iii)

[MP PMT 2001]

 $(c) \quad Potassium \ trioxalato \ aluminate \ (III)$ 

 $(d) \quad Potassium \ trioxalato \ aluminate \ (1V)$ 

36.

The I.U.P.A.C. name of  $K_3[\operatorname{Ir}(C_2O_4)_3]$  is



# UNIVERSAL SELF SCORER 912 Co-ordination Chemistry

(b) Potassium hexacyanoferrate (III) (c) Potassium hexacyanoiron (II) (d) Tripotassium hexacyanoiron (II) Which compound is zero valent metal complex [KCET 2005] (a) $[Cu(NH_3)_4]SO_4$ (b) $[Pt(NH_3)_2Cl_2]$ (c) $[Ni(CO)_4]$ (d) $K_3[Fe(CN)_6]$	12.	Which of the following isomeric pairs shows ionization isomerism  (a) $[Co(NH_3)_6][Cr(CN)_6]$ and $[Cr(NH_3)_6][Co(CN)_6]$ (b) $[Cr(H_2O)_6]Cl_3$ and $[Cr(H_2O)_5Cl]Cl_2.H_2O$ (c) $[Pt(NH_3)_2Cl_2]$ and $[Pt(NH_3)_4][PtCl_4]$ (d) $[Co(NH_3)_5Br]SO_4$ and $[Co(NH_3)_5SO_4]Br$ Among the following ions which one has the highest paramagnetism[IIT 1993; I (a) $[Cr(H_2O)_6]^{3+}$ (b) $[Fe(H_2O)_6]^{2+}$
(c) Potassium hexacyanoiron (II) (d) Tripotassium hexacyanoiron (II) Which compound is zero valent metal complex  (a) $[Cu(NH_3)_4]SO_4$ (b) $[Pt(NH_3)_2Cl_2]$		(a) $[Co(NH_3)_6][Cr(CN)_6]$ and $[Cr(NH_3)_6][Co(CN)_6]$ (b) $[Cr(H_2O)_6]Cl_3$ and $[Cr(H_2O)_5Cl]Cl_2.H_2O$ (c) $[Pt(NH_3)_2Cl_2]$ and $[Pt(NH_3)_4][PtCl_4]$ (d) $[Co(NH_3)_5Br]SO_4$ and $[Co(NH_3)_5SO_4]Br$
(c) Potassium hexacyanoiron (II) (d) Tripotassium hexacyanoiron (II) Which compound is zero valent metal complex  (a) $[Cu(NH_3)_4]SO_4$ (b) $[Pt(NH_3)_2Cl_2]$	12.	(a) $[Co(NH_3)_6][Cr(CN)_6]$ and $[Cr(NH_3)_6][Co(CN)_6]$ (b) $[Cr(H_2O)_6]Cl_3$ and $[Cr(H_2O)_5Cl]Cl_2.H_2O$ (c) $[Pt(NH_3)_2Cl_2]$ and $[Pt(NH_3)_4][PtCl_4]$
(c) Potassium hexacyanoiron (II) (d) Tripotassium hexacyanoiron (II) Which compound is zero valent metal complex [KCET 2005]	12.	(a) $[Co(NH_3)_6][Cr(CN)_6]$ and $[Cr(NH_3)_6][Co(CN)_6]$ (b) $[Cr(H_2O)_6]Cl_3$ and $[Cr(H_2O)_5Cl]Cl_2.H_2O$
(c) Potassium hexacyanoiron (II) (d) Tripotassium hexacyanoiron (II)	12.	(a) $\left[Co\left(NH_3\right)_6\right]\left[Cr\left(CN\right)_6\right]$ and $\left[Cr(NH_3)_6\right]\left[Co(CN)_6\right]$
(c) Potassium hexacyanoiron (II)	12.	
	12.	Which of the following isomeric bairs shows ionization isomerism
		(d) Unpaired
(a) Potassium hexacyanoferrate (II)	-	(c) Paired (d) Uppgired
* * *	2005]	(b) Both paired and unpaired
(d) Potassium ammine cyano peroxo dioxo chromatic (IV)		(a) No
• • • • • • • • • • • • • • • • • • • •	11.	Paramagnetic co-ordination compounds contain electrons
(a) Potassium ammine dicyano dioxoperoxochromate (VI)		(c) $[Co(NH_3)_5 NO_2]Cl_2$ (d) $[Pt(NH_3)_2 Cl_2]$
[DCE 2003]		(a) $[Cr(NH_3)_4 Cl_2]Cl$ (b) $[Co(en)_2 Cl_2]Cl$
The IUPAC name of $K_2[Cr(CN)_2O_2(O)_2(N\!H_3)]$ is	10.	Which one of the following will not show geometrical isomerism
(d) Hexamminechloride		(d) Low oxidation and higher oxidation states
(c) Hexamminechromium (II) ion		(b) Complex cation and complex anion (c) Inner sphere and outer sphere
(a) Hexamminechromium (VI) ion (b) Hexamminechromium (III) ion		(a) Cis and Trans structure
	-	between the [UPSEAT 2002]
( ) [ ( ) ( ) ( ) ( ) ( ) ( ) ( ) ( ) (	9.	(c) Three unpaired $e^-$ (d) No unpaired $e^-$ Coordination isomerism is caused by the interchange of ligands
(d) $[Co(CN)_k]^{3-}, MnO_3$		
(c) $TiO_3$ , $MnO_2$		(a) One unpaired $e^-$ (b) Two unpaired $e^-$
(b) $CrO_2Cl_2, MnO_4^-$	8.	$[Ti(H_2O)_6]^{+3}$ is paramagnetic in nature due to [RPMT 2002]
(a) $[Fe(CN)_6]^{3-}$ , $[Co(CN)_6]^{3-}$		(a) $[Cr(NI_3)_6]Cl_3$ (b) $[Co(NI_3)_5]DrDO_4$ (c) $[Cr(en)_2Cl_2]$ (d) $[Cr(en)_3Cl_3]$
highest possible oxidation state is [IIT-JEE (Screening) 2004]	7.	Which would exhibit ionisation isomerism [MP PET 1997] (a) $\left[Cr(NH_3)_6\right]Cl_3$ (b) $\left[Co(NH_3)_5Br\right]SO_4$
The pair of the compounds in which both the metals are in the	_	(c) Linkage (d) Ionization
(d) None of these		(a) Geometrical (b) Optical
(c) Pentamminentrocobalt (II) chloride		isomers [MP PMT 1993; MP PET 1995, 2001]
	6.	$[Co(NH_3)_4 Cl_2]NO_2$ and $[Co(NH_3)_4 Cl.NO_2]Cl$ are
		(a) Linkage (b) Geometrical (c) Ionization (d) Optical
		[MP PMT 1993, 94, 95; MP PET 1997; MP PET/PMT 1998]
		which type of isomerism
• •	5.	$\left[ {\it Co(NH_{3})_{5}\it Br} \right] {\it SO}_{4} \; {\it and} \; \left[ {\it Co(NH_{3})_{5}\it SO}_{4} \right] \! \it Br \; {\it are examples of}$
**		(c) Linkage isomers (d) Coordination isomers
		(a) Geometrical isomers (b) Optical isomers
(a) + 2 (b) + 3	4.	$[Co(NH_3)_5 NO_2]Cl_2$ and $[Co(NH_3)_5 (ONO)]Cl_2$ are related to each other as
The oxidation state of $Co$ in $\left[Co(H_2O)_5Cl\right]^{2+}$ is		
(d) Potassium dicyanosilver (II)		(c) $[Cr(NH_3)_6]Cl_3$ (d) $[Cr(en)_2Cl_2]^+$
(c) Potassium dicyanoargentate (l)	-	(a) $\left[Cr(NH_3)_6\right]\left[Co(CN)_6\right]$ (b) $\left[Co(en)_2Cl_2\right]$
(b) Potassium silver cyanide	3.	(d) 4 Which would exhibit co-ordination isomerism
- ' '		(a) Zero (b) 2 (c) 3 (d) 4
		(Atomic no. of $Co = 27$ ) [CBSE PMT 2003]
	2.	The number of unpaired electrons in the complex ion $\left[CoF_6\right]^{3-}$ is
•		
(-)		·
· ·		geometric isomerism ( $A$ and $B$ are monodentate ligands)  (a) $[MA_5B]$ (b) $[MA_2B_4]$
•	1.	Which one of the following octahedral complexes will not show
	(a) $+1$ (b) $+2$ (c) $+3$ (d) $+4$ What is the structural formula of lithium tetrahydridoaluminate (a) $A[LiH_4]$ (b) $Al_2[LiH_4]_3$ (c) $Li[AlH_4]$ (d) $Li[AlH_4]_2$ IUPAC name for $K[Ag(CN)_2]$ is (a) Potassium argentocyanide (b) Potassium dicyanoargentate (l) (d) Potassium dicyanoargentate (l) (d) Potassium dicyanosilver (II) The oxidation state of $Co$ in $[Co(H_2O)_5Cl]^{2+}$ is (a) $+2$ (b) $+3$ (c) $+1$ (d) $+4$ The chemical formula of diammine silver ( $IIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIII$	(a) $+1$ (b) $+2$ (c) $+3$ (d) $+4$ What is the structural formula of lithium tetrahydridoaluminate (a) $Al[LiH_4]$ (b) $Al_2[LiH_4]_3$ 2.  (c) $L[AIH_4]$ (b) $Al_2[LiH_4]_3$ 2.  (d) $L[AIH_4]$ (e) $L[AIH_4]_2$ 11UPAC name for $K[Ag(CN)_2]$ is (a) Potassium argentocyanide (b) Potassium dicyanoargentate (1) (d) Potassium dicyanoargentate (1) (d) Potassium dicyanosilver (11) (d) Potassium dicyanosilver (11) (e) Potassium dicyanosilver (11) (f) Potassium dicyanosilver (11) (h) $+1$ (h)



(c)  $[Cu(H_2O)_{\epsilon}]^{2+}$ (d)  $\left[Zn(H_2O)_{\epsilon}\right]^{2+}$ (b) Only  $[Fe(CN)_6]^{3-}$  is paramagnetic Amongst  $Ni(CO)_4$ ,  $\left[Ni(CN)_4\right]^{2-}$  and  $\left[NiCl_4\right]^{2-}$ (c) Only  $[FeF_6]^{3-}$  is paramagnetic [IIT 1991] 14. (d) Both are diamagnetic (a)  $Ni(CO)_4$  and  $[NiCl_4]^{2-}$  are diamagnetic and Which of the following is paramagnetic [AFMC 1997] 26.  $[Ni(CN)_{A}]^{2-}$  is paramagnetic (a)  $\left[Ni(CO)_{4}\right]$ (b)  $[Co(NH_3)_6]^{3+}$ (b)  $[NiCl_4]^{2-}$  and  $[Ni(CN)_4]^{2-}$  are diamagnetic and (c)  $[Ni(CN)_{4}]^{2-}$ (d)  $\left[NiCl_{A}\right]^{2-}$  $Ni(CO)_4$  is paramagnetic The total number of possible isomers for the complex compound 27.  $\left[Cu^{II}(NH_3)_4\right]\left[Pt^{II}Cl_4\right]$  are (c)  $Ni(CO)_A$  and  $[Ni(CN)_A]^{2-}$  are diamagnetic and [CBSE PMT 1998; DPMT 2004; ] & K CET 2005]  $[NiCl_{4}]^{2-}$  is paramagnetic (b) 4 (d) 6 (c) 5 (d)  $Ni(CO)_4$  is diamagnetic and  $[NiCl_4]^{2-}$  and  $[Ni(CN)_4]^{2-}$ 28. Which one of the following shows maximum paramagnetic character are paramagnetic (a)  $\left[ Cr(H_2O)_6 \right]^{3+}$ (b)  $[Fe(CN)_6]^{4-}$  $[Co(NH_3), Cl_2]^+$  exhibits 15. (c)  $[Fe(CN)_{6}]^{3-}$ (d)  $[Cu(H_2O)_6]^{2+}$ (a) Geometrical isomerism (b) Optical isomerism (c) Bonding isomerism (d) Ionisation isomerism  $[Co(NH_3)_6][Cr(C_2O_4)_3]$ complexes 29. The compound which does not show paramagnetism is 16.  $[Cr(NH_3)_6][Co(C_2O_4)_3]$ [AMU 2002] [IIT 1992] (a) Linkage isomerism (b) Geometrical isomerism (b)  $\left[Ag(NH_3), Cl\right]$ (a)  $\left[ Cu(NH_3)_4 \right] Cl_3$ (c) Coordination isomerism (d) Ionisation isomerism (c) *NO* Which of the following exhibits highest molar conductivity 30. [MP PET 1994] The number of geometrical isomers for  $[Pt(NH_3), Cl_2]$  is 17. (a)  $\left[Co(NH_3)_6\right]Cl_3$ (b)  $\left[Co(NH_3)_5 Cl\right]Cl_2$ [CBSE PMT 1995] (c)  $\left[Co(NH_3)_{4}Cl_{2}\right]Cl$ (d)  $\left[Co(NH_3)_2 Cl_3\right]$ (a) Two (b) One (c) Three (d) Four Which of the following compounds is colourless 31.  $\left[ Cr(H_2O)_6 Cl_3 \right]$  and The pair of complex compounds [MP PET 1994] 18. (a)  $Cu_2(CH_3COO)_4.2H_2O$  (b)  $Cu_2Cl_2$  $[Cr(H_2O)_5 Cl]Cl_2H_2O$  are an example of (c)  $CuSO_{4}.5H_{2}O$ (a) Linkage isomerism (b) Ionisation isomerism (d)  $[Cu(NH_3)_4]SO_4.4H_2O$ (c) Coordination isomerism (d) Hydrate isomerism The type of magnetism exhibited by  $[Mn(H_2O)_6]^{2+}$  ion is 32. The number of geometrical isomers of the complex 19.  $\left[Co(NO_2)_2(NH_3)_2\right]$  is [IIT 1994] [CBSE PMT 1997] (a) Paramagnetism (b) Diamagnetism (b) 3 (c) Both (a) and (b) (d) None of these (c) 4 (d) o The number of isomers possible for square planar complex The type of isomerism present in nitropentamine chromium (III) 20.  $K_2[PdClBr_2(SCN)]$  is [MP PET 1994] chloride is [AIEEE 2002] (a) Optical (b) Linkage (b) 3 (d) Polymerisation (c) lonization (c) 4 [MP PMT 2001] The number of unpaired electrons in  $Ni(CO)_4$  is Which of the following compounds exhibits linkage isomerism 21. 34. (a)  $[Co(en)_3]Cl_3$ (b)  $[Co(NH_3)_6[Cr(CN)_6]$ [AIIMS 1997] (c)  $[Co(en)_2 NO_2 Cl]Br$ (d)  $[Co(NH_3)_5 Cl]Br_2$ (a) Zero (b) One Pick out from the following complex compounds, a poor electrolytic (c) Three (d) Five 22. conductor in solution [MP PMT 1994] If the magnetic moment of  $[Ag(CN)_2]^{-1}$  is zero, then the number 35. (a)  $K_2[PtCl_6]$ (b)  $[Co(NH_3)_3(NO_2)_3]$ of unpaired electrons will be [MP PET 1995] (c)  $K_4[Fe(CN)_6]$ (d)  $\left[Cu(NH_3)_A\right]SO_A$ (b) 2 [MP PET 2003] The possible number of optical isomers in  $\left[Co(en)_2Cl_2\right]^+$  are 23.  $[Pt(NH_3)_4 Cl_2]Br_2$  and  $[Pt(NH_3)_4 Br_2]Cl_2$  are related to each (a) 2 (b) 3 (c) 4 (d) 6 [MP PET 1996: AFMC 2000: CBSE PMT 2001] Magnetic moment of  $[Cu(NH_3)_4]^{2+}$  ion is [RPET 2003] 24. (a) Optical isomers (b) Coordinate isomers (c) lonization isomers (d) Linkage isomers (c) 2.23

[RPET 1999]

What is true for  $[Fe(CN)_6]^{3-}$  and  $[FeF_6]^{3-}$ 

(a) Both are paramagnetic

25.

Which of the following complex will show geometrical as well as 37. optical isomerism (*en* = ethylene diamine)

[KCET 1996]

	ERSAL	914 Co-ordination	Chei	mistry
	(a)	$Pt(NH_3)_2Cl_2$	(b)	$[Pt(NH_3)Cl_4]$
	(c)	$[Pt(en)_3]^{4+}$	(d)	$[Pt(en)_2]Cl_2$
38.	Whi	ch of the following complexe	s is d	iamagnetic
				[RPMT 1997]
	(a)	$Ni(CO)_4$	(b)	$Ni(Cl_4)^{2-}$
	(c)	$Ni(Br)_4^{2-}$	(d)	$NiCl_2.4H_2O$
39.	Whi	ch one has the highest paran	nagne	tism [AMU 2001]
	(a)	$Ni(CO)_4$	(b)	$[Ni(NH_3)_4]Cl_2$
	(c)	$[Ni(NH_3)_6]Cl_2$	(d)	$[Cu(NH_3)_4]Cl_2$
40.		ch of the following coord cal isomerism	liantio	on compounds would exhibit [CBSE PMT 2004]
	(a)	trans-dicyanobis (ethylened	iamine	•
	(b)	tris-(ethylenediamine) cobal		
	(c)	•	,	de
41.	(d)	diamminedichloroplatinum ch of the following does not	. ,	ontical icomer
41.	VVIII	ch of the following does not	liave	[AllMS 2004]
	(a)	$[Co(NH_3)_3Cl_3]$		
	(b)	$[Co(en)_3]Cl_3$		
	(c)	$[Co(en)_2Cl_2]Cl$	(d)	$[Co(en)(NH_3)_2Cl_2]Cl$
42.	Cha ison		linatio	on sphere yields which types of [DCE 2002]
	(a)	Optical	` '	Geometrical
40	(c)	lonisation	(d)	
43.		es of isomerism shown by $C(NH_3)_5 NO_2 Cl_2$ is		[DCE 2002]
	(a)	Optical	(b)	lonisation
	(c)	Geometrical	(d)	Linkage
44.		ch of the following will not g		_
				$[Co(NH_3)_4 Cl_2]Cl$
		$[Co(NH_3)_5 Cl]Cl_2$		
45.		many ions are produced fro		
	(a) (c)	6 3	(b)	4 2
46.		colour of $CoCl_3.5NH_3.H$	. ,	
•	(a)	Orange yellow	(b)	
	(c)	Green	(d)	Violet
47	(e)	Pink		nested to be a narramannetic

complex

48.

49.

50.

(a)  $[Ni(H_2O)_6]^{2+}$ 

(c)  $\left[ Zn(NH_3)_4 \right]^{2+}$ 

(c)  $[Pt(en)Cl_2]$ 

 $BaCl_2$  (aq.)

(a) 1

(c) 3

 $AgNO_3$  in aqueous mediun

chloropentammine cobalt (III) chloride

(c)  $[Cr(NH_3)_5 Cl]SO_4$ (d) Both (b) and (c) The number of precipitable halide ions in the sample 51.  $[Pt(NH_3)Cl_2Br]Cl$  will be (a) 2 (b) 3 (c) 4 (d) 1 52. The colour of tetrammine copper (11) sulphate is (a) Blue (b) Red (c) Violet (d) Green The number of ions per mole of a complex  $[CoCl_2.5 NH_3]Cl_2$  in 53. aqueous solution will be [MP PMT 2001] (a) Nine (b) Four (c) Three (d) Two How many unpaired electrons are present in the central metal ion of 54.  $[CoCl_4]^{2-}$ [Orissa JEE 2005] (b) 4 (a) 3 (d) 2 (c) 5 What is the magnetic moment of  $K_3[FeF_6]$ 55. [Orissa JEE 2005] (b) 4.89 BM (a) 5.91 BM (c) 3.87 BM (d) 6.92 BM (i)  $K_4[Fe(CN)_6]$  (ii)  $K_3[Cr(CN)_6]$  (iii)  $K_3[Fe(CN)_6]$ 56. (iv)  $K_2[Ni(CN)_4]$ Choose the complex which is paramagnetic [Kerala CET 2005] (b) (i), (iii) and (iv) (i), (ii) and (iii) (i), (iii) and (iv) (d) (i), (ii) and (iv) (c) (e) (ii) and (iv) In coordination compound  $[Co(en)_2 Cl_2]Cl$  which is false 57. [Kerala CET 2005] Show geometrical Isomerism (b) Shanp openicaddsomerism Show ionic Isomerism (c) (d) A octahedral complex A cationic complex [RPET 1999] **Hybridisation and Geometry** The correct structural formula of zeise's salt is (a)  $K^{+} \left[ PtCl_{3} - \eta^{2} - (C_{2}H_{4}) \right]$ (b)  $K_2 \left[ PtCl_3 - \eta^2 - C_2 H_4 \right]$ (c)  $K^{+} \left[ PtCl_{2} - \eta^{2} - (C_{2}H_{4}) \right] Cl^{-}$ 

(d)  $K^+[PtCl_3(C_2H_4]^-$ 

[MP PMT 1991, 2000]

[MP PMT 1994]

(b)  $\left[Ni(CO)_4\right]$ 

Which one of the following will give a white precipitate with

How many ions will be produced in solution from one molecule of

Which of the following complex will give white precipitate with

(a)  $[Co(NH_3)_4 SO_4]NO_2$  (b)  $[Cr(NH_3)_5 SO_4]Cl$ 

(d) 4

(a)  $\left[Co(NH_3)_5Cl\right](NO_2)_5$  (b)  $\left[Pt(NH_3)_5Cl_2\right]$ 

(d)  $\left[ Co(NH_3)_6 \right]^{+3}$ 

(d)  $\left[Pt(NH_3)_{4}\right]Cl_2$ 

2. The correct order of hybridisations of central atom in

 $N\!H_3, [PtCl_4\,]^{2-}, PCl_5 \ \ {\rm and} \ \ BCl_3 \ \ {\rm is}$  [MP PMT 2003]

(a)  $dsp^2, dsp^3, sp^2$  and  $sp^3$ 

(b)  $sp^3, sp^3, sp^3d$  and  $sp^2$ 

(c)  $dsp^2$ ,  $sp^2$ ,  $sp^3$  and  $dsp^3$ 

(d)  $dsp^2$ ,  $sp^3$ ,  $sp^2$  and  $dsp^3$ 

3. One mole of the complex compound  $Co(NH_3)_5 Cl_3$ , gives 3 moles of ions on dissolution in water. One mole of the same complex reacts with two moles of  $AgNO_3$  solution to yield two moles of AgCl(s). The structure of the complex is

[AIEEE 2003]



(a)  $[Co(NH_3)_5 Cl]Cl_2$ The shape of  $\left[Cu(NH_3)_4\right]^{2+}$  is square planar,  $Cu^{2+}$  in this 16. (b)  $[Co(NH_3)_3 Cl_3].2NH_3$ [NCERT 1989; RPET 1999] (a)  $sp^3$  hybridised (b)  $dsp^2$  hybridised (c)  $[Co(NH_3)_4 Cl_2]Cl.NH_3$ (d)  $[Co(NH_3)_4 Cl]Cl_2.NH_3$ (d)  $sp^3d^2$  hybridised (c)  $sp^3d$  hybridised The geometry of  $Ni(CO)_4$  and  $Ni(PPh_3)_2 Cl_2$  are Cuprammonium ion  $\left[Cu(NH_3)_A\right]^{2+}$  is 17. [IIT-JEE 1999; DCE 2002] [MP PMT 1997; KCET 2002] (a) Both square planar (a) Tetrahedral (b) Square planar Tetrahedral and square planar respectively (c) Triangular bipyramid (d) Octahedral (c) Both tetrahedral In the complex  $[SbF_5]^{2-}$ ,  $sp^3d$  hydridisation is present. Geometry Square planar and tetrahedral respectively of the complex is [Pb. PMT 2000] 18. Which complex has square planar structure [IIPMER 2002] (b) Square bipyramidal (a) Square pyramidal  $Ni(CO)_4$ (b)  $[NiCl_{4}]^{2-}$ (c) Tetrahedral (d) Square (c)  $[Ni(H_2O)_6]^{2+}$ (d)  $[Cu(NH_3)_4]^{2+}$ 6. The type of hybridization involved in the metal ion of  $[Ni(H_2O)_6]^{2+}$  complex is  $[Pt(NH_3)_A]Cl_2$  is 19. [DCE 2001] (a) Square planar (a)  $d^3sp^2$ (b)  $sp^{3}d^{2}$ (b) Tetrahedral (c) Pyramidal (d) Pentagonal (d)  $dsp^2$ A complex involving  $dsp^2$  hybridization has 20. In the formation of  $K_4 Fe(CN)_6$ , the hybridisation involved is 7. (a) A square planar geometry (b) A tetrahedral geometry (a)  $sp^2$ (b)  $d^2 s p^3$ (c) An octahedral geometry (c)  $d^3sp^2$ (d)  $d^4 p$ (d) Trigonal planar geometry Which of the following is not true for metal carbonyls 8. A tetrahedral complex ion is formed due to ...... hybridization 21. [MP PET 1993] (a)  $sp^2$ (b)  $sp^3$ (a) The oxidation state of the metal in the carbonyls is zero (c)  $dsp^2$ (d)  $d^2sp^3$ (b) The secondary carbonyls are obtained from photodecomposition Back bonding is involved in which of the organometallic compounds 22. (c) Metal carbonyls are single bonded species (b)  $Mg^{2+}(C_5H_5^-)_2$ (a)  $[(CH_3)_3Al]_2$ (d)  $d\pi - p\pi$  overlap is observed in metal carbonyls (d)  $[(C_5H_5)_2Fe]$ (c) R - Mg - XAn octahedral complex is formed, when hybrid orbitals of the 9. [DCE 2003] following type are involved  $d^2sp^3$  hybridisation leads to 23. (a)  $sp^3$ (b)  $dsp^2$ (a) Hexagonal shape (b) Trigonal bipyrimidal (d) Tetrahedral shape (c) Octahedral shape (c)  $sp^3d^2$ (d)  $sp^2d$ Chromium hexacarbonyl is an octahedral compound involving 24. Which one is an example of octahedral complex 10. (a)  $sp^3d^2$ (b)  $dsp^2$ [MP PET 2000] (a)  $FeF_6^{3-}$ (b)  $Zn(NH_3)_4^{2+}$ (c)  $d^2sp^3$ (d)  $d^3sp^2$  orbitals (c)  $Ni(CN)^{2-}_{\Lambda}$ (d)  $Cu(NH_3)^{2+}_{4}$  $[CoF_6]^{-3}$  is formed by ..... hybridization 25. Which of the following complexes has a square planar geometry 11. (b)  $d^3 s p^2$ (a)  $d^2sp^3$ (a)  $Ag(NH_3)_2^+$ (b)  $Cu(en)_{2}^{2+}$ (c)  $d^2sp^3$ (d)  $sp^3d^2$ (c)  $[MnCl_A]^{2-}$ (d)  $Ni(CO)_{A}$ 26. The species havoing tetrahedral shape is [IIT-JEE (Screening) 2004] The shape of  $\left[Fe(CN)_6\right]^{4-}$  ion is 12. (a)  $[PdCl_{4}]^{2-}$ (b)  $[Ni(CN)_4]^{2-}$ (a) Hexagonal (b) Pyrimidal (c)  $[Pd(CN)_{\Lambda}]^{2-}$ (d)  $[NiCl_{4}]^{2-}$ (c) Octahedral (d) Octagonal What is the shape of  $Fe(CO)_5$  [CBSE PMT 2000] 13.  $[Ni(CO)_4], [Ni(CN)_4]^{2-}, [NiCl_4]^{2-}$ 27. species. the (a) Linear (b) Tetrahedral hybridization states at the Ni atom are, respectively (c) Square planar (d) Trigonal bipyramidal [CBSE PMT 2004; MP PMT 1992; BHU 1995; AFMC 1997] What type of hybridization is involved in  $[Fe(CN)_6]^{3-}$ 14. (a)  $sp^3, sp^3, dsp^2$ (d)  $dsp^2$ ,  $sp^3$ ,  $sp^3$ [AMU 1999] (d)  $sp^3, dsp^2, sp^3$ (c)  $sp^3, dsp^2, dsp^2$ (a)  $d^2sp^3$ (b)  $dsp^2$ (At. no. of Ni = 28) (c)  $sp^3d^2$ The bond in  $K_4[Fe(CN)_6]$  are: [MP PET 2004] 28. (a) All ionic The example of  $dsp^2$  hybridisation is All covalent (b) [MP PET 1999; AlIMS 2001]

lonic and covalent

29.

lonic, covalent and co-ordiante covalent Hybridization of Fe in  $K_3Fe(CN)_6$  is [DCE 2002]

4

5.

(a)  $Fe(CN)_6^{3-}$ 

(c)  $Zn(NH_3)_4^{2+}$ 

(b)  $Ni(CN)_{4}^{2-}$ 

(d)  $FeF_6^{3-}$ 

	916 Co-ordination Chemistry		
30.	(a) $sp^3$ (b) $dsp^3$ (c) $sp^3d^2$ (d) $d^2sp^3$ The complex ion which has no ' $d$ ' electrons in the central metal	43.	(c) $d^3$ (in weak as well as in strong fields) (d) $d^5$ (in strong ligand field) The geometry of $Ni(CO)$ and $Ni(PPh)$ $Cl$ are
	atom is [IIT-JEE Screening 2001] (a) $[MnO_4]^-$ (b) $[Co(NH_3)_6]^{3+}$ (c) $[Fe(CN)_6]^{3-}$ (d) $[Cr(H_2O)_6]^{3+}$ Which of the following statement is correct		<ul> <li>(a) Both square planer</li> <li>(b) Tetrahedral and square planar respectively</li> <li>(c) Both tetrahedral</li> <li>(d) Square planar and tetrahedral respectively</li> </ul>
31.	(a) $\left[Cu(NH_3)_6\right]^{2+}$ is a colourless ion		., .
			Complexes and complex stability
	(b) $\left[Zn(H_2O)_6\right]^{2+}$ ion is blue coloured (c) $\left[Ni(CN)_4\right]^{2-}$ ion has a tetrahedral shape	1.	Which of the following complexes has the highest stability constant at 298 K
32.	(d) Nickel dimethyl glyoxides is red in colour Which of the following shall form an octahedral complex		(a) $[CdCl_4]^{2-}$ (b) $[CdBr_4]^{2-}$
	[DCE 2001] (a) $d^4$ (low spin) (b) $d^8$ (high spin)	•	(c) $\left[CdI_4\right]^{2-}$ (d) $\left[Cd\left(CN\right)_4\right]^{2-}$ The most stable ion is
22	(c) $d^6$ (low spin) (d) None of these Which one of the following is a strong field ligand	2.	The most stable ion is [AIEEE 2002] (a) $[Fe(OH)_3]^{3-}$ (b) $[Fe(Cl)_6]^{3-}$
33.	(a) $CN^-$ (b) $NO_2^-$	3.	(c) $[Fe(CN)_6]^{3-}$ (d) $[Fe(H_2O)_6]^{3+}$ The most stable complex among the following is
	(c) $en$ (d) $NH_3$		[MP PMT 2002]
34.	The strongest ligand in the following is [MP PET 1995]		(a) $K_3[Al(C_2O_4)_3]$ (b) $[Pt(en)_2]Cl_2$
	(a) $CN^-$ (b) $Br^-$		(c) $Ag(NH_3)_2Cl$ (d) $K_2(Ni(EDTA)]$
35. 36.	(c) $HO^-$ (d) $F^-$ The neutral ligand is  (a) Chloro (b) Hydroxo  (c) Ammine (d) Oxalato  The ligands which can get attached to the central metal ion through more than one atom are called	4.	Which of the following factors tends to increase the stability of metal ion complexes  (a) Higher ionic radius of the metal ion  (b) Higher charge/size ratio of the metal ion  (c) Lower ionisation potential of the metal ion  (d) Lower basicity of the ligand
	(a) Ambident ligands (b) Polydentate ligands	5.	$CuSO_4$ reacts with KCN solution and forms: [DPMT 2004]
	(c) Chelate ligands (d) Neutral ligands		(a) $K_3[Cu(CN)_4]$ (b) $Cu(CN)$
37.	A strong ligand gives a complex which is generally called  (a) High spin  (b) High energy		(c) $Cu(CN)_2$ (d) $K_4[Cu(CN)_6]$
38.	(c) Low spin (d) Stable $CN^-$ is a strong field ligand. This is due to the fact that	6.	A co-ordination complex compound of cobalt has the molecular formulae containing five ammonia molecules, one nitro group and two chlorine atoms for one cobalt atom. One mole of this compound produces three mole ions in an aqueous solution on reacting with
	(a) It can accept electron from metal species (b) It forms high spin complexes with metal species		excess of $AgNO_3$ , $AgCl$ precipitate. The ionic formula for this complex would be:
	<ul><li>(c) It carries negative charge.</li><li>(d) It is a pseudohalide</li></ul>		[DPMT 2004; Kerala PMT 2004] (a) $[Co(NH_3)_5(NO_2)]Cl_2$
39.	Considering $H_2O$ as a weak field ligand, the number of unpaired		(a) $[Co(NH_3)_5(NO_2)]Cl_2$ (b) $[Co(NH_3)_5Cl][Cl(NO_2)]$
	electrons in $[Mn(H_2O)_6]^{2+}$ will be (At. No. of $Mn=25$ ) asas a		(c) $[Co(NH_3)_4(No_2)Cl][(NH_3)Cl]$
	ajkk [CBSE PMT 2004]		(d) $[Co(NH_3)_5][(NO_2)_2Cl_2]$
	(a) Two (b) Four (c) Three (d) Five	7.	In any ferric salt, on adding potassium ferrocyanide, a prussian blue colour is obtained, which is
40.	Which of the following is a $\pi$ - complex  (a) Trimethyl aluminium (b) Ferrocene		[BIT 1992; BHU 2002]

[KCET 1996; AIIMS 2003] 8.

 $NH_3$  (b) CO

(a)  $NH_3$ 

(c) Diethyl zinc

( )

(d) Nickel carbonyl

(c) F

41.

(d) Ethylene diamine

**42.** The value of the 'spin only' magnetic moment for one of the following configurations is 2.84 BM. the correct one

[AIEEE 2005]

(a)  $d^4$  (in strong ligand field)

Which of the following is a  $\,\pi-$ acid ligand

(b)  $d^4$  (in weak ligand field)

(a)  $K_3 Fe(CN)_6$ 

(b)  $KFe[Fe(CN)_6]$ 

(d)  $FeSO_4.Fe(CN)_6$ 

(d)  $Fe_4[Fe(CN)_6]_3$ 

**8.** Prussian blue is formed when

[CBSE PMT 1990]

(a) Ferrous sulphate reacts with  $FeCl_3$ 

(b) Ferric sulphate reacts with  $K_4[Fe(CN)_6]$ 

(c) Ferrous ammonium sulphate reacts with  $\ensuremath{\mathit{FeCl}}_3$ 

(d) Ammonium sulphate reacts with  $\ensuremath{\mathit{FeCl}}_3$ 

9. Complex salt can be made by the combination of  $[Co^{III}(NH_3)_5\,Cl]^X \ \ \text{with} \qquad \qquad [\text{RPMT 2000, AFMC 2002}]$ 



- (a)  $PO_4^{3-}$
- $Cl^{-}$ (b)
- (c)  $2Cl^{-}$
- (d)  $2K^{+}$
- 10. Which reagent can be used to identify nickel ion
  - (a) Resorcinol
  - (b) Dimethyl glyoxime [DMG]
  - (c) Diphenyl benzidine
  - (d) Potassium ferrocyanide
- Dimethyl glyoxime forms a coloured complex with 11.
  - (a) Ag
- (b) *Ni*
- (c) Cr
- (d) Zn
- Silver chloride dissolves in excess of  $NH_4OH$  . The cation present 12. in this solution is [EAMCET 1998]
  - (a)  $Ag^+$
- (b)  $[Ag(NH_3)_2]^+$
- $[Ag(NH_3)_4]^+$
- (d)  $\left[Ag(NH_3)_6\right]^4$
- Silver sulphide dissolved in a solution of sodium cynamide to form 13. [AMU 1999]
  - (a)  $Na[Ag(CN)_2]$
- (b)  $Na_3[Ag(CN)_4]$
- (c)  $Na_5[Ag(CN)_6]$
- (d)  $Na_2[Ag(CN)_2]$
- Which one will give  $Fe^{3+}$  ions in solution 14.
  - (a)  $\left[ Fe(CN)_{\epsilon} \right]^{3}$
  - (b)  $Fe_2(SO_4)_3$
  - (c)  $[Fe(CN)_{\epsilon}]^{4-}$
  - (d)  $NH_4(SO_4)_2$ .  $FeSO_4$ .  $6H_2O$
- 15. The cation that does not form an amine complex with excess of
  - (a)  $Cd^{2+}$
- (b)  $Al^{3+}$
- $Cu^{2+}$ (c)
- (d)  $Ag^+$

## Application of organometallics

- Ziegler-Natta catalyst is an organometallic compound of which metal
- (b) Zirconium
- (c) Rhodium
- (d) Titanium
- In the case of small cuts, bleeding is stopped by applying potash 2. alum. Here alum acts as

[KCET (Med.) 2001]

- (a) Fungicide
- (b) Disinfectant
- (c) Germicide
- (d) Coagulating agent
- The formula of ferrocene is 3.
  - (a)  $\left[ Fe(CN)_6 \right]^{4-}$
- (b)  $[Fe(CN)_6]^{3-}$
- (c)  $[Fe(CO)_5]$
- (d)  $[(C_5H_5), Fe]$
- Which of the following is formed when n-butyl lithium reacts with tin (11) chloride [AFMC 2001]
  - (a) LiBr
- (b)  $Et_4Pb$
- (c)  $(C_4H_9)_4Sn$
- (d)  $(C_2H_5)_4Pb$
- Which of the following organo-silicon compound on hydrolysis will give a three dimensional silicone

[Orissa JEE 2003]

- (a)  $R_3SiCl$
- (b) RSiCl<sub>3</sub>
- (c) SiCl<sub>4</sub>
- (d)  $R_2SiCl_2$
- Which one is not an organometallic compound 6.

[] & K CET 2005; Pb. CET 2003]

- (a) RMgX
- (b)  $C_2H_5ONa$

- $(CH_3)_4 Sn$ (c)
- (d)  $KC_AH_O$
- The complex used as an anticancer agent is 7.

[AIIMS 2003]

- (a) trans  $-[Co(NH_3)_3 Cl_3]$  (b) cis  $-[PtCl_2(NH_3)_2]$
- (c)  $cis K_2[PtCl_2Br_2]$
- (d)  $Na_2CO_3$
- The compound that is not olefinic organometallic is 8.
  - (a)  $K[C_2H_4PtCl_3]3H_2O$  (b)  $Be(CH_2)_2$
  - (c)  $(C_2H_4PtCl_3)_2$
- (d)  $C_4H_4Fe(CO)_3$
- Among the following, which is not the  $\pi$ -bonded organometallic 9. compound [CBSE PMT 2003]
  - (a)  $(CH_3)_4 Sn$
- (b)  $K[PtCl_3(\eta^2 C_2H_4)]$
- (c)  $Fe(\eta^5 C_5H_5)_2$

11.

16.

18.

- (d)  $Cr(\eta^6 C_6H_6)_2$
- Wilkinson's catalyst is used in 10.
  - (a) Polymerization
- (b) Condensation (d) Hydrogenation
- (c) Halogenation
- What is the use of tetraethyl lead (a) As a catalyst in addition reaction of alkenes
- (b) As a catalyst in polymerization reaction of alkenes
- (c) For reducing knocking
- (d) For creating knocking
- 12. Which of the following is an organo-metallic compound

[MP PMT 2001]

- (a) Lithium ethoxide
- (b) Ethyl lithium
- (c) Lithium acetate
- (d) Lithium carbide
- Which of the following is an organometallic compound 13.

[AIIMS 1997]

- (a)  $Ti(C_2H_5)_4$
- (b)  $Ti(OC_2H_5)_4$
- (c)  $Ti(OCOCH_3)_4$
- (d)  $Ti(OC_6H_5)_4$
- 14. Which of the following is not an organometallic compound

[MP PET 1996; BHU 2002]

- (a) Ethyl magnesium bromide
- Tetraethyl lead
- (c) Sodium ethoxide
- Trimethyl aluminium
- An organometallic compound amongst the following is 15.
  - Ferrocene [J & K CET 2005]
- (b)  $CaC_2$
- Tetraethyl lead (TEL)
- (d) All of these
- Which of the following does not have a metal carbon bond

[CBSE PMT 2004]

- (a)  $K[Pt(C_2H_4)Cl_3]$
- (b)  $Ni(CO)_A$
- (c)  $Al(OC_2H_5)_3$
- (d)  $C_2H_5MgBr$
- Coordination compounds have great importance in biological 17. systems. In this context which of the following statements is [AIEEE 2004] incorrect
  - Cyanocobalamin is  $B_{12}$  and contains cobalt.
  - Haemoglobin is the red pigment of blood and contains iron.
  - Chlorophylls are green pigments in plants and contains calcium.
  - Carboxypepticase-A is an enzyme and contains zinc. Ziegler-Natta catalyst is
    - [Pb. CET 2004] (b)  $K[PtCl_3(C_2H_4)]$
  - (a)  $(Ph_3P)_3RhCl$ (c)  $[Al_2(C_2H_6)_6] + TiCl_4$
- (d)  $[Fe(C_2H_5)_2]$
- Dimethyl glyoxime gives a red precipitate with  $Ni^{2+}$ , which is used for its detection. To get this precipitate readily the best pH range is [AIIMS 2004]
  - (a) < 1

- (b) 2-3
- (c) 3-4
- (d) 9-11
- The  $\pi$ -bonded organo metallic compound which has ethene as 20. one of its component is [] & K CET 2005]
  - (a) Zeise's salt
- (b) Ferrocene



- Dibenzene chromium
- (d) Tetraethyl tin.

# Critical Thinking

## Objective Questions

- The coordination number and oxidation state of  $K_3[Cr(C_2O_4)_3]$  are, respectively [CBSE PMT 1995]
  - (a) 4 and +2
- (b) 6 and +3
- (c) 3 and +3
- (d) 3 and 0
- The complex compounds which result from the coordination of [BHU 1999] carbon monoxide are known as
  - (a) Electronic
- (b) Carbonvls
- (c) Carbonates
- (d) Carbon permono
- Mixture X = 0.02 mol of  $[Co(NH_3)_5 SO_4]Br$  and 0.02 mol of 3.  $[Co(NH_3)_5Br]SO_4$  was prepared in 2 litre of solution
  - 1 *litre* of mixture X + excess  $AgNO_3 \rightarrow Y$ .
  - 1 *litre* of mixture  $X + \text{ excess } BaCl_2 \rightarrow Z$

Number of moles of Y and Z are

(a) 0.01, 0.01

[IIT JEE 2003] (b) 0.02, 0.01

- (c) 0.01, 0.02
- (d) 0.02, 0.02
- Which of the following organometallic compound is  $\sigma$  and  $\pi$ bonded [MH CET 2001; PCET 2002]
  - (a)  $Fe(CH_3)_3$
  - (b)  $[Fe(\eta^5 C_5H_5)_2]$
  - (c)  $[Co(CO)_5 NH_2]^{2+}$
  - (d)  $K[PtCl_3(n^2 C_2H_4)]$
- In  $Ni(NH_3)_4$   $SO_4$ , the valency and coordination number of Niwill be respectively
  - (a) 3 and 6
- (b) 2 and 4
- (c) 4 and 2
- (d) 4 and 4
- Co-ordination number of Fe in the complexes  $[Fe(CN)_6]^{4-}$ , 6.  $[Fe(CN)_6]^{3-}$  and  $[FeCl_4]^-$  would be respectively
  - (a) 2, 3, 3
- (b) 6, 6, 4
- (c) 6, 3,3
- (d) 6, 4, 6
- On hydrolysis  $(Me)_2 SiCl_2$  will produce 7.

[IIT-JEE 2003]

- (a)  $(Me)_2 Si(OH)_2$
- (b)  $(Me)_2 Si = O$
- (c)  $-[-O-(Me)_2Si-O-]_n$
- (d)  $Me_2SiCl(OH)$
- Which of the following represents a chelating ligand 8.

[JIPMER 2002]

- (a)  $H_2O$
- (b) OH-
- (c) DMG
- (d)  $Cl^-$
- $CuSO_4$  reacts with KCN solution and forms 9.

[MP PMT 1992; IIT 1996; UPSEAT 2001, 02]

- (a)  $Cu(CN)_2$
- (b) Cu(CN)
- (c)  $K_2[Cu(CN)_4]$
- (d)  $K_3 \left[ Cu(CN)_4 \right]$
- In the process of extraction of gold, 10.

Roasted gold ore  $+CN^- + H_2O \xrightarrow{O_2} [X] + OH^-$ 

 $[X] + Zn \rightarrow [Y] + Au$ 

Identify the complexes [X] and [Y][IIT-IEE 2003]

- (a)  $X = [Au(CN)_2]^-, Y = [Zn(CN)_4]^{2-}$
- (b)  $X = [Au(CN)_A]^{3-}, Y = [Zn(CN)_A]^{2-}$
- $X = [Au(CN)_2]^-, Y = Zn(CN)_6]^{4-}$
- (d)  $X = [Au(CN)_4]^-, Y = [Zn(CN)_4]^{2-}$
- A solution of potassium ferrocyanide would contains ..... ions

(b) 3

(c) 4

- (d) 5
- Which one of the following complexes is an outer orbital complex
  - (a)  $[Co(NH_3)_6]^{3+}$
- (b)  $[Mn(CN)_6]^{4-}$
- (c)  $[Fe(CN)_6]^{4-}$
- (d)  $[Ni(NH_3)_6]^{2+}$

Atomic nos : Mn = 25, Fe = 26, Co = 27, Ni = 28

Which one of the following has largest number of isomers?

[AIEEE 2004]


- (a)  $[Ir(PP_3)_2H(CO)]^{2+}$
- (b)  $[Co(NH_3)_5 Cl]^{2+}$
- (c)  $[Ru(NH_3)_4 Cl_2]^+$
- $[Co(en), Cl_2]^{\dagger}$


(R = alkyl group; en = ethylenediamine)


- Which kind of isomerism is exhibited by octahedral  $Co(NH_3)_4 Br_2 Cl$  ? [IIT-JEE Screening 2005]
  - (a) Geometrical and Ionization
  - Geometrical and Optical
  - Optical and Ionization
  - Geometrical only (d)
- Which one of the following is expected to exhibit optical isomerism 15. (en = ethylenediamine)
  - (a)  $cis [Pt(NH_3)_2 Cl_2]$
- (b)  $trans [Co(en)_2 Cl_2]$
- (c)  $trans [Pt(NH_3)_2 Cl_2]$
- (d)  $cis [Co(en)_2 Cl_2]$
- $[EDTA]^{4-}$  is a: 16.
- [UPSEAT 2004] (b) Bidentate ligand
- (a) Monodentate ligand (c) Quadridentate ligand
- (d) Hexadentate ligand
- Which Mheero 1993 in statements is incorrect? 17.

[KCET 2004]

- (a) In  $K_3[Fe(CN)_6]$ , the ligand has satisfied only the secondary valency of ferric ion.
- (b) In  $K_3[Fe(CN)_6]$ , the ligand has satisfied both primary and secondary valencies of ferric ion.
- (c) In  $K_4[Fe(CN)_6]$ , the ligand has satisfied both primary and secondary valencies of ferrous ion.
- (d) In  $[Cu(NH_3)_4]SO_4$ , the ligand has satisfied only the secondary valency of copper.
- Which of the following is considered as an anticancer species. 18.









An aqueous solution of CoCl2 on addition of excess of 19. concentrated HCl turns blue due to formation of



#### [AIIMS 2005]

- (a)  $[Co(H_2O)_4 Cl_2]$
- (b)  $[Co(H_2O)2Cl_4]^{2-}$
- (c)  $[CoCl_4]^{2-}$
- (d)  $[Co(H_2O)2Cl_2]$
- **20.** The correct order for the wavelength of absorption in the visible region is [AllMS 2005]
  - (a)  $[Ni(NO_2)_6]^{4-} < [Ni(NH_3)_6]^{2+} < [Ni(H_2O)_6]^{2+}$
  - (b)  $[Ni(NO_2)_6]^{4-} < [Ni(H_2O)_6]^{2+} < [Ni(NH_3)_6]^{2+}$
  - (c)  $[Ni(H_2O)_6]^{2+} < [Ni(NH_3)_6]^{2+} < [Ni(NO_2)_6]^{4-}$
  - (d)  $[Ni(NH_3)_6]^{2+} < [Ni(H_2O)_6]^{2+} < [Ni(NO_2)_6]^{4-}$
- In which of the following pairs both the complexes show optical isomerism [AIIMS 2005]
  - (a)  $cis-[Cr(C_2O_4)_2Cl_2]^{3-}$ ,  $cis-[Co(NH_3)_4Cl_2]$
  - (b)  $[Co(en)_3]Cl_3$ , cis- $[Co(en)_2Cl_2]Cl$
  - (c) [PtCl(dien)]Cl,  $[NiCl_2Br_2]^{2-}$
  - (d)  $[Co(NO_3)_3(NH_3)_3]$ , cis- $[Pt(en)_2Cl_2]$
- 22. Which of the following compounds shows optical isomerism

#### [AIEEE 2005; CBSE PMT 2005]

11.

- (a)  $[Cu(NH_3)_4]^{2+}$
- (b)  $[ZnCl_4]^2$
- (c)  $[Cr(C_2O_4)_3]^{3-}$
- (d)  $[Co(CN)_6]^{3-}$



# Assertion & Reason

### For AIIMS Aspirants

Read the assertion and reason carefully to mark the correct option out of the options given below :

- (a) If both assertion and reason are true and the reason is the correct explanation of the assertion.
- (b) If both assertion and reason are true but reason is not the correct explanation of the assertion.
- (c) If assertion is true but reason is false.
- (d) If the assertion and reason both are false.
- (e) If assertion is false but reason is true.
- Assertion : Potassium ferrocyanide and potassium ferricyanite both are diamagnetic.
  - Reason : Both have unpaired electrons.
- **2.** Assertion :  $NF_3$  is a weaker ligand than  $N(CH_3)_3$

Reason :  $NF_3$  ionizes to give  $F^-$  ions ionsiionsionsioiiioioioioioioioioioio

ions in aqueous solution.

Reason

- 3. Assertion : The  $[Ni(en)_3]Cl_2$  (en = ethylene diamine) has lower stability than  $[Ni(NH_3)_6]Cl_2$ 
  - : In  $[Ni(en)_3]Cl_2$  the geometry of Ni is trigonal
    - bipyramidal
- 4. Assertion : The ligands nitro and nitrito are called ambidentate
  - ligands.
  - Reason : These ligands give linkage isomers.
- **5.** Assertion : Geometrical isomerism is also called *cis-trans* 
  - isomerism.
  - Reason : Tetrahedral complexes show geometrical isomerism.

# **6.** Assertion : $(en)_2 Co \stackrel{NH}{C} \rho(en)_2$ is named as tetrak

(ethylene diamine)  $\mu$  -hydroxo-imido dicobalt (III) ion

Reason : In naming polynuclear complexes *i.e.*, containing two or more metal atoms joined by bridging ligands, the word  $\mu$  is added with hyphen before the name of

such ligands.

7. Assertion :  $[Fe(CN)_6]^{3-}$  is weakly paramagnetic while  $[Fe(CN)_6]^{4-}$  is diamagnetic.

Reason :  $[Fe(CN)_6]^{3-}$  has +3 oxidation state while  $[Fe(CN)_6]^{4-}$  has +2 oxidation state.

**8.** Assertion :  $H_2N - NH_{2}$  is a chelating ligand.

Reason : A chelating ligand must possess two or more lone pairs at such a distance that it may form suitable strain free rings at the metal ion.

**9.** Assertion :  $[Ti(H_2O)_6]^{3+}$  is coloured while  $[Sc(H_2O)_6]^{3+}$  is

Reason : d-d transition is not possible in  $[Sc(H_2O)_6]^{3+}$ .

10. Assertion : All the octahedral complexes of  $Ni^{2+}$  must be outer orbital complexes.

Reason : Outer orbital octahedral complexes are given by weak ligands.

Assertion : Potassium ferrocyanide is diamagnetic whereas potassium fericyanide is paramagnetic.

Reason : Crystal field splitting in ferrocyanide ion is greater

than that of ferricyanide ion.

[AIIMS 2005]



#### **Basic Terms**

1	b	2	С	3	b	4	b	5	d
6	а	7	b	8	С	9	b	10	С
11	d	12	С	13	d	14	b	15	b
16	b	17	а	18	С	19	С	20	b
21	b	22	С	23	b	24	d	25	а
26	d	27	а	28	b	29	b	30	С
31	b	32	С	33	b	34	b	35	d
36	d	37	b	38	b	39	d	40	b
41	d	42	b	43	d	44	b	45	а
46	а	47	d	48	С	49	b	50	d
51	а	52	b	53	d	54	а	55	d
56	d	57	a	58	d	59	а	60	d
61	С	62	d						

#### Nomenclature, oxidation State and EAN number

1	h	2	·	3	C	1	h	5	Ч
	U		·	J	·	-	U	J	u



6	С	7	b	8	а	9	а	10	а
11	С	12	a	13	b	14	a	15	b
16	С	17	d	18	b	19	d	20	а
21	b	22	b	23	а	24	b	25	а
26	d	27	С	28	b	29	а	30	а
31	а	32	b	33	а	34	b	35	С
36	b	37	а	38	а	39	b	40	С
41	а	42	b	43	С	44	b	45	b
46	d	47	а	48	b	49	С	50	С
51	b	52	b	53	С	54	d	55	b
56	а	57	b	58	С				

# **Isomerism and Magnetic properties**

1	а	2	d	3	а	4	С	5	С
6	d	7	b	8	а	9	а	10	С
11	d	12	d	13	b	14	С	15	а
16	b	17	а	18	d	19	а	20	b
21	С	22	b	23	b	24	а	25	а
26	d	27	d	28	а	29	С	30	а
31	b	32	а	33	С	34	а	35	d
36	С	37	d	38	а	39	d	40	b
41	а	42	С	43	d	44	а	45	b
46	е	47	а	48	d	49	С	50	С
51	d	52	а	53	С	54	а	55	а
56	С	57	С						

# **Hybridisation and Geometry**

1	а	2	b	3	а	4	b	5	а
6	b	7	b	8	d	9	С	10	а
11	b	12	С	13	d	14	а	15	b
16	b	17	d	18	d	19	а	20	а
21	b	22	d	23	С	24	С	25	С
26	d	27	d	28	d	29	d	30	а
31	d	32	С	33	а	34	а	35	С
36	а	37	С	38	d	39	d	40	b
41	b	42	а	43	С				

# Complexes and complex stability

1	d	2	d	3	d	4	b	5	a
6	а	7	d	8	b	9	С	10	b
11	b	12	b	13	а	14	b	15	b

# Application of organometallics

1	d	2	d	3	d	4	С	5	b
6	b	7	b	8	а	9	а	10	d
11	С	12	b	13	а	14	С	15	d
16	С	17	С	18	С	19	d	20	а

# **Critical Thinking Questions**

1	b	2	b	3	а	4	b	5	b
6	b	7	С	8	С	9	d	10	а
11	d	12	d	13	d	14	а	15	d
16	d	17	ad	18	а	19	С	20	a
21	b	22	С						

## **Assertion & Reason**

1	d	2	С	3	d	4	а	5	С
6	е	7	b	8	е	9	а	10	b
11	С								