

Class 11

2017-18



CHEMISTRY

FOR JEE MAIN & ADVANCED

SECOND
EDITION



Topic Covered
Stoichiometry and
Redox Reactions

Exhaustive Theory ◀
(Now Revised)

Formula Sheet ◀

9000+ Problems ◀
based on latest JEE pattern

2500 + 1000 (New) Problems ◀
of previous 35 years of
AIEEE (JEE Main) and IIT-JEE (JEE Adv)

5000+ Illustrations and Solved Examples ◀

Detailed Solutions ◀
of all problems available

Planceess Concepts

Tips & Tricks, Facts, Notes, Misconceptions,
Key Take Aways, Problem Solving Tactics

Planceessential

Questions recommended for revision




1. STOICHIOMETRY AND REDOX REACTIONS

THE MOLE AND EQUIVALENT CONCEPT

1. INTRODUCTION



Some important terminologies required to master the concepts of chemistry are as follows:

- (a) **Chemical Equation:** It is the symbolic representation of a true chemical reaction. The equation provides qualitative and quantitative information about a chemical change in a simple manner. For e.g. in the reaction,



The '+' sign on left hand side means 'react with' and on the right of arrow as 'produces'. The substances which react are called 'reactants' and those produced in the reaction are called 'products' and they are represented on the LHS and RHS of the arrow respectively.

- (b) **Thermo Chemical Equation:** The equations which represent chemical as well as thermal changes are called thermo chemical equations. The energy change is shown by putting a value on the RHS of the reaction. For eg.



- (c) **Molecule and Atom:** A molecule is defined as the smallest possible particle of a substance which has all the properties of that substance and can exist freely in nature. On the other hand an atom is the smallest particle of an element (made up of still smaller particles like electrons, protons, neutrons, etc.) which can take part in a chemical reaction. It may or may not exist free in nature.
- (d) **Atomicity of an Element:** The term refers for the number of atoms present in one molecule of an element, e.g., atomicity of H_2 , O_3 , S_8 , P_4 is 2, 3, 8 and 4 respectively.
- (e) **Mole:** A mole is defined as the amount of matter that contains as many objects (atoms, molecules, electron, proton or whatever, objects we are considering) as the same number of atoms in exactly 12 g of C^{12} . This number is also known as **Avogadro's number (N_A)**. Avogadro's number = 6.023×10^{23} molecule/mole. Thus, 1 mole of an entities contains N_A particles of that entity.

Number of objects (N) in n mole = number of mole (n) \times number of objects per mole (N_A)

$$N = n \times N_A = \frac{w}{M} \times N_A, \left[n = \frac{w}{M} \right] \text{ where } N_A \text{ is Avogadro's number.}$$

- (f) **Molar Mass:** The mass in gram of 1 mole of a substance is known as molar mass or molecular weight of substance.

- (g) **Atomic Weight:** The atomic mass of an element is defined as the average relative mass of its atoms as compared to the mass of a carbon atom taken as 12. Note that the atomic weight of an element is a relative weight of one atom and not the absolute weight.
- (h) **Gram Molecular Weight:** Molecular weight or gram molecular weight of a substance is weight of N_A -molecules of that substance in grams.

Illustration 1: Calculate the mass of single atom of sulphur and a single molecule of carbon dioxide.

(JEE MAIN)

Sol: Know the gram atomic and the molecular mass of sulphur and CO_2 and then divide by N_A .

$$\text{Mass of one sulphur atom} = \frac{\text{Gram atomic mass}}{6.02 \times 10^{23}} = \frac{32}{6.02 \times 10^{23}} = 5.33 \times 10^{-23} \text{ g}$$

$$\text{Formula of carbon dioxide} = \text{CO}_2$$

$$\text{Molecular mass of CO}_2 = 12 + 2 \times 16 = 44$$

$$\text{Gram-molecular mass of CO}_2 = 44 \text{ g}$$

$$\text{Mass of one molecule of CO}_2 = \frac{\text{Gram molecular mass}}{6.02 \times 10^{23}} = \frac{44}{6.02 \times 10^{23}} = 7.308 \times 10^{-23} \text{ g}$$

Illustration 2: How many electrons are present in 1.6 g of methane?

(JEE ADVANCED)

Sol: Know the molar mass of methane, the moles and the no. of molecules. Then calculate no. of electrons for the solved no. of molecules.

$$\text{Gram molecular mass of methane, (CH}_4\text{)} = 12 + 4 = 16 \text{ g}$$

$$\text{Number of moles in 1.6 g of methane} = \frac{1.6}{16} = 0.1$$


$$\text{Number of molecule of methane in 0.1 mole} = 0.1 \times 6.023 \times 10^{23} = 6.02 \times 10^{22}$$

$$\text{One molecules of methane has} = 6 + 4 = 10 \text{ electrons}$$

$$\text{So, } 6.02 \times 10^{22} \text{ molecules of methane have} = 10 \times 6.02 \times 10^{22} \text{ electrons} = 6.02 \times 10^{23} \text{ electrons}$$

2. STOICHIOMETRY

The quantitative aspect, dealing with mass and volume relations among reactants and products is termed stoichiometry. Consider for example, the reaction represented by a balanced chemical equation:



$$\text{Weight ratio : } 4\text{g} \quad 32\text{g} \quad 36\text{g}$$

$$\text{Volume ratio : } 2 \text{ vol} \quad 1 \text{ vol} \quad 2 \text{ vol} \quad (\text{valid only for gaseous state at same P and T})$$

The given reaction suggests the combination ratio of reactants and formation ratio of products in terms of:

- (a) **Mole ratio:** 2 mol H_2 reacts with 1 mol of O_2 to form 2 mol of H_2O vapors.
- (b) **Molecular ratio:** 2 molecule of H_2 reacts with 1 molecule of O_2 to form 2 molecules of H_2O vapors.
- (c) **Weight ratio:** 4 g H_2 reacts with 32 g O_2 to form 36 g of H_2O vapors.

- (d) **Volume ratio:** In gaseous state 2 volume H_2 reacts with 1 volume O_2 to form 2 volume H_2O vapors at same conditions of P and T.

Therefore, coefficients in the balanced chemical reaction can be interpreted as the relative number of moles, molecules or volume (if reactants are gases) involved in the reaction. These coefficients are called stoichiometrically equivalent quantities and may be represented as:



Or Mole of H_2 : Mole of O_2 : Mole of H_2O = 2: 1: 2


Where the symbol \equiv is taken to mean 'stoichiometrically equivalent to'. The stoichiometric relation can be used to give conversion factors for relating quantities of reactants and products in a chemical reaction.

2.1 Some Important Laws

- (a) **Law of conservation of mass:** "In all physical and chemical changes, the total mass of the reactants is equal to that of the products" or "matter can neither be created nor destroyed."
- (b) **Law of constant composition/definite proportion:** "A chemical compound is always found to be made up of the same elements combined together in the same fixed ratio by weight".
- (c) **Law of multiple proportions:** "When two elements combine together to form two or more chemical compounds, then the weight of one of the elements which combine with a fixed weight of the other bear a simple ratio to one another".
- (d) **Law of reciprocal proportions:** The ratio of the weights of two elements A and B which combine with a fixed weight of the third element C is either the same or a simple multiple of the ratio of the weights of A and B which directly combine with each other.
- (e) **Gay-Lussac's law of gaseous volumes:** "When gases react together, they always do so in volumes which bear a simple ratio to one another and to the volumes of the products, if gaseous, all measurements are made under the same conditions of temperature and pressure".

Illustration 3: What mass of sodium chloride would be decomposed by 9.8 g of sulphuric acid, if 12 g of sodium bisulphate and 2.75 g of hydrogen chloride were produced in a reaction assuming that the law of conservation of mass is true? **(JEE MAIN)**

Sol: Apply the law of conservation of mass.



According to law of conservation of mass, Total mass of reactant = Total mass of product

Let the mass of NaCl decomposed be x, so


$$x + 9.8 = 12 + 2.75$$

$$= 14.75$$

$$x = 4.95 \text{ g}$$

Illustration 4: How much volume of oxygen will be required for complete combustion of 40 mL of acetylene (C_2H_2) and how much volume of carbon dioxide will be formed? All volumes are measured at NTP. **(JEE ADVANCED)**

Solution: Write the balanced chemical reaction and from the given data determine the volume.



So, for complete combustion of 40 mL of acetylene, 100 mL of oxygen are required and 80 mL of carbon dioxide is formed.

2.2 Avogadro's Hypothesis

"Equal volumes of all gases/vapors under similar conditions of temperature and pressure contain equal number of molecules."

This statement leads to the following facts:

- (a) One mole of all gases contain Avogadro's number of molecules, i.e., 6.023×10^{23} molecules.
- (b) The volume of 1 mole of gas at NTP or STP is 22.4 litre.
- (c) NTP or STP refers for $P = 1 \text{ atm}$, $T = 0^\circ\text{C}$ or 273 K .
- (d) Molecular weight = $2 \times$ vapour density (for gaseous phase only)

It provides a method to determine the atomic weights of gaseous elements.

2.3 Dulong and Petit's Law

This law is valid for metals only. According to this law, atomic weight \times specific heat (in cal/g) ≈ 6.4 . Also, heavier the element, lesser will be its specific heat. Therefore, $C_{\text{Hg}} < C_{\text{Cu}} < C_{\text{Al}}$.

2.4 Equivalent Weight

For comparing reacting weights of substances participating in a chemical reaction, chemists coined the term 'equivalent weight'. The substances react in their equivalent weight ratios.

2.4.1 Equivalent Weight of an Element or Compound in a Non-Redox Change

Equivalent weight of an element is its weight which reacts with 1 part by weight of hydrogen, 8 parts by weight of oxygen and 35.5 parts by weight of chlorine. This definition leads to following important generalisations.

(a) **Equivalent weight of an element:** $E = \frac{\text{Atomic weight of element}}{\text{Valence of element}}$

(b) **Equivalent weight of an ionic compound:** $(E) = \frac{\text{Formula weight of compound}}{\text{Total charge on cationic or anionic part}}$

Also, Eq. wt. of compound

$$E = \text{Eq. wt. of I part} + \text{Eq. wt. of II part}$$

(c) **Equivalent weight of an acid or base:**

$$E_{\text{Acid}} = \frac{\text{Molecular weight}}{\text{Basicity}}$$

Basicity = Number of H-atoms replaced from one molecule of acid

$$E_{\text{Base}} = \frac{\text{Molecular weight}}{\text{Acidity}}$$

Acidity = Number of OH-groups replaced from one molecule of base.

(d) **Equivalent weight of acid salt:** $= \frac{\text{Molecular weight of acid salt}}{\text{Replaceable H-atom in it}}$

An acid salt is one which has replaceable H-atom, e.g., NaHCO_3 , NaHSO_4 , Na_2HPO_4 , Na_2HPO_3 is not an acid salt, since it does not have replaceable H-atom.

Note: An acid salt possesses acidity as well as basicity both.

(i) **Equivalent weight of basic salt:**
$$= \frac{\text{Molecular weight of basic salt}}{\text{Replacable OH gps in basic salt}}$$

A basic salt is one which has replaceable OH gps e.g., Ca(OH)Cl , $\text{Al(OH)}_2\text{Cl}$, Al(OH)Cl_2 etc.

2.4.2 Equivalent Weight of an Element or Compound in a Redox Change

For a redox change, the equivalent weight of a substance is given by,

$$\text{Equivalent weight of an oxidant or reductant} = \frac{\text{Molecular weight}}{\text{Number of electrons lost or gained by one molecule of oxidant or reductant}}$$

2.4.3 Gram Equivalent Weight

The equivalent weight of a substance expressed in grams is called gram eq. wt. or one gram equivalent. Now we can define gram-equivalent (g meq) in gms of a substance whose equivalent weight is as follows: No. of equivalents = $\frac{g}{E}$

Illustration 5: An unknown element forms an oxide. What will be the equivalent mass of the element if the oxygen content is 20% by mass? **(JEE MAIN)**

Sol: Use the equation of equivalent weight.

$$\text{Equivalent mass of element} = \frac{\text{Mass of element}}{\text{Mass of oxygen}} \times 8 = \frac{80}{20} \times 8 = 32$$

Illustration 6: The equivalent weight of a metal is double than that of oxygen. How many times is the weight of its oxide greater than the weight of metal? **(JEE ADVANCED)**

Sol: First calculate the equivalent weight of the metal and from the molecular formula, determine the ratio.

$$\text{Equivalent mass of metal} = 16 = \frac{x}{n}$$

Where x = atomic mass of metal


N = valency of metal

Molecular formula of metal oxide = M_2O_n

$$\frac{\text{Mass of metal oxide}}{\text{Mass of metal}} = \frac{2(16n) + 16(n)}{2(16n)} = 1.5$$

2.5 The Limiting Reagent

The reagent producing the least number of moles of products is the limiting reagent. For example, consider a chemical reaction given below, containing 10 mol of H_2 and 7 mol of O_2 . Since, 2 mol H_2 reacts with 1 mol O_2 , thus,



It is thus, evident that the reaction stop only after consumption of 5 moles of O_2 since, no further amount of H_2 is left to react with unreacted O_2 . The substance that is completely consumed in a reaction is called **limiting reagent** because it determines or limits, the amount of product. The other reactants present in excess are sometimes called as **excess reagents**.

Calculation of limiting reagent

- By calculating the required amount by the equation and comparing it with given amount. [Useful when only two reactants are there]
- By calculating amount of anyone product obtained taking each reactant one by one irrespective of other reactants. The one giving least product is limiting reagent.
- Divide given moles of each reactant by their stoichiometric coefficient, the one with least ratio is limiting reagent. [Useful when numbers of reactants are more than two].


2.6 Reaction Yield

The theoretical yield of a product is the maximum quantity that can be expected on the basis of stoichiometry of a chemical equation. The percentage yield is the percentage of a theoretical yield actually achieved. The lower yield of a chemical reaction is due to side reactions.

$$\text{Percentage yield} = \frac{\text{Actual yield}}{\text{Theoretical yield}} \times 100$$

Illustration 7: 10 mL N_2 and 25 mL H_2 at same P and T are allowed to react to give NH_3 quantitatively. Predict (i) the volume of NH_3 formed, (ii) limiting reagent. **(JEE MAIN)**

Sol: Frame the reaction and lay down the conditions, due to which volume of NH_3 can be found which leads to the limiting reagent.



Limiting reagent is H_2 .

Illustration 8: A chloride of an element contains 49.5% chlorine. The specific heat of the element is 0.056. Calculate the equivalent mass, valency and atomic mass of the element. **(JEE ADVANCED)**

Sol: Calculate mass of the metal from the given percentage and the equivalent mass of the metal. Using the Dulong and Petit's law, specific heat gives the atomic mass of the metal and then the valency can be found.

Mass of chlorine in the metal chloride = 49.5%

Mass of metal = $(100 - 49.5) = 50.5$

Equivalent mass of the metal = $\frac{\text{Mass of metal}}{\text{Mass of chlorine}} \times 35.5 = \frac{50.5}{49.5} \times 35.5 = 36.21$

According to Dulong and Petit's law,

Approximate atomic mass of the metal = $\frac{6.4}{\text{Specific heat}} = \frac{6.4}{0.056} = 114.3$

$$\text{Valency} = \frac{\text{Approximate atomic mass}}{\text{Equivalent mass}} = \frac{114.3}{36.21} = 3.1 \approx 3$$

Hence, exact atomic mass = $36.21 \times 3 = 108.63$

3. METHODS OF EXPRESSING CONCENTRATION OF SOLUTION

(a) **Strength of Solution:** Amount of solute present in one litre solution

$$S = \frac{\text{Weight of solute}}{\text{Volume of solution in litre}} = \frac{w}{V \text{ in (l)}}$$

$$= \text{Normality} \times \text{Equivalent weight}$$

$$= \text{Molarity} \times \text{Molecular weight}$$

(b) **Mass Percentage or Percent by Mass:**

$$\%(w/w) \text{ Mass percentage of solute} = \frac{\text{Mass of solute}}{\text{Mass of solution}} \times 100$$

(c) **Percent Mass by Volume:** $\%(w/v) = \frac{\text{Mass of solute}}{\text{Volume of solution}} \times 100$

(d) **Parts Per Million (ppm):** $= \frac{\text{Mass of solute}}{\text{Mass of solution}} \times 10^6$

(e) **Molarity:** It is expressed as moles of solute contained in one litre of solution or it is also taken as millimoles of solute in 1000 cc(mL) of solution. It is denoted by M.

$$\text{Molarity} = \frac{\text{Moles of solute}}{\text{Litres of solution}} = \frac{\text{Millimoles of solute}}{\text{Millilitres of solution}}; \quad M = \frac{n_B}{V_{lt}} = \frac{g_B / m_B}{V_{lt}}$$

(f) **Molality:** It is the number of mole present in 1kg solvent.

$$\text{Molality}(m) = \frac{\text{No. of moles of solute}}{\text{Weight (in kg) of solvent}}$$

Let w_A grams of the solute of molecular mass m_A be present in w_B grams of the solvent, then

$$\text{Molality}(m) = \frac{w_A}{m_A \times w_B} \times 1000$$

(g) **Normality:** It is define as number of equivalent of a solute present in one litre of solution.

$$N = \frac{\text{Equivalent of solute}}{\text{Volume of solution in litre}} = \frac{\text{Weight of solute}}{\text{Equivalent weight of solute} \times V \text{ in litre}}$$

$$N = \frac{w}{E \times V \text{ in (l)}} = \frac{w \times 1000}{E \times V \text{ in mL}}$$

Note: A striking fact regarding equivalent and milli equivalent is equivalent and milli equivalent of reactants react in equal number to give same number of equivalent or milli equivalent of products separately.

(h) **Formality:** Since molecular weight of ionic solids is not determined accurately due to their dissociative nature and therefore molecular weight of ionic solid is often referred as formula weight and molarity as formality. Formality = $\frac{\text{Wt. of solute}}{\text{Formula wt.} \times V(\text{in l})}$ i.e., molarity

(i) **Specific Gravity of Solution:**
$$= \frac{\text{Weight of solution}}{\text{Volume of solution}}$$
 i.e., weight of 1 mL solution.

PLANCESS CONCEPTS

- Molality, % by weight, mole fractions are independent of temperature since these involve weights.
- Rest all, i.e., normality, molarity, % by volume, % by strength and strength are temperature dependent, normally decrease with increase in temperature since volume of solution increases with T.
- Molar solution having normality 1N and molarity 1M respectively.
- On diluting a solution, eq. meq. mole or m mole of solute do not change however N and M change.

Aishwarya Karnawat (JEE 2012, AIR 839)

(j) **Mole Fraction:** It is the fractional part of the moles that is contributed by each component to the total number of moles that comprises the solution. In containing n_A moles of solvent and n_B moles of solute. Mole

$$\text{fraction of B} = x_B = \frac{n_B}{n_A + n_B}$$

$$\text{Mole fraction of A} = x_A = \frac{n_A}{n_A + n_B}$$

(k) **Ionic Strength:** The ionic strength (μ) of the solution obtained by mixing two or more ionic compounds is given by: $\mu = \frac{1}{2} \sum cZ^2$. Where c is the concentration (molarity) of that ion and Z is its valence.

Illustration 9: 30 mL of 0.1 N BaCl_2 is mixed with 40 mL of 0.2 N $\text{Al}_2(\text{SO}_4)_3$. How many g of BaSO_4 are formed?

(JEE MAIN)

Sol: Frame the reaction and place the given data to find the milliequivalents at the end of the reaction. Using the formula below, weight of BaSO_4 can be found.

	BaCl_2	+	$\text{Al}_2(\text{SO}_4)_3$	\longrightarrow	BaSO_4	+	AlCl_3
Meq. before	30×0.1		40×0.2		0		0
reaction	= 3		= 8		= 0		= 0
Meq. after reaction	0		5		3		3

3 Meq. of BaCl_2 reacts with 3 Meq. of $\text{Al}_2(\text{SO}_4)_3$ to produce 3 Meq. of BaSO_4 and 3 Meq. of AlCl_3

$$\therefore \text{Meq. of } \text{BaSO}_4 \text{ formed} = \frac{W_{\text{BaSO}_4}}{E_{\text{BaSO}_4}} \times 1000 = 3$$

$$W_{\text{BaSO}_4} \text{ formed} = \frac{3 \times 233}{2 \times 1000} = 0.3495 \text{ g}$$

Illustration 10: 500 mL of aM solution and 250 mL of bM solution of the same solute are mixed and diluted to 2 litre. The diluted solution shows the molarity 1.6 M. If a: b is 2: 5, then calculate a and b. **(JEE ADVANCED)**

Sol: Using the mixture molarity formula $\frac{M_1 \times V_1 + M_2 \times V_2}{V_1 + V_2}$, a and b is calculated.

$$\frac{500 \times a + 250 \times b}{2000} = 1.6; 500a + 250b = 3200$$

If, $\frac{a}{b} = \frac{2}{5}$ then $\frac{500 \times b \times 2}{5} + 250b = 3200$; $450b = 3200$; $b = 7.11$

Similarly, $500a + \frac{250 \times 5a}{2} = 3200$; $\therefore a = \frac{3200}{1125} = 2.84$


4. SOME CHARACTERISTIC APPLICATION OF MOLE CONCEPT

4.1 Gravimetric Analysis

Gravimetric analysis is an analytical technique based on the measurement of mass of solid substances and or volume of gaseous species. Gravimetric analysis is divided into three parts.

- (a) Mass–Mass Relationship:** It relates the mass of a species (reactant or product) with the mass of another species (reactants or products)

Let us consider a chemical reaction,



Suppose the mass of NaHCO_3 being heated is 'x' g and we want to calculate the weight of Na_2CO_3 being produced by heating of 'a' g NaHCO_3 .

The moles of $\text{NaHCO}_3 = \frac{x}{84}$

According to the above balanced equation 2 moles of NaHCO_3 upon heating gives 1 mole of Na_2CO_3

- (b) Mass–Volume Relationship:** It relates the mass of a species (reactant or product) and the volume of a gaseous species (reactant or product) involved in a chemical reaction. Suppose we are provided with 'a' gms of NaHCO_3 in a vessel of capacity VL and the vessel is heated, so that decomposes as $2\text{NaHCO}_3 \xrightarrow{\Delta} \text{Na}_2\text{CO}_3 + \text{H}_2\text{O} + \text{CO}_2$

Now, we want to calculate the volume of CO_2 gas being reduced.

Moles of NaHCO_3 taken $= \frac{x}{84}$

Now, since 2 moles of NaHCO_3 gives 1 mole of CO_2 at STP. Thus

Moles of CO_2 produced $= \frac{1}{2} \times \frac{x}{84}$

As we know that 1 mole of any gas at STP occupies a volume of 22.4 L.

So, volume of CO_2 produced $= \left(\frac{1}{2} \times \frac{x}{84} \times 22.4 \right) \text{L}$

- (i) Volume–Volume Relationship:** It relates the volume of gaseous species (reactants or products) with the volume of another gaseous species (reactant or product) involved in a chemical reaction.

Illustration 11: An ore containing Mn_2O_3 is analysed for the manganese content by quantitatively converting the manganese to Mn_3O_4 and weighing it. A 1.52 g sample of ore yields 0.126 g Mn_3O_4 . Calculate the percent of Mn and Mn_2O_3 in the sample. **(JEE ADVANCED)**

Sol: From the given data, find out the amount of Mn_2O_3 and calculate the %.

Equate the no. of moles of Mn_2O_3 with the no. of moles of Mn and hence find % of Mn. $3\text{Mn}_2\text{O}_3 \longrightarrow 2\text{Mn}_3\text{O}_4$

Mole ratio $\text{Mn}_2\text{O}_3 : \text{Mn}_3\text{O}_4 :: 3 : 2$

$$\therefore \text{Moles of } \text{Mn}_2\text{O}_3 = \frac{3}{2} \times \text{Moles of } \text{Mn}_3\text{O}_4 = \frac{3}{2} \times \frac{0.126}{229} = 5.253 \times 10^{-4}$$

$$\therefore \text{Amount of } \text{Mn}_2\text{O}_3 = 5.253 \times 10^{-4} \times 158 = 0.13 \text{ g}$$

$$\therefore \% \text{ of } \text{Mn}_2\text{O}_3 = \frac{0.13}{1.52} \times 100 = 8.58$$

Also, $\text{Mn}_2\text{O}_3 \longrightarrow 2\text{Mn}$

$$\therefore \text{Mole of Mn} = 2 \times \text{Mole of } \text{Mn}_2\text{O}_3 = 2 \times 5.253 \times 10^{-4} = 10.506 \times 10^{-4}$$

$$\therefore \text{Amount of Mn} = 10.506 \times 10^{-4} \times 55$$

$$\therefore \% \text{ Mn} = \frac{0.09}{1.52} \times 100 = 5.29$$

Illustration 12: A 1.0 g sample of pure organic compound containing chlorine is fused with Na_2O_2 convert chlorine to NaCl . The sample is then dissolved in water, and the chloride precipitated with AgNO_3 , giving 1.96 g of AgCl . If the molecular weight of organic compound is 147, how many chlorine atoms does each molecule contain?

(JEE MAIN)

Sol: Calculate the moles of chloride ions in silver chloride and the organic compound; and compare the ratio.

$$\text{Moles of AgCl} = \text{Moles of chloride} = \frac{1.96}{143.5} = 0.0136$$

$$\text{Moles of organic compound} = \frac{1}{147} = 6.8 \times 10^{-3}$$

$$\text{Chlorine atoms in each molecules of organic compound} = \frac{0.0136}{6.8 \times 10^{-3}} = 2$$

4.2 Volumetric Analysis

It is the process of determination of conc. of a solution with the help of another solution of known conc. It may also be defined as experimental method of determination of volume of a solution of known strength needed for a definite volume of another solution of unknown strength.

Titration: It is an operation forming the basis of volumetric analysis. The addition of measured amount of a solution of one reagent (called the **titrant**) from a burette to a definite amount of another reagent (called **analyte**) until the reaction between them is complete, i.e., till the second reagent (analyte) is completely used up, i.e., upto end point.

Type of Titrations: There are four general classes of volumetric titrations.

- Acid-Base Titration:** Acid or base solutions are titrated against a standard solution of a strong base or strong acid using suitable acid-base indicator.
- Precipitation Titration:** In such titration, the titrant forms an insoluble product with analyte e.g., titration of chloride ions against AgNO_3 solution.
- Complexometric Titrations:** In such titrations, the titrant is a complexing agent and forms a water-soluble complex with the analyte, usually containing a metal ion. The titrant is often a chelating agent, e.g., ethylenediaminetetraacetic acid (EDTA).
- Redox Titrations:** These involves the titration of an oxidizing agent against a reducing agent or vice versa.

Standard Solution: It is the solution of known strength.

Primary Standard Solution: The solution for which conc. is known is called primary standard solution.

Note: For primary standard solution,

- Solute should not be reactive towards solvent or air.

(b) Solute should not be hygroscopic.

(c) Temperature should be constant.

In **acids**, oxalic acid ($\text{H}_2\text{C}_2\text{O}_4 \cdot 2\text{H}_2\text{O}$), benzoic acid ($\text{C}_6\text{H}_5\text{COOH}$), sulphamic acid (HNNH_2SO_3), etc. are taken as primary standard solution.

In **bases**, washing soda ($\text{Na}_2\text{CO}_3 \cdot 10\text{H}_2\text{O}$), borax ($\text{Na}_2\text{B}_4\text{O}_7$), etc are taken as primary standard solution.

In **oxidizing agents**, only potassium dichromate ($\text{K}_2\text{Cr}_2\text{O}_7$) is taken as primary standard solution.

In **reducing agents**, hypo ($\text{Na}_2\text{S}_2\text{O}_3 \cdot 5\text{H}_2\text{O}$), Mohr's salt ($\text{FeSO}_4 \cdot (\text{NH}_4)_2\text{SO}_4 \cdot 6\text{H}_2\text{O}$), sodium oxalate ($\text{Na}_2\text{C}_2\text{O}_4$), etc are taken as primary standard solutions.

End point: End point of titration is normally detected by a sudden change in color of the solution.

Indicator: These compound mixed in the solution in very small amount, which responses the sudden change in color of the solution and show the end point of titration. In acid–base titration, the indicators used are either weak organic acid or weak organic bases. Some examples are

Acidic Indicator: Phenolphthalein, litmus paper etc.

Basic Indicator: Methyl orange, methyl red, etc.

Acid	Base	Indicator
Strong	Strong	Any
Strong	Weak	Methyl orange, methyl red, etc
Weak	Strong	Phenolphthalein etc

Principle of Titration: Titration means stoichiometry and hence its problems may be solved by mole as well as equivalent concept. But for simplicity equivalent concept is preferred, according to which the number of g-equivalents of all reactants reacted will be equal and the same number of g-equivalents of each products will form. The number of g-equivalents of substances may be determined by using the following formulae:

$$\text{Number of g-equivalents} = \frac{\text{Wt. (in gm)}}{\text{Gm. eq.wt.}} = \frac{VS}{1000} = \frac{\text{Vol. of gas}}{\text{Eq. vol.}} = \text{Mole} \times x - \text{factor}$$

Where, S = strength in normality

Illustration 13: 30 mL of a certain solution of Na_2CO_3 and NaHCO_3 , required 12 mL of 0.1 N H_2SO_4 using phenolphthalein as indicator. In presence of methyl orange, 30 mL of same solution required 40 mL of 0.1 N H_2SO_4 . Calculate the amount of Na_2CO_3 per litre in mixture. **(JEE MAIN)**

Sol: Use titration principles to understand the numerical. Find out the milliequivalents of H_2SO_4 and Na_2CO_3 .

Phenolphthalein as indicator:

Meq. of H_2SO_4 used = $12 \times 0.1 = 1.2$ for 30 mL mixture

$$\therefore \frac{1}{2} \text{ Meq. of } \text{Na}_2\text{CO}_3 \text{ in 30 mL mixture} = 1.2 \quad \dots(i)$$

Methyl orange as indicator: This time fresh solution is titrated with H_2SO_4 using methyl orange as indicator. By equating the data of the bases with the required acid, solve the milliequivalents and then calculate the strength of the bases.

$$\begin{aligned} \text{Meq. of } \text{Na}_2\text{CO}_3 + \text{Meq. of } \text{NaHCO}_3 &= \text{Meq. of } \text{H}_2\text{SO}_4 \text{ used} \\ &= 40 \times 0.1 = 4 \quad \dots(ii) \end{aligned}$$

By Eq. (i)

Meq. of $\text{Na}_2\text{CO}_3 = 2.4$

$$\therefore \frac{w}{53} \times 1000 = 2.4$$

or $w_{\text{Na}_2\text{CO}_3} = 0.1272 \text{ g in } 30 \text{ mL}$

\therefore Strength of $\text{Na}_2\text{CO}_3 = 4.24 \text{ g litre}^{-1}$

Also, Meq. of $\text{NaHCO}_3 = 4 - 2.4 = 1.6$; $\frac{w}{84} \times 1000 = 1.6$

$\therefore w_{\text{NaHCO}_3} = 0.1344 \text{ g in } 30 \text{ mL}$

Strength of $\text{NaHCO}_3 = \frac{0.1344 \times 1000}{30} = 4.48 \text{ g litre}^{-1}$



Illustration 14: 0.5 g mixture of $\text{K}_2\text{Cr}_2\text{O}_7$ and KMnO_4 was treated with excess of KI in acidic medium. Iodine liberated required 150 cm^3 of 0.10N solution of thiosulphate solution for titration.

Find the percentage of $\text{K}_2\text{Cr}_2\text{O}_7$ in the mixture.

(JEE MAIN)

Solution: Determine the equivalent weight of chromate and permanganate solution and compare the mili. Eq of each components to determine the % of chromate.

Reactions of $\text{K}_2\text{Cr}_2\text{O}_7$ and KMnO_4 with KI may be given as :



Thus equivalent wt. of $\text{K}_2\text{Cr}_2\text{O}_7 = \frac{294}{6} = 49$

Equivalent weight of $\text{KMnO}_4 = \frac{158}{5} = 31.6$

m.eq. of $\text{K}_2\text{Cr}_2\text{O}_7 + \text{m.eq. of } \text{KMnO}_4 = \text{m.eq. of } \text{I}_2 = \text{m.eq. of hypo.}$

Let the mass of $\text{K}_2\text{Cr}_2\text{O}_7 = x \text{ g}$

Mass of $\text{KMnO}_4 = (0.5-x)\text{g}$

$$\frac{x}{49} + \frac{(0.5-x)}{31.6} = 150 \times 0.1 \times 10^{-3} \times 2 = 0.03$$

$$\% \text{ of } \text{K}_2\text{Cr}_2\text{O}_7 = \frac{0.0732}{0.5} \times 100 = 14.64$$

4.3 Double Indicators Titration

For the titration of alkali mixtures (e.g., $\text{NaOH} + \text{Na}_2\text{CO}_3$) or $(\text{Na}_2\text{CO}_3 + \text{NaHCO}_3)$, two indicators phenolphthalein and methyl orange are used. This will be discussed in detail in Ionic Equilibrium.

4.4 Eudiometry

Eudiometry or gas analysis involves the calculation based on gaseous reactions in which the amounts of gases are represented by their volumes, measured at STP. Some basic assumptions for calculations

(a) Gay-Lussac's law of volume combination holds good.

(b) For non-reacting mixture. Amagat's law holds good. According to this, the total volume of a non-reacting gaseous mixture is equal to the sum of partial volumes of all the component gases. The volume of solids or liquids is considered to be negligible in comparison to the volumes of gases.

Thus, we can summarize the above points as – eudiometry involves volume measurement during the reaction. Since, Volume of gas, V is directly proportional to number of moles at constant P, T and thus, volume ratio of gases can be directly used in place of mole ratio for analysis.

Illustration 15: A mixture of ethane (C_2H_6) and ethene (C_2H_4) occupies 40 litre at 1.00 atm and at 400 K. The mixture reacts completely with 130 g of O_2 to produce CO_2 and H_2O . Assuming ideal gas behaviour, calculate the mole fraction of C_2H_4 and C_2H_6 in the mixture. **(JEE MAIN)**

Sol: Using the Ideal gas equation, find out no. of moles. Frame the balanced combustion reactions of the hydrocarbons and lay down the values. Calculate the mole fraction accordingly.

For a gaseous mixture of C_2H_6 and C_2H_4



$$PV = nRT$$

$$\therefore 1 \times 40 = n \times 0.082 \times 400$$

$$\text{Total mole of } (C_2H_6 + C_2H_4) = 1.2195$$

Let mole of C_2H_6 and C_2H_4 be a and b respectively.

$$a + b = 1.2195 \quad \dots(i)$$



$$\therefore \text{Mole of } O_2 \text{ needed for complete reaction of mixture} = 7a/2 + 3b$$

$$\therefore \frac{7a}{2} + 3b = \frac{130}{32} \quad \dots(ii)$$

By Eqs. (i) and (ii), $a = 0.808$

$$b = 0.4115$$

$$\therefore \text{Mole fraction of } C_2H_6 = \frac{0.808}{1.2195} = 0.66 \text{ and Mole fraction of } C_2H_4 = 0.34$$

4.5 To Represent Concentration of H_2O_2 Solution

(a) In percentage: The amount of H_2O_2 present in 100 mL H_2O_2 solution is H_2O_2 concentration in percentage of H_2O_2 solution.

(b) In volume: The volume of O_2 at STP given by 1 mL H_2O_2 solution on decomposition is H_2O_2 concentration of H_2O_2 in volume.

Note:

(i) Direct conversions can be made by using following relations

- % strength = $\frac{17}{56} \times \text{volume strength}$
- Volume strength = $5.6 \times \text{Normality}$
- Volume strength = $11.2 \times \text{Molarity}$

(ii) The volume strength of H_2O_2 solution decreases on long standing due to decomposition of H_2O_2 and O_2 .

Illustration 16: Report the concentration of 1.5 N solution of H_2O_2 in terms of volume. **(JEE MAIN)**

Sol: From the given equivalent of H_2O_2 , calculate the weight and then the volume of O_2 . This itself can solve the volume strength of H_2O_2 .

$$\therefore \text{Equivalent of } H_2O_2 \text{ in 1 litre solution} = 1.5$$

$$\therefore w_{H_2O_2} \text{ in 1 litre solution} = 1.5 \times \frac{34}{2} = 25.5 \text{ g}$$

$$\therefore \text{Volume of O}_2 \text{ obtained by 1000 mL H}_2\text{O}_2 \text{ solution} = \frac{22400 \times 25.5}{68} = 8400 \text{ mL}$$

$$\therefore \text{Volume strength of H}_2\text{O}_2 = \frac{8400}{1000} = 8.4$$

4.6 To Represent the Concentration of Oleum

(100 – X%) of oleum means 'X' g H₂O reacts with equivalent amount of free SO₃ to give H₂SO₄.

Illustration 17: 0.5 g of fuming H₂SO₄ (oleum) is diluted with water. The solution requires 26.7 mL of 0.4N NaOH for complete neutralization. Find the % of free SO₃ in the sample of oleum. Also report % of oleum solution.

(JEE ADVANCED)

Sol: Principle of titration is used. Equation of oleum and the base in terms of their milliequivalents is done wherein the amount of oleum is found. % of SO₃ is thus found. Reaction of SO₃ with H₂O gives H₂SO₄. Lay down the calculated and the given values and solve the % of oleum.


Fuming H₂SO₄ contains H₂SO₄ and SO₃. Both react with NaOH. Let a g and b g SO₃ be present.

$$\text{For reaction, } \therefore \text{Meq. of H}_2\text{SO}_4 + \text{Meq. of SO}_3 = \text{Meq. of NaOH}; \frac{a}{98/2} \times 1000 + \frac{b}{80/2} \times 1000 = 26.7 \times 0.4$$

$$\therefore 80a + 98b = 41.87 \quad \dots(i)$$

$$\text{Also, } a + b = 0.5 \quad \dots(ii)$$

$$\therefore \% \text{ of SO}_3 = \frac{0.1039}{0.5} \times 100 = 20.78\%$$



$$80 \text{ g SO}_3 = 18 \text{ g H}_2\text{O}$$

$$\therefore 20.78 \text{ g SO}_3 = \frac{18 \times 20.78}{80} = 4.68$$

$$\% \text{ of oleum} = 100 + 4.68 = 104.68\%$$

4.7 To Determine Hardness of Water

Water, which gives foams easily with soap is called soft water and if not then hard water. The hardness of water is due to the presence of bicarbonates, chlorides and sulphates of Ca and Mg. The extent of hardness is known as **degree of hardness** defined usually as the no. of parts by weight of CaCO₃ present per million parts by weight of water. Hardness is expressed in ppm i.e., 1 ppm = 1 part of CaCO₃ in 10⁶ part of hard water.

Note: The reason for choosing CaCO₃ as the standard to express hardness, inspite of the fact that CaCO₃ is not soluble in water but its molecular weight is 100 which makes calculation easy.

4.8 Mass Balance Equations

The principle of mass balance is based on the law of conservation of mass, i.e., the number of atoms of an element remains constant in a chemical reaction.

4.9 Charge Balance Equations

The principle of charge balance equations is based on the principle of electroneutrality, i.e., all solution are electrically neutral since sum of positive charges equals the sum of negative charges.

4.10 Saponification Value

It is the amount of KOH in mg required to neutralize a fatty acid obtained by the hydrolysis of 1 g of oil.

REDOX REACTIONS

1. INTRODUCTION

Molecular Equations: $2\text{FeCl}_3 + \text{SnCl}_2 \rightarrow 2\text{FeCl}_2 + \text{SnCl}_4$

The reactants and products have been written in molecular forms; thus, the equation is termed as **molecular equation**.


Ionic Equations: The reactions in which the reactants and products are present in the form of ions are called **ionic reactions**.

For example: $2\text{Fe}^{3+} + 6\text{Cl}^- + \text{Sn}^{2+} + 2\text{Cl}^- \rightarrow 2\text{Fe}^{2+} + 4\text{Cl}^- + \text{Sn}^{4+} + 4\text{Cl}^-$

Or $2\text{Fe}^{3+} + \text{Sn}^{2+} \rightarrow 2\text{Fe}^{2+} + \text{Sn}^{4+}$

Illustration 18: Represent the following equation in ionic form.

(JEE MAIN)



Sol: Knowing the oxidation numbers of the elements present, balanced ionic form can be represented. In this equation except H_2O , all are ionic in nature. Representing these compounds in ionic forms,



2K^+ ions and 13SO_4^{2-} ions are common on both sides, so these are cancelled. The desired ionic equation reduces to, $\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 6\text{Fe}^{2+} \rightarrow 6\text{Fe}^{3+} + 2\text{Cr}^{3+} + 7\text{H}_2\text{O}$

Phenomenon of Oxidation and Reduction:

Oxidation or de-electronation is a process which liberates electrons.

Reduction or electronation is a process which gains electrons.

Oxidation	Reduction
a. $\text{M} \longrightarrow \text{M}^{n+} + n\text{e}^-$	$\text{M}^{n+} + n\text{e}^- \longrightarrow \text{M}$
b. $\text{M}^{n_1+} \longrightarrow \text{M}^{n_2+} + (n_2 - n_1)\text{e}^- \quad (n_2 > n_1)$	$\text{M}^{n_2+} + (n_2 - n_1)\text{e}^- \longrightarrow \text{M}^{n_1+} \quad (n_2 > n_1)$
c. $\text{A}^{n-} \longrightarrow \text{A} + n\text{e}^-$	$\text{A} + n\text{e}^- \longrightarrow \text{A}^{n-}$
d. $\text{A}^{n_1-} \longrightarrow \text{A}^{n_2-} + (n_1 - n_2)\text{e}^-$	$\text{A}^{n_2-} + (n_1 - n_2)\text{e}^- \longrightarrow \text{A}^{n_1-}$

Note: M may be an atom or a group of atoms; A may be atom or a group of atoms.

Oxidizing and Reducing Agent:

(a) If an element is in its highest possible oxidation state in a compound, it can function as an oxidizing agent, e.g. KMnO_4 , $\text{K}_2\text{Cr}_2\text{O}_7$, HNO_3 , H_2SO_4 , HClO_4 etc.

(b) If an element is in its lowest possible oxidation state in a compound, it can function as a reducing agent, e.g. H_2S , FeSO_4 , $\text{Na}_2\text{S}_2\text{O}_3$, SnCl_2 etc.

- (c) If an element is in its intermediate oxidation state in a compound, it can function both as an oxidizing agent as well as reducing agent, e.g. H_2O_2 , H_2SO_3 , HNO_3 , SO_2 etc.
- (d) If highly electronegative element is in its higher oxidation state in a compound, that compound can function as a powerful oxidizing agent, e.g. KClO_4 , KClO_3 , KIO_3 etc.
- (e) If an electronegative element is in its lowest possible oxidation state in a compound or in free state, it can function as a powerful reducing agent, e.g. I^- , Br^- , N_3^- etc.

2. MODERN CONCEPT OF OXIDATION AND REDUCTION

According to the modern concept, loss of electrons is oxidation whereas gain of electrons is reduction. Oxidation and reduction can be represented in a general way as shown below:




Figure 1.1: Oxidation and Reduction

PLANCESS CONCEPTS

- In a redox process the valency of the involved species changes. The valency of a reducing agent increases while the valency of an oxidising agent decreases in a redox reaction. The valency of a free element is taken as zero.
- Redox reaction involves two half reactions, one involving loss of electron or electrons (oxidation) and the other involving gain of electron or electrons (reduction).

Saurabh Gupta (JEE 2010, AIR 443)


3. ION ELECTRON METHOD FOR BALANCING REDOX REACTIONS

This method involves the following steps:

- (a) Divide the complete equations into two half reactions
 - (i) One representing oxidation
 - (ii) The other representing reduction
- (b) Balance the atoms in each half reaction separately according to the following steps
 - (i) Balance all atoms other than oxygen and hydrogen
 - (ii) To balance oxygen and hydrogen
- (c) **Acidic Medium**
 - (i) Add H_2O to the side which is oxygen deficient to balance oxygen atoms
 - (ii) Add H^+ to the side which is hydrogen deficient to balance H atoms

(d) Basic Medium


- (i) Add OH^- to the side which has less negative charge
- (ii) Add H_2O to the side which is oxygen deficient to balance oxygen atoms
- (iii) Add H^+ to the side which is hydrogen deficient





(JEE MAIN)

Sol:


Step 1: Select the oxidant, reductant atoms and write their half reactions, one representing oxidation and other reduction. i.e., $\text{C}^{+3} \longrightarrow 2\text{C}^{+4} + 2\text{e}^-$



Step 2: Balance the no. of electrons and add the two equation.



Step 3: Write complete molecule of the reductant and oxidant from which respective redox atoms were obtained.




Step 4: Balance other atoms if any (except H and O).

In above example K is unbalanced, therefore,



Step 5: Balance O atom using H_2O on desired side.



4. OXIDATION STATE AND OXIDATION NUMBER

4.1 Oxidation State

It is defined as the charge (real or imaginary) which an atom appears to have when it is in combination. In the case of electrovalent compounds, the oxidation number of an element or radical is the same as the charge on the ion.

4.2 Oxidation Number

- (a) Oxidation number of an element in a particular compound represents the number of electrons lost or gained by an element during its change from free state into that compound or Oxidation number of an element in a particular compound represent the extent of oxidation or reduction of an element during its change from free state into that compound.
- (b) Oxidation number is given positive sign if electrons are lost. Oxidation number is given negative sign if electrons are gained.
- (c) Oxidation number represent real change in case of ionic compounds. However, in covalent compounds it represents imaginary charge.

Rules for Calculation of Oxidation Number:

Following rules have been arbitrarily adopted to decide oxidation number of elements on the basis of their periodic properties.

- (a) In uncombined state or free state, oxidation number of an element is zero.
- (b) In combined state oxidation number of-
 - (i) F is always -1 .
 - (ii) O is -2 . In peroxide it is -1 , in superoxides it is $-1/2$. However in F_2O it is $+2$.
 - (iii) H is $+1$. In ionic hydrides it is -1 . (i.e., IA, IIA and IIIA metals).
 - (iv) Halogens as halide is always -1 .
 - (v) Sulphur as sulphide is always -2 .
 - (vi) Metal is always $+ve$.
 - (vii) Alkali metals (i.e., IA group – Li, Na, K, Rb, Cs, Fr) is always $+1$.
 - (viii) Alkaline earth metals (i.e., IIA group – Be, Mg, Ca, Sr, Ba, Ra) is always $+2$.
- (c) The algebraic sum of the oxidation number of all the atoms in a compound is equal to zero. e.g. $KMnO_4$.

$$\text{Ox. no. of K} + \text{Ox. no. of Mn} + (\text{Ox. no. of O}) \times 4 = 0$$

$$(+1) + (+7) + 4 \times (-2) = 0$$
- (d) The algebraic sum of all the oxidation no. of elements in a radical is equal to the net charge on the radical. e.g. CO_3^{2-} .

$$\text{Oxidation no. of C} + 3 \times (\text{Oxidation no. of O}) = -2(4) + 3 \times (-2) = -2$$
- (e) Oxidation number can be zero, $+ve$, $-ve$ (integer or fraction)
- (f) Maximum oxidation no. of an element is = Group no. (Except O and F)
 Minimum oxidation no. of an element is = Group no. -8 (Except metals)
 Redox reactions involve oxidation and reduction both. Oxidation means loss of electrons and reduction means gain of electrons. Thus redox reactions involve electron transfer and the number of electrons lost are same as the number of electrons gained during the reaction. This aspect of redox reaction can serve as the basis of a pattern for balancing redox reactions.

Oxidation number of Mn in $KMnO_4$: Let the oxidation number of Mn be x . Now we know that the oxidation numbers of K is $+1$ and that of O is -2 .



Now to the sum of oxidation numbers of all atoms in the formula of the compound must be zero, i.e. $+1 + x - 8 = 0$.
 Hence, the oxidation number of Mn in $KMnO_4$ is $+7$.

Illustration 20: What is the oxidation number of Cr in $K_2Cr_2O_7$?

(JEE MAIN)

Sol: Let the Ox. no. of Cr in $K_2Cr_2O_7$ be x .

We know that, Ox. no. of K = $+1$

Ox. no. of O = -2

So, $2(\text{Ox. no. K}) + 2(\text{Ox. no. Cr}) + 7(\text{Ox. no. O}) = 0$

$$2(+1) + 2(x) + 7(-2) = 0$$

$$\text{or } +2 + 2x - 14 = 0$$

or $2x = +14 - 2 = +12$

or $x = +\frac{12}{2} = +6$ Hence, oxidation number of Cr in is +6.

Illustration 21: H_2S act only as reductant, whereas SO_2 acts as oxidant and reductant both. **(JEE ADVANCED)**

Sol: Oxidation number of S is -2 in H_2S . It can increase only oxidation number up to $+6$.

Oxidation number of S is $+4$ in SO_2 . It can increase or decrease as it lies between maximum $(+6)$ and minimum (-2) oxidation number of S.

Illustration 22: Which compound amongst the following has the highest oxidation number of Mn? KMnO_4 , K_2MnO_4 , MnO_2 and Mn_2O_3 . **(JEE MAIN)**

Sol:

		Ox. no. of Mn
KMnO_4	$+1+x-8=0 \quad x=+7$	$+7$
K_2MnO_4	$+2+x-8=0 \quad x=+6$	$+6$
MnO_2	$x-4=0 \quad x=+4$	$+4$
Mn_2O_3	$2x-6=0 \quad x=+3$	$+3$

Thus, the highest oxidation number for Mn is in KMnO_4 .

4.3 Balancing of Redox Reactions by Oxidation State Method


This method is based on the fact that the number of electrons gained during reduction must be equal to the number of electrons lost during oxidation. Following steps must be followed while balancing redox equations by this method.

- Write the skeleton equation (if not given, frame it) representing the chemical change.
- With the help of oxidation number of elements, find out which atom is undergoing oxidation/reduction, and write separate equations for the atom undergoing oxidation/reduction.
- Add the respective electrons on the right for oxidation and on the left for reduction equation. Note that the net charge on the left and right side should be equal.
- Multiply the oxidation and reduction reactions by suitable integers so that total electrons lost in one reaction is equal to the total electrons gained by other reaction.
- Transfer the coefficients of the oxidizing and reducing agents and their products as determined in the above step to the concerned molecule or ion.
- By inspection, supply the proper coefficient for the other formulae of substances not undergoing oxidation and reduction to balance the equation.


Illustration 23: $\text{Cr}_2\text{O}_7^{2-} + \text{I}^- + \text{H}^+ \longrightarrow \text{Cr}^{3+} + \text{I}_2$

(JEE MAIN)


Sol: (i) Find the oxidation state of atoms undergoing redox change




(ii) Balance the number of atoms undergoing redox change.



(iii) Find the change in oxidation state and balance the change in oxidation states by multiplying the species with a suitable integer.



As the decrease in oxidation state of chromium is 6 and increase in oxidation state of iodine is 2, so we will have to multiply I^- / I_2 by 3 to equalize the changes in oxidation state.



(iv) Find the total charges on both the sides and also find the difference of charges.

$$\text{Charge on LHS} = -2 + 6 \times (-1) = -8$$

$$\text{Charge on RHS} = 2 \times (+3) = +6$$

$$\text{Difference in charge} = +6 - (-8) = 14$$

(v) Now, as the reaction is taking place in acidic medium, we will have to add the ions, to H^+ the side falling short in positive charges, so we will add 14H^+ and LHS to equalize the charges on both sides.



(vi) To equalize the H and O atoms, add $7\text{H}_2\text{O}$ on RHS






Illustration 24: Balance the following equation by oxidation number method:



(JEE ADVANCED)

Sol: Writing oxidation numbers of all atoms,




Oxidation numbers of Cl and I have changed.




Decrease in Ox. no. of Cl = 2 units per Cl_2 molecule

Increase in Ox. no. of I = 2 units per IO_3^- molecule




To balance oxygen, 2OH^- ions be added on LHS and one H_2O molecule on RHS. Hence, the balanced equation is




5. TYPES OF REACTIONS

The redox reactions are of the following types:

- (a) **Combination reactions:** A compound is formed by chemical combination of two or more elements. The combination of an element or compound with oxygen is called combustion. The combustion and several other combinations which involve change in oxidation state are called redox reactions.




- (b) **Decomposition reactions:** Decomposition is the reverse process of combination, it involves the breakdown of the compound into two or more components. The product of decomposition must contain at least one component in elemental state.




In above example, there is no change in oxidation state of potassium. Thus, it should be noted that the decomposition does not result into change in the oxidation number of each element.

- (c) **Displacement reactions:** The reactions in which an atom or ion in a compound is displaced by another atom or ion are called displacement reactions. The displacement reactions are of 2 types:

- (i) **Metal displacement:** In these reactions, a metal in a compound is replaced by another metal in an uncombined state. It is found that a metal with stronger reducing character can displace the other metal having a weaker reducing character.





- (ii) **Non-metal displacement:** These displacement reactions generally involve redox reactions, where the hydrogen is displaced. Alkali and alkaline earth metals are highly electropositive, they displace hydrogen from cold water.




- (d) **Disproportionation and Oxidation-Reduction:** One and the same substance may act simultaneously as an oxidizing agent with the result that a part of it gets oxidized to a higher state and rest of it is reduced to lower state of oxidation. Such a reaction, in which a substance undergoes simultaneous oxidation and reduction is called disproportionation and the substance is said to **disproportionate**.


The following are some of the examples of disproportionation:




- (e) Oxidation state of chlorine lies between -1 to $+7$; thus out of ClO^- , ClO_2^- , ClO_3^- , ClO_4^- ; ClO_4^- does not undergo disproportionation because in this oxidation state of chlorine is highest, i.e., $+7$. Disproportionation of the other oxoanions are:



FORMULAE SHEET





RULES IN BRIEF

The following are the definitions of 'mole' represented in the form of equations:

(a) Number of moles of molecules = $\frac{\text{Weight in g}}{\text{Molecular weight}}$

(b) Number of moles of atoms = $\frac{\text{Weight in g}}{\text{Atomic weight}}$

(c) Number of moles of gases = $\frac{\text{Volume at NTP}}{\text{Standard molar volume}}$

(Standard molar volume is the volume occupied by 1 mole of any gas at NTP, which is equal to 22.4 litres.)


(d) Number of moles of atoms / molecules / ions / electrons = $\frac{\text{No. of atoms / molecules / ions / electrons}}{\text{Avogadro constant}}$

(e) Number of moles of solute = Molarity \times Volume of solution in litres

Or No. of millimoles = Molarity \times Volume in mL.

$$\frac{\text{Millimoles}}{1000} = \text{moles}$$

(f) For a compound $M_x N_y$, x moles of N = y moles of M




Solved Examples

JEE Main/Boards

Example 1: Calculate the composition of 109% oleum.

Sol: Let the mass of SO_3 in the sample be 'w' g, then the mass of H_2SO_4 would be $(100 - w)$ g. On dilution,



Moles of SO_3 in oleum = $\frac{w}{80}$ = Moles of H_2SO_4 formed after dilution.

$$\therefore \text{Mass of } \text{H}_2\text{SO}_4 \text{ formed on dilution} = \frac{98w}{80}$$

$$\text{Total mass of } \text{H}_2\text{SO}_4 \text{ present in oleum after dilution} = \frac{98w}{80} + (100 - w) = 109; \quad w = 40$$

Thus oleum sample contains 40% SO_3 and 60% H_2SO_4 .

Example 2: 20g of a sample of $\text{Ba}(\text{OH})_2$ is dissolved in 10 mL of 0.5 N HCl sol. The excess of HCl was titrated with 0.2 N NaOH. The volume of NaOH used was 10 cc. Calculate the percentage of $\text{Ba}(\text{OH})_2$ in the sample.

Sol: The titration principle is applied wherein milli-equivalents of the neutralization reactions is calculated.

Solving further, one gets the mass and % of the base.

$$\text{Milli eq. of HCl initially} = 10 \times 0.5 = 5$$

$$\text{Milli eq. of NaOH consumed}$$

$$= \text{Milli eq. of HCl in excess} = 10 \times 0.2 = 2$$

$$\therefore \text{Milli eq. of HCl consumed}$$

$$= \text{Milli eq. of } \text{Ba}(\text{OH})_2 = 5 - 2 = 3$$

$$\therefore \text{Eq. of } \text{Ba}(\text{OH})_2 = \frac{3}{1000} = 3 \times 10^{-3}$$

$$\text{Mass of } \text{Ba}(\text{OH})_2 = 3 \times 10^{-3} (171/2) = 0.2565 \text{ g}$$

$$\% \text{Ba}(\text{OH})_2 = (0.2565/20) \times 100 = 1.28\%$$

Example 3: One litre of mixture of CO and CO_2 is passed through red hot charcoal in tube. The new volume becomes 1.4 litre. Find out % composition of original mixture by volume. All measurements are made at same P and T.

Sol: Assuming the mixture contents as a and b, the reaction is framed and values are laid down.


Let the mixture contains

$$\text{CO} = a \text{ litre}; \quad \text{CO}_2 = b \text{ litre}$$

$$\therefore a + b = 1$$

...(i)

On passing the mixture over charcoal only CO_2 reacts as:



Vol. before reaction b 0

Vol. after reaction 0 2b

$$\therefore a + 2b = 1.4$$


By Eqs. (i) and (ii)

$$a = 0.6 \text{ litre} \quad \text{or} \quad a = 60\%$$



$$b = 0.4 \text{ litre} \quad \text{or} \quad b = 40\%$$

Example 4: 0.5 g sample containing is treated with HCl liberating Cl_2 . The is passed into a Sol. of KI and 30.0 cm³ of 0.1 M are required to titrate the liberated iodine. Calculate the percentage of in sample. (At. wt. of Mn = 55).

Sol: Principle of titration is involved in which equating the neutralization reactions is done and milliequivalents of each species is calculated. Thus, weight is calculated and the purity is found.



Redox change are: $2\text{e}^- + \text{I}_2 \longrightarrow 2\text{I}^-$



The reactions suggest that,

Meq. of MnO_2 = Meq. of Cl_2 formed

= Meq. of I_2 liberated = Meq. of $\text{Na}_2\text{S}_2\text{O}_3$ used

$$\therefore \frac{w}{M/2} \times 1000 = 0.1 \times 1 \times 30$$

[$\because N_{\text{Na}_2\text{S}_2\text{O}_3} = M_{\text{Na}_2\text{S}_2\text{O}_3}$ since valency factor = 1, see redox changes for $\text{Na}_2\text{S}_2\text{O}_3$]

$$\text{Or } w_{\text{MnO}_2} = \frac{0.1 \times 1 \times 30 \times M}{2000} = \frac{0.1 \times 1 \times 30 \times 87}{2000}$$

$$(\because M_{\text{MnO}_2} = 87); w_{\text{MnO}_2} = 0.1305$$

$$\therefore \text{Purity of } \text{MnO}_2 = \frac{0.1305}{0.5} \times 100 = 26.1\%$$

Example 5: 10 mL mixture of CH_4 , C_2H_4 and C_3H_8 in the ratio 1: 1.5: 2.5 respectively is burnt in excess of air. Calculate the volume of air used and volume of CO_2 formed after combustion. All measurements are made at same P and T.




Sol: Using the given ratios, find the volumes of the hydrocarbons and frame the balanced combustion reactions.

The calculated O_2 level is 1/5th of the air. Hence volume of CO_2 is found.

$$\text{Volume of } \text{CH}_4 = \frac{1 \times 10}{5} = 2 \text{ mL}$$

$$\text{Volume of } \text{C}_2\text{H}_4 = \frac{1.5 \times 10}{5} = 3 \text{ mL}$$

$$\text{Volume of } \text{C}_3\text{H}_8 = \frac{2.5 \times 10}{5} = 5 \text{ mL}$$



\therefore Volume of O_2 needed

$$= 2 \times 2 + 2 \times 3 + 3 \times 5 = 38 \text{ mL}$$






Since, O_2 is 1/5th part of air

$$\therefore V_{\text{air}} = \frac{25 \times 100}{20} = 125 \text{ mL}$$

Volume of CO_2 formed



$$= 2 \times 1 + 2 \times 3 + 3 \times 5 = 23 \text{ mL}$$

Example 6: Select the species acting as reductant and oxidant in the reaction given below:





Sol: Calculate the oxidation numbers, find the loss/gain of electrons and thus identify the respective oxidants and reductants.

In a conjugate pair oxidant has higher oxidation number.





\therefore PCl_3 is reductant and Cl_2 is oxidant.

\therefore In a conjugate pair of redox, the one having higher ox. no. is oxidant.



Oxidant is AlCl_3 and reductant is K.




∴ SO₂ is oxidant and H₂S is reductant.

(iv) No change in ox. no. of either of the conjugate pair.


∴ None is oxidant or reductant.

(v) For I₂ : I₂⁰ → 2I⁺³ and I₂⁰ + 2e⁻ → 2I⁻¹ I₂ acts as oxidant and reductant both.



Example 7: Balance the following reaction





Sol: Here NO₃⁻ is undergoing reduction and Al is undergoing oxidation.




by balancing each half reaction, we get



by multiplying equation (iii) by 3 and equation (iv) by 8, we get



by combining these equations, we get



Example 8: The composition of a sample of wurtzite is Fe_{0.93}O_{1.00}. What percentage of iron is present in the form of Fe III?

Sol: Oxidation no. of Fe in wurtzite is $\frac{200}{93} = 2.15$

It is an intermediate value between two oxidation state of Fe as, Fe (II) and (III).


Let percentage of Fe (III) be a, then

$$2 \times (100 - 0) + 3 \times a = 2.15 \times 100 \text{ Or } a = 15$$

∴ Percentage of Fe(III) = 15%


Example 9: A 5.0 cm³ solution of H₂O₂ liberates 0.508g of iodine from an acidified KI solution. Calculate the strength of H₂O₂ solution in term of volume strength at STP.

Sol: Volume strength is the volume of oxygen released from 1 mL of hydrogen peroxide solution.



$$\frac{w}{17} \times 1000 = \left[\frac{0.508}{\frac{254}{2}} \right] \times 1000$$

$$\therefore w = 0.068 \text{ g}$$





$$\therefore 34 \text{ g H}_2\text{O}_2 \text{ gives } 11.2 \text{ litre O}_2$$

$$\therefore 0.068 \text{ g gives } \frac{11.2 \times 0.068}{34} = 0.0224 \text{ litre} = 22.4 \text{ ml O}_2$$

$$\therefore \text{Volume strength of H}_2\text{O}_2 = \frac{22.4}{5} = 4.48\%$$

Example 10: A 1.100 g sample of copper ore is dissolved and the Cu²⁺ is treated with excess KI. The liberated I₂ requires 12.12 mL of 0.10 M Na₂S₂O₃ solution for titration. What is % copper by mass in the ore?

Sol: The titration reaction is framed to identify the loss/gain of electrons. The milliequivalents of the respective ions are equated and the amount is calculated. % can be found by dividing the whole weight.



$$\text{Meq. of Cu}^{2+} = \text{Meq. of liberated I}_2 = \text{Meq. of Na}_2\text{S}_2\text{O}_3 \\ = 12.12 \times 0.1 \times 1 = 1.212$$



$$\therefore \frac{w_{\text{Cu}^{2+}}}{63.6/1} \times 1000 = 1.212$$

$$\therefore w_{\text{Cu}^{2+}} = 0.077 \text{ g} = w_{\text{Cu}} \quad (\text{Cu} \xrightarrow{\text{H}_2\text{SO}_4} \text{CuSO}_4)$$

$$\therefore \% \text{ Cu} = \frac{0.077}{1.10} \times 100 = 7\%$$

JEE Advanced/Boards

Example 1: Chile salt petre, a source of NaNO₃ also contains NaIO₃. The NaIO₃ can be used as source of iodine, produced in the following reactions.



One litre of chile salt petre solution containing 5.80g NaIO₃ is treated with stoichiometric quantity of NaHSO₃. Now an additional amount of same solution is added to reaction mixture to bring about the second reaction.

How many grams of NaHSO_3 are required in step I and what additional volume of chile salt petre must be added in step II to bring in complete conversion of I^- to I_2 ?

Sol: The titration reaction is used to identify the loss/gain of electrons. The milliequivalents of the respective species are equated and the amount is calculated. Stepwise calculation gives the volume of NaIO_3 .

Meq. of NaHSO_3 = Meq. of NaIO_3

$$= N \times V = \frac{5.8}{198/6} \times 1000$$

[Et. wt. of NaI = $M/6$ because $\text{I}^{3+} + 6e \longrightarrow \text{I}^-$]

Meq. of NaHSO_3 = 175.76

$$\therefore W_{\text{NaHSO}_3} = \frac{175.76 \times 104}{2000} = 9.14 \text{ g}$$

Also Meq. of formed in I step using valence factor 6 = 175.76

In II step valence factor of IO_3^- is 1 and valence factor of is 5.

Thus, Meq. of formed using valence factor 1 = $\frac{175.76}{6}$

Also Meq. of NaIO_3 used in step II = $\frac{175.76}{6}$

$$\therefore N \times V = \frac{175.76}{6}; \Rightarrow \frac{5.8}{198/5} \times V = \frac{175.76}{6}$$

$$\therefore V_{\text{NaIO}_3} = 200 \text{ mL}$$

Example 2: What amount of substance containing 60% NaCl , 37% KCl should be weighed out for analysis so that after the action of 25 mL of 0.1 N AgNO_3 solution, excess of Ag^+ is back titrated with 5 mL of NH_4SCN . Given that 1 mL of NH_4SCN = 1.1 mL of AgNO_3 .

Sol: Let a g of the given sample be weighed out for the purpose. The reaction of the chlorides with AgNO_3 will give AgCl . The unreacted Ag^+ ions will get consumed by NH_4SCN to give AgSCN . Proceeding accordingly, equate the milliequivalents and calculate a.

$$\therefore \text{Wt. of NaCl} = \frac{60}{100} \times a = 0.6 \text{ a g}$$

$$\therefore \text{Wt. of KCl} = \frac{37}{100} \times a = 0.37 \text{ a g}$$

Now this mixture reacts with AgNO_3 , the excess of AgNO_3 is back titrated with NH_4SCN . Meq. of AgNO_3 added to mixture

$$= 25 \times 0.1 = 2.5$$

Normality of NH_4SCN can be derived as

Meq. of NH_4SCN = Meq. of AgNO_3

$$N \times 1 = 0.1 \times 1.1$$

$$N = 0.11$$

Meq. of AgNO_3 left = Meq. of NH_4SCN

$$= 5 \times N$$

$$\therefore \text{Meq. of AgNO}_3 \text{ left} = 5 \times 0.11 = 0.55$$

$$\therefore \text{Meq. of AgNO}_3 \text{ used for mixture} = 2.5 - 0.55 = 1.95$$

Meq. of KCl + Meq. of NaCl is mixture

$$= 1.95; \frac{0.73a}{74.5} \times 1000 + \frac{0.6a}{58.5} \times 1000 = 1.95$$

$$\therefore a = 0.128 \text{ g}$$

Example 3: NaOH and Na_2CO_3 are dissolved in 200 mL aqueous solution. In the presence of phenolphthalein indicator, 17.5 mL of 0.1 N HCl are used to titrate this solution. Now methyl orange is added in the same sol. titrated and it requires 2.5 mL of the same HCl . Calculate the normality of NaOH and Na_2CO_3 and their mass present in the solution.

Sol: The titration of a simple acid and a base using an indicator is seen over here. The milliequivalents of the acid is calculated and equated with that of the base. The volume and the mass is thus calculated.

Milli equivalent (a) of HCl used in the presence of phenolphthalein indicator.

$$= N \times V \text{ (mL)} = 0.1 \times 17.5 = 1.75$$

$$1.75 \text{ (a)} = \text{milli. eq. of NaOH} + \frac{1}{2} \text{ milli eq. of Na}_2\text{CO}_3 \quad \dots \text{ (i)}$$

Milli eq. (b) of HCl used in the presence of methyl orange indicator

$$= N \times V \text{ (mL)} = 0.1 \times 2.5 = 0.25$$

$$0.25 \text{ (b)} = \frac{1}{2} \text{ milli eq. of Na}_2\text{CO}_3 \quad \dots \text{ (ii)}$$

For Na_2CO_3 solution; from equation (ii)

$$\text{Milli eq. of acid used by Na}_2\text{CO}_3 = 2b = 2 \times 0.25 = 0.5$$

Volume of Na_2CO_3 solution = 200 mL

Suppose, Normality of Na_2CO_3 = N

$$\text{Milli equivalents of} = N \times V \text{ (mL)} = 200 N$$

$$\text{Putting equivalents of acid and Na}_2\text{CO}_3 \text{ equal } 200 N = 0.5$$

$$\text{Or (Normality of Na}_2\text{CO}_3 \text{ solution) } N = \frac{1}{400}$$

$$\text{Mass of Na}_2\text{CO}_3 = N \times E \times V \text{ (litre)}$$

$$(E \text{ for Na}_2\text{CO}_3 = 53) = \frac{1}{400} \times 5 \times 0.2 = 0.0265 \text{ gram}$$

For NaOH Sol.; from equation (i) and (ii)

Milli eq. acid used by NaOH = $a - b = 1.75 - 0.25 = 1.50$

Volume of NaOH solution = 200 mL

Suppose, Normality of NaOH solution = N

Milli eq. of NaOH = $N \times V \text{ (mL)} = 200 N$

Putting the milli eq. of NaOH and acid used equal 200
 $N = 1.5$

(Normality of NaOH Sol.) $N = \frac{1.5}{200}$

Mass of NaOH = $N \times E \times (V \text{ litres})$

$= \frac{1.5}{200} \times 40 \times 0.2$ (E for NaOH = 40) = 0.06 g

Example 4: The molarity and molality of a solution are M and m respectively. If the molecular weight of the solute is M' . Calculate the density of the solution in terms of M, m and M' .

Sol: Let weight of solute be w g and weight of solvent be W g, volume of solution be V mL and density be D. Substitute as follows.

$$\therefore M = \frac{w \times 1000}{M' \times V} \quad \dots(i)$$

$$m = \frac{w \times 1000}{M' \times W} \quad \dots(ii)$$

$$D = \frac{w + W}{V} \quad \dots(iii)$$

By Eq. (i) $w = \frac{MM'V}{1000} \quad \dots(iv)$


By Eq. (ii) $W = \frac{w \times 1000}{M' \times m}$

By Eq. (iv) $W = \frac{MM'V \times 1000}{1000 \times M' \times m} = \frac{MV}{m} \quad \dots(v)$

$$\therefore \text{By Eq. (iii)} D = \frac{\frac{MM'V}{1000} + \frac{MV}{m}}{V} \therefore D = M \left[\frac{1}{m} + \frac{M'}{1000} \right]$$


Example 5: 1.249 g of a sample of pure BaCO_3 and impure CaCO_3 containing some CaO was treated with dil. HCl and it evolved 168 mL of CO_2 at NTP. From this solution BaCrO_3 was precipitated, filtered and washed. The precipitate was dissolved in dilute sulphuric acid and diluted to 100 mL. 10 mL of this solution when treated with KI solution, liberated iodine which required exactly 20 mL of 0.05 N $\text{Na}_2\text{S}_2\text{O}_3$. Calculate the percentage of CaO in the sample.

Sol: An acid-base titration accompanied with iodine titration gives the following equation.



Calculating the equivalents of the involved species gives their amount and the %.

$$= \frac{168}{22400} = 7.5 \times 10^{-3} \quad \dots(i)$$



Eq. of $\text{Na}_2\text{S}_2\text{O}_3$ = Eq. of I_2 = Eq. of BaCr_2O_7

$$= \frac{20 \times 10^{-3} \times 0.05 \times 100}{10} = 1 \times 10^{-2}$$

$$\text{Moles of BaCr}_2\text{O}_7 = \frac{1}{6} \times 10^{-2}$$

$$\text{Moles of BaCrO}_4 = \frac{2}{6} (1 \times 10^{-2})$$

$$\text{Moles of BaCO}_3 = \frac{1}{3} \times 10^{-2} = 3.33 \times 10^{-3} \quad \dots(ii)$$

Weight of BaCO_3 = 0.650 gm

From equation (i) and (ii) we get $\Rightarrow n_{\text{CaCO}_3} = 4.17 \times 10^{-3}$



Weight of CaCO_3 = $100 \times 4.17 \times 10^{-3} = 0.417$ g

Weight of CaO = $1.249 - 0.656 - 0.417 = 0.176$

$$\% \text{ of CaO} = \frac{0.176}{1.249} \times 100 = 14.09 \%$$

Example 6: Find out the percentage of oxalate ion in a given sample of oxalate salt of which 0.3 g dissolved in 100 mL of water required 90 mL of N/20 KMnO_4 for complete oxidation.

Sol: Redox changes are



\therefore Meq. of oxalate ion = Meq. of KMnO_4

$$\frac{w}{E} \times 1000 = 90 \times \frac{1}{20}; E_{\text{C}_2\text{O}_4^{2-}} = \frac{\text{Ionic wt.}}{2} \therefore \frac{w}{88} \times 1000 = \frac{9}{2}$$

$$\therefore w_{\text{C}_2\text{O}_4^{2-}} = 0.198 \text{ g}$$

\therefore 0.3 g $\text{C}_2\text{O}_4^{2-}$ sample has oxalate ion = 0.198 g

$$\therefore \text{Percentage of C}_2\text{O}_4^{2-} \text{ in sample} = \frac{0.198 \times 100}{0.3} = 66\%$$

Example 7: Balance the following redox equation, $\text{AsO}_3^{3-} + \text{MnO}_4^- \longrightarrow \text{AsO}_4^{3-} + \text{MnO}_2$ using ion-electron method (alkaline medium)

Sol: (i) Identify the oxidation and reduction halves.


Reduction half reaction: $\text{MnO}_4^- \longrightarrow \text{MnO}_2$

Oxidation half reaction: $\text{AsO}_3^{3-} \longrightarrow \text{AsO}_4^{3-}$


(ii) Atoms of the element undergoing oxidation and reduction are already balanced.

(iii) Balancing O atoms,

Reduction half reactions:



Oxidation half reactions:




(iv) Balancing H atoms, H atoms are already balanced in both the half reactions.

(v) Balancing charge,




Reduction half reaction:




Oxidation half reaction:



(vi) Multiply equation (i) by 3 and equation (ii) by 2 and then add (i) and (ii).




Example 8: 1 g sample of AgNO_3 is dissolved in 50 mL of water. It is titrated with 50 mL of KI solution. The AgI precipitated is filtered off. Excess of KI in filtrate is titrated with $\text{M}/10 \text{ KIO}_3$ in presence of 6M HCl till all I^- converted into ICl . It requires 50 mL of $\text{M}/10 \text{ KIO}_3$ solution. 20 mL of the same stock solution of KI requires 30 mL of $\text{M}/10 \text{ KIO}_3$ under similar conditions. Calculate % of AgNO_3 in sample. The reaction is:



Sol: Follow the reaction $\text{AgNO}_3 + \text{KI} \longrightarrow \text{AgI} + \text{KNO}_3$

1. Ag present in AgNO_3 is removed as AgI by adding 50 mL KI of which 20 mL requires 30 mL of $\text{M}/10 \text{ KIO}_3$.
2. The solution contains KI unused. The unused KI is converted into ICl by KIO_3 .

\therefore Meq. of KI in 20 mL = Meq. of KIO_3



\therefore Meq. of KI in 50 mL added to AgNO_3

$$\therefore \text{Eq. wt. of KI} = \frac{\text{M}}{2} = \frac{30 \times 4 \times 50}{10 \times 20} = 30$$

Now, Meq. of KI left unused by $\text{AgNO}_3 = 30 - 20$

\therefore Mole ratio of AgNO_3 and KI

\therefore Meq. of $\text{AgNO}_3 = 10$

Reaction is 1: 1 and thus if Eq.

$$\therefore \frac{w}{170/2} \times 1000 = 10 \quad \left| \quad \text{Wt. of KI is } \text{M}/2, \right.$$



$w = 0.85 \text{ g}$ then Eq. wt. of $\text{AgNO}_3 = \text{M}/2$

\therefore Percentage of purity of AgNO_3 in sample

$$= \frac{0.85 \times 100}{1} = 85\%$$

Example 9: Selenium in a 10.0 gm soil sample is distilled as the tetrabromide, which is collected in an aqueous solution, where it is hydrolysed to SeO_3^{2-} . The SeO_3^{2-} is estimated iodometrically, requiring 4.5 mL of standard $\text{Na}_2\text{S}_2\text{O}_3$ solution for the titration. If 1 mL of $\text{Na}_2\text{S}_2\text{O}_3 = 0.049 \text{ mg}$ of $\text{K}_2\text{Cr}_2\text{O}_7$, what is the concentration of Se in the soil in ppm?

Sol: Follow the reaction



$$1\text{mL Na}_2\text{S}_2\text{O}_3 \equiv \frac{0.049 \times 10^{-3} \times 6}{294} \text{ eq. of } \text{K}_2\text{Cr}_2\text{O}_7$$

$$\equiv \frac{0.049 \times 10^{-3} \times 6 \times 10^3}{294} \text{ Meq. of } \text{K}_2\text{Cr}_2\text{O}_7$$

$\therefore 4.5 \text{ mL Na}_2\text{S}_2\text{O}_3$

$$= \frac{0.049 \times 10^{-3} \times 6 \times 10^3 \times 4.5}{294} \text{ Meq. of } \text{K}_2\text{Cr}_2\text{O}_7 = 4.5 \times 10^{-3}$$

Meq. of $\text{K}_2\text{Cr}_2\text{O}_7$ or Meq. of $\text{Na}_2\text{S}_2\text{O}_3$

Meq. of Se = Meq. of SeO_3^{2-} = Meq. of KI = Meq. of I_2 = Meq. of $\text{Na}_2\text{S}_2\text{O}_3$

$$\frac{w_{\text{Se}}}{79} \times 1000 \times 4 = 4.5 \times 10^{-3}$$

$$\therefore w_{\text{Se}} = 8.8875 \times 10^{-5} \text{ g}$$

$$\therefore \text{ppm} = \frac{8.8875 \times 10^{-5} \times 10^6}{10} = 8.8875$$

Q.20 2.5 moles of sulphuryl chloride were dissolved in water to produce sulphuric acid and hydrochloric acid. How many moles of KOH will be required to completely neutralise the solution?

Q.21 100 g of a sample of common salt containing contamination of NH_4Cl and MgCl_2 to the extent of 2% each by mass is dissolved in water. How much volume of 5% by mass of AgNO_3 solution ($d = 1.04 \text{ g cm}^{-3}$) is required to precipitate all chloride ions?

Q.22 A mixture of formic acid and oxalic acid is heated with concentrated H_2SO_4 . The gases produced are collected and on treatment with KOH solution, the volume of the gases decreased by 1.6th. Calculate the molar ratio of the two acids in the original mixture.

Q.23 The mean molecular mass of a mixture of methane (CH_4) and ethene (C_2H_4) in the molar ratio of x: y is found to be 20. What will be the mean molecular mass if the molar ratio of the gases is reversed?

Q.24 1 g sample of KClO_3 was heated under such conditions that a part of it decomposes as $2\text{KClO}_3 \longrightarrow 2\text{KCl} + 3\text{O}_2$ while the remaining part decomposes as $4\text{KClO}_3 \longrightarrow 3\text{KClO}_4 + \text{KCl}$.

If net oxygen obtained is 146.8 mL at STP.

Calculate the mass of KClO_4 in the residue.

Q.25 A mixture of FeO and Fe_3O_4 was heated in air to constant mass and it was found to gain 5% in its mass. Find the composition of the initial mixture.

Q.26 Equal masses of zinc (at. mass 65) and iodine (at. mass 127) were allowed to react till completion of reaction to form ZnI_2 . Which substance is left unreacted and to what fraction of its original mass?




Q.27 Two gram each of P_4 and O_2 are allowed to react till none of the reactant is left. If the products are P_4O_6 and P_4O_{10} . Calculate the mass of each of the product.

Q.28 A piece of aluminium weighing 2.7 g was heated with 100 mL of H_2SO_4 (25% by mass, $d = 1.18 \text{ g cm}^{-3}$). After complete dissolution of metal, the solution is diluted by adding water to 500 mL.

What is the molarity of free H_2SO_4 in resulting solution?












Q.29 Chemical reaction between ferrous oxalate and KMnO_4 has been given in the form of three partial equations. Write the complete balanced equation and thus find out the volume of 0.5 M KMnO_4 required to

completely react with 1.5 mol of FeC_2O_4 .












Redox Reactions






Q.1 Indicate the oxidation number of underlined in each case:




Q.2 Write complete balanced equation for the following in acidic medium by ion-electron method:









Q.3 Write complete balanced equation for the following in basic medium by ion-electron method:





Q.4 Balance the following equations by oxidation method:



Q.5 Define disproportionation? Give one example.

Q.6 Define difference between ion electron method and oxidation method?

Q.7 What is the most essential conditions that must be satisfied in a redox reaction?


Q.8 Does the oxidation number of an element in any molecule or any poly atomic ion represents the actual charge on it?

Q.9 What is redox couple?




Q.10 Calculate the standard e.m.f. of the cells formed by different combinations of the following half cells.



Q.11 Balance the following equations in acidic medium by both oxidation number and ion electron methods & identify the oxidants and the reductants.



Q.12 Write the half reactions for the following redox reactions:



Q.13 Define oxidation & reduction in term of oxidation number.

Q.14 Discuss the following redox reactions?

(i) Combination reactions

(ii) Decomposition reactions

(iii) Displacement reactions

(iv) Disproportionation reaction

Q.15 What is the difference between valence and oxidation number?

Q.16 H_2S acts only as reducing agent while SO_2 can act both as a reducing agent and oxidising agent. Explain.

Q.17 What are half reactions? Explain with examples?

Q.18 Explain the term:

(i) Oxidation

(ii) Reduction

(iii) Oxidizing agent

(iv) Reducing agent

Exercise 2

Mole Concept

Single Correct Choice Type

Q.1 If 'x' gms of an element A reacts with 16 gms of oxygen then the equivalent weight of element A is

(A) $\frac{x}{4}$

(B) $\frac{x}{2}$

(C) x

(D) 2x

Q.2 The mass of CO containing the same amount of oxygen as in 88 gms of CO_2 is

(A) 56 gms

(B) 28 gms

(C) 112 gms

(D) 14 gms

Q.3 When 8 gms of oxygen reacts with magnesium then the amount of MgO formed is

(A) 18 gm

(B) 20 gm

(C) 24 gm

(D) 32 gm

Q.4 One gram of the silver salt of an organic dibasic acid yields, on strong heating 0.5934 g of silver. If the weight percentage of carbon in it 8 times the weight percentage of hydrogen and one half the weight percentage of oxygen, determine the molecular formula of the acid. [Atomic weight of Ag = 108]


(A) $\text{C}_4\text{H}_6\text{O}_4$

(B) $\text{C}_4\text{H}_6\text{O}_6$

(C) $\text{C}_2\text{H}_6\text{O}_2$

(D) $\text{C}_5\text{H}_{10}\text{O}_5$

Q.5 Mass of sucrose $C_{12}H_{22}O_{11}$ produced by mixing 84 gm of carbon, 12 gm of hydrogen and 56 liter O_2 at 1 atm and 273 K according to given reaction, is



- (A) 138.5 (B) 155.5 (C) 172.5 (D) 199.5

Q.6 40 gm of carbonate of an alkali metal or alkaline earth metal containing some inert impurities was made to react with excess HCl solution. The liberated CO_2 occupied 12.315 lit. at 1 atm and 300 K. The correct option is

- (A) Mass of impurity is 1 gm and metal is Be
(B) Mass of impurity is 3 gm and metal is Li
(C) Mass of impurity is 5 gm and metal is Be
(D) Mass of impurity is 2 gm and metal is Mg

Q.7 An hydride of nitrogen decomposes to give nitrogen and hydrogen. It was formed that one volume of the hydride gave one volume of N_2 and 2 volume of H_2 at STP. The hydride of nitrogen is

- (A) NH_3 (B) N_2H_6 (C) NH_2 (D) N_2H_4


Q.8 5 volumes of a hydrocarbon on complete consumed 10 volumes of oxygen giving 5 volumes of CO_2 at STP. The hydrocarbon is

- (A) C_2H_6 (B) C_2H_4 (C) CH_4 (D) C_2H_4

Q.9 The percentage by mole of NO_2 in a mixture of $NO_2(g)$ and $NO(g)$ having average molecular mass 34 is

- (A) 25% (B) 20% (C) 40% (D) 75%

Q.10 The minimum mass of mixture of A_2 and B_4 required to produce at least 1 kg of each product is (Given At. mass of 'A' = 10; At. mass of 'B' = 120)



- (A) 2120 gm (B) 1060 gm (C) 560 gm (D) 1660 gm

Q.11 74 gm of a sample on complete combustion given 132 gm CO_2 and 54 gm of H_2O . The molecular formula of the compound may be

- (A) C_5H_{12} (B) $C_4H_{10}O$ (C) $C_3H_{10}O_2$ (D) $C_3H_7O_2$

Q.12 The volume of oxygen used when x gms of Zn is converted to ZnO is

- (A) $\frac{2x}{65} \times 5.6$ litres (B) $\frac{x}{65} \times 5.6$ litres
(C) $\frac{4x}{65} \times 5.6$ litres (D) None of these

Q.13 A sample of clay was partially dried and then contained 50% silica and 7% water. The original clay contained 12% water. The silica in original sample is

- (A) 51.69 (B) 47.31
(C) 63.31 (D) None of these

Q.14 The mass of CO_2 produced from 620 mixture of $C_2H_4O_2$ and O_2 , prepared produce maximum energy is (combustion reaction is exothermic)

- (A) 413.33 gm (B) 593.04 gm
(C) 440 gm (D) 320 gm

Q.15 In the quantitative determination of nitrogen, N_2 gas liberated from 0.42 gm of a sample of organic compound was collected over water. If the volume of N_2 gas collected was 100/11 mL at total pressure 860 mm Hg at 250 K, % by mass of nitrogen in the organic compound is

[Aq. tension at 250 K is 24 mm Hg and $R = 0.08 \text{ L atm mol}^{-1} \text{ K}^{-1}$]

- (A) $\frac{10}{3}\%$ (B) $\frac{5}{3}\%$ (C) $\frac{20}{3}\%$ (D) $\frac{100}{3}\%$

Q.16 300 mL of 0.1 M HCl and 200 mL of 0.3 M H_2SO_4 are mixed. The normality of the resulting mixture is

- (A) 0.4 N (B) 0.1 N (C) 0.3 N (D) 0.2 N

Q.17 The volume of water which should be added to 300 mL of 0.5 M NaOH solution so as to get a solution of 0.2 M is

- (A) 550 mL (B) 350 mL (C) 750 mL (D) 450 mL

Q.18 The mole fraction of a solution containing 3.0 gms of urea per 250 gms of water would be

- (A) 0.00357 (B) 0.99643
(C) 0.00643 (D) None of these

Q.19 The mass of P_4O_{10} produced if 440 gm of P_4S_3 is mixed with 384 gm of O_2 is $P_4S_3 + O_2 \longrightarrow P_4O_{10} + SO_2$

- (A) 568 gm (B) 426 gm
(C) 284 gm (D) 396 gm

Q.20 Calculate percentage change in M_{avg} of the mixture, if PCl_5 undergo 50% decomposition. $PCl_5 \longrightarrow PCl_3 + Cl_2$

- (A) 50% (B) 66.66%
(C) 33.33% (D) Zero

Q.21 The mass of Mg_3N_2 produced if 48 gm of Mg metal is reacted with 34 gm NH_3 gas is $\text{Mg} + \text{NH}_3 \longrightarrow \text{Mg}_3\text{N}_2 + \text{H}_2$

- (A) $\frac{200}{3}$ (B) $\frac{100}{3}$ (C) $\frac{400}{3}$ (D) $\frac{150}{3}$

Q.22 The molarity of a solution of conc. HCl containing 36.5% by weight of HCl would be

- (A) 16.75 (B) 17.75 (C) 15.75 (D) 14.75

Q.23 0.35 gms of a sample of $\text{Na}_2\text{CO}_3 \cdot x\text{H}_2\text{O}$ were dissolved in water and the volume was made to 50 mL of this solution required 9.9 mL of $\frac{N}{10}$ HCl for complete neutralization. Calculate the value of x.

- (A) 1 (B) 2 (C) 3 (D) None of these

Q.24 1.2 gms of a sample of washing soda was dissolved in water and volume was made upto 250 cc. 25 cc of this solution when titrated against N/10 HCl for required 17 mL. The percentage of carbonate is given sample is

- (A) Approximately 70% (B) Approximately 66%
(C) Approximately 76% (D) None of these

Q.25 The number of carbon atoms present in a signature, if a signature written by carbon pencil weights 1.2×10^{-3} g is

- (A) 12.40×10^{20} (B) 6.02×10^{19}
(C) 3.01×10^{19} (D) 6.02×10^{20}

Q.26 The average atomic mass of a mixture containing 79 mole % of ^{24}Mg is 24.31. % mole of ^{26}Mg is

- (A) 5 (B) 20 (C) 10 (D) 15

Q.27 25 cc of solution containing NaOH and Na_2CO_3 when titrated against N/10 HCl. Using phenolphthalein as indicator required 40 cc. of HCl. The same volume of mixture when titrated against N/10 HCl using methyl orange required 45cc of this HCl. The amount of NaOH and Na_2CO_3 in one mixture is

- (A) NaOH = 28 gm/L ; Na_2CO_3 = 10.6 gm/L
(B) NaOH = 10.6 gm/L ; Na_2CO_3 = 28 gm/L
(C) NaOH = 14 gm/L ; Na_2CO_3 = 5.3 gm/L
(D) None of these

Q.28 0.5 gms of a mixture of K_2CO_3 and Li_2CO_3 requires 30 mL of 0.25 NHCl solution for neutralization. The percentage composition of mixture would be

- (A) K_2CO_3 = 96%; Li_2CO_3 = 4%
(B) K_2CO_3 = 4%; Li_2CO_3 = 96%
(C) K_2CO_3 = 50%; Li_2CO_3 = 25%
(D) K_2CO_3 = 50%; Li_2CO_3 = 74%

Q.29 How many mL of a 0.05 M KMnO_4 solution are required to oxidise 2.0 g of FeSO_4 in a dilute acid solution?

- (A) 5.263 (B) 0.5263
(C) 52.63 (D) None of these

Redox Reaction

Single Correct Choice Type

Q.1 The equivalent weight of FeSO_4 when it is oxidised by acidified KMnO_4 will be equal to

- (A) M_0 of FeSO_4 (B) $\frac{M_0 \text{FeSO}_4}{2}$
(C) $2M_0 \text{FeSO}_4$ (D) $\frac{M_0 \text{FeSO}_4}{4}$

Q.2 The equivalent weight of $\text{K}_2\text{Cr}_2\text{O}_7$ when it is converted Cr^{3+} will be equal to

- (A) $M_{\text{K}_2\text{Cr}_2\text{O}_7}$ (B) $\frac{M_{\text{K}_2\text{Cr}_2\text{O}_7}}{3}$
(C) $\frac{M_{\text{K}_2\text{Cr}_2\text{O}_7}}{4}$ (D) $\frac{M_{\text{K}_2\text{Cr}_2\text{O}_7}}{6}$

Q.3 The amount of H_2S that can be oxidised to sulfur on oxidation using 1.58 gm of KMnO_4 as oxidising agent in acidic medium will be

- (A) 0.85 gms (B) 1.7 gms
(C) 0.425 gms (D) None of these

Q.4 The amount of nitric acid required to oxidise 127 gms of I_2 to I_2O_5 will be _____. Assume that during the reaction HNO_3 gets converted to NO_2 .

- (A) 12.7 (B) 3.15 (C) 315 (D) 31.5

Q.5 10 mL of oxalic acid was completely oxidised by 20 mL of 0.02 M KMnO_4 . The normality of oxalic acid solution is

- (A) 0.05 N (B) 0.1 N (C) 0.2 N (D) 0.025 N

Q.6 0.2 g of a sample of H_2O_2 required 10 mL of 1N KMnO_4 in a titration in the presence of H_2SO_4 . Purity of H_2O_2 is

- (A) 25% (B) 65% (C) 85% (D) None of these

Q.7 The number of moles of KMnO_4 that will be needed to react completely with one mole of ferrous oxalate in acidic solution is

- (A) $\frac{2}{5}$ (B) $\frac{3}{5}$ (C) $\frac{4}{5}$ (D) 1

Q.8 A metal oxide is reduced by heating it in a stream of hydrogen. It is found that after complete reduction, 3.15 g of the oxide has yielded 1.05 g of the metal. We may deduce that

- (A) The atomic weight of the metal is 8
(B) The atomic weight of the metal is 4
(C) The equivalent weight of the metal is 4
(D) The equivalent weight of the metal is 8

Q.9 Oxidation involves

- (A) Gain of electrons
(B) Loss of electrons
(C) Increase in the valency of negative part
(D) Decrease in the valency of positive part

Q.10 The oxidation number of Cr in $\text{K}_2\text{Cr}_2\text{O}_7$

- (A) +2 (B) -2 (C) +6 (D) -6

Q.11 When $\text{K}_2\text{Cr}_2\text{O}_7$ is converted into $\text{K}_2\text{Cr}_2\text{O}_4$ the change in oxidation number of Cr is

- (A) 0 (B) 6 (C) 4 (D) 3

Q.12 White P reacts with caustic soda. The products are PH_3 and NaH_2PO_2 . This reaction is an example of

- (A) Oxidation (B) Reduction
(C) Oxidation and reduction (D) Neutralization

Q.13 The oxidation number of carbon in CH_2O is

- (A) -2 (B) +2 (C) 0 (D) +4

Q.14 The oxidation number of C in CH_4 , CH_3Cl , CH_2Cl_2 , CHCl_3 , and CCl_4 are respectively

- (A) 0, 2, -2, 4, -4 (B) -4, -2, 0, +2, +4
(C) 2, 4, 0, -2, -4 (D) 4, 2, 0, -2, -4

Q.15 Which of the following reactions is not redox type

- (A) $2\text{BaO} + \text{O}_2 \longrightarrow 2\text{BaO}_2$
(B) $4\text{KClO}_3 \longrightarrow 2\text{KClO}_4 + \text{KCl}$
(C) $\text{BaO}_2 + \text{H}_2\text{SO}_4 \longrightarrow \text{BaSO}_4 + \text{H}_2\text{O}_2$
(D) $\text{SO}_2 + 2\text{H}_2\text{S} \longrightarrow 2\text{H}_2\text{O} + 3\text{S}$

Q.16 In which of the following compounds iron has lowest oxidation state

- (A) $\text{K}_4\text{Fe}(\text{CN})_6$ (B) K_2FeO_4
(C) Fe_2O (D) $\text{Fe}(\text{CO})_5$

Q.17 Select the compound in which chlorine is assigned the oxidation number +5

- (A) HClO (B) HClO_2 (C) HClO_3 (D) HClO_4

Q.18 If three electrons are lost by a metal iron M^{3+} its final oxidation number would be

- (A) 0 (B) +2 (C) +5 (D) +6

Q.19 The oxidation number of Mn in MnO_4^- is

- (A) +7 (B) -5 (C) -7 (D) +5

Q.20 The oxidation number of carbon in CHCl_3 is

- (A) +2 (B) +4 (C) +4 (D) -3

Q.21 Pb^{2+} loses two electrons in a reaction. What will be the oxidation number of lead after the reaction?

- (A) +2 (B) 0 (C) +4 (D) -2


Q.22 The oxidation number of carbon in $\text{C}_{12}\text{H}_{22}\text{O}_{11}$ is

- (A) 0 (B) -6 (C) +2 (D) +6

Q.23 The oxidation state of sulphur in SO_4^{2-} is


- (A) +2 (B) +4 (C) +5 (D) +6

Q.24 If the following reaction 'X' is



- (A) 1e^- (B) 2e^- (C) 3e^- (D) 4e^-

Q.25 In the following reaction the value of 'X' is



- (A) 4e^- (B) 3e^- (C) 2e^- (D) 1e^-

Q.26 The oxidation state of sulphur in $S_2O_7^{2-}$ is
(A) +6 (B) -6 (C) -2 (D) +2

Q.27 The oxidation number and covalency of sulphur in S_8 are respectively
(A) 0 & 2 (B) 0 & 8 (C) 6 & 8 (D) 6 & 2

Q.28 The oxidation state of nitrogen in N_3H is
(A) $1/3$ (B) +3 (C) -1 (D) $-1/3$

Q.29 The oxidation number of iron in potassium ferricyanide is
(A) +1 (B) +2 (C) +3 (D) +4

Q.30 Oxidation number of hydrogen in MH_2 is
(A) +1 (B) -1 (C) +2 (D) -2

Q.31 The oxidation state of phosphorus varies from
(A) -1 to +1 (B) -3 to +3 (C) -3 to +5 (D) -5 to +1

Q.32 Select the compound in which chlorine is assigned the oxidation number +5
(A) $HClO_4$ (B) $HClO_2$ (C) $HClO_3$ (D) HCl

(B) 33.6 l $H_2(g)$ is produced regardless of temperature and pressure for every mole Al that reacts
(C) 67.2 l $H_2(g)$ at STP is produced for every mole Al that reacts
(D) 11.2 $H_2(g)$ at STP is produced for every mole HCl (aq) consumed

Q.4 How many moles of magnesium phosphate, $Mg_3(PO_4)_2$ will contain 0.25 mole of oxygen atoms (2006)
(A) 0.02 (B) 3.125×10^{-2}
(C) 1.25×10^{-2} (D) 2.5×10^{-2}

Q.5 If 10^{21} molecules are removed from 200 mg of CO_2 , then the number of moles of CO_2 left are (1983)
(A) 2.85×10^{-3} (B) 28.8×10^{-3}
(C) 0.288×10^{-3} (D) 1.68×10^{-2}

Q.6 In standardization of $Na_2S_2O_3$ using $K_2Cr_2O_7$ by iodometry, the equivalent weight of $K_2Cr_2O_7$ is (2000)
(A) $\frac{MW}{2}$ (B) $\frac{MW}{3}$ (C) $\frac{MW}{6}$ (D) $\frac{MW}{1}$

Q.7 The molarity of a solution obtained by mixing 750 mL of 0.5(M) HCl with 250 mL of 2(M) HCl will be: (2013)
(A) 0.875 M (B) 1.00 M (C) 1.75 M (D) 0.975 M

Q.8 A gaseous hydrocarbon gives upon combustion 0.72 g of water and 3.08 g of CO_2 . The empirical formula of the hydrocarbon is: (2013)
(A) C_2H_4 (B) C_3H_4 (C) C_6H_5 (D) C_7H_8

Q.9 Experimentally it was found that a metal oxide has formula $M_{0.98}O$. Metal M, present as M^{2+} and M^{3+} in its oxide. Fraction of the metal which exists as M^{3+} would be: (2013)
(A) 7.01 % (B) 4.08 % (C) 6.05 % (D) 5.08 %

Q.10 The ratio of masses of oxygen and nitrogen in a particular gaseous mixture is 1 : 4. The ratio of number of their molecule is 3 (2014)
(A) 1 : 4 (B) 7 : 32 (C) 1 : 8 (D) 3 : 16

Q.11 The molecular formula of a commercial resin used for exchanging ions in water softening is $C_6H_7SO_3Na$ (Mol. Wt. 206). What would be the maximum uptake of Ca^{2+} ions by the resin when expressed in mole per gram resin? (2015)
(A) $\frac{1}{103}$ (B) $\frac{1}{206}$ (C) $\frac{2}{309}$ (D) $\frac{1}{412}$

Previous Years' Questions

Mole Concept

Q.1 If we consider that $1/6$, in place of $1/12$, mass of carbon atom is taken to be the relative atomic mass unit, the mass of one mole of a substance will (2002)
(A) Decrease twice
(B) Increase two fold
(C) Remain unchanged
(D) Be a function of the molecular mass of the substance

Q.2 A molar solution is one that contains one mole of a solute in (1986)
(A) 1000 g of the solvent (B) One litre of the solvent
(C) One litre of the solution (D) 22.4 litres of the solution

Q.3 In the reaction,
 $2Al(s) + 6HCl(aq) \longrightarrow 2Al^{3+}(aq) + 6Cl^{-}(aq) + 3H_2(g)$ (2007)
(A) 6 l HCl (aq) is consumed for every 3L $H_2(g)$ produced

Q.12 At 300 K and 1 atm, 15 mL of a gaseous hydrocarbon requires 375 mL air containing 20% O_2 by volume for complete combustion. After combustion the gases occupy 330 mL. Assuming that the water formed is in liquid form and the volumes were measured at the same temperature and pressure, the formula of the hydrocarbon is: **(2016)**

- (A) C_2H_{12} (B) C_4H_8 (C) C_4H_{10} (D) C_3H_6

Redox Reactions

Q.13 Several blocks of magnesium are fixed to the bottom of a ship to **(2003)**

- (A) Keep away the sharks
(B) Make the ship lighter
(C) Prevent action of water and salt
(D) Prevent puncturing by under-sea rocks

Q.14 Which of the following chemical reactions depicts the oxidizing behaviour of H_2SO_4 ? **(2006)**

- (A) $2HI + H_2SO_4 \rightarrow I_2 + SO_2 + 2H_2O$
(B) $Ca(OH)_2 + H_2SO_4 \rightarrow CaSO_4 + 2H_2O$
(C) $NaCl + H_2SO_4 \rightarrow NaHSO_4 + HCl$
(D) $2PCl_5 + H_2SO_4 \rightarrow 2POCl_3 + 2HCl + SO_2Cl_2$

Q.15 The oxidation number of carbon in CH_2O is **(1982)**

- (A) -2 (B) +2 (C) 0 (D) 4

Q.16 The oxidation state of chromium in the final product formed by the reaction between KI and acidified potassium dichromate solution is **(2005)**

- (A) +4 (B) +6 (C) +2 (D) +3

Q.17 When $KMnO_4$ acts as an oxidising agent and ultimately forms $[MnO_4]^{2-}$, MnO_2 , Mn_2O_3 , Mn^{+2} then the number of electrons transferred in each case respectively is **(2002)**

- (A) 4, 3, 1, 5 (B) 1, 5, 3, 7
(C) 1, 3, 4, 5 (D) 3, 5, 7, 1

Q.18 Which of the following is a redox reaction **(2002)**

- (A) $NaCl + KNO_3 \rightarrow NaNO_3 + KCl$
(B) $CaC_2O_4 + 2HCl \rightarrow CaCl_2 + H_2C_2O_4$
(C) $Mg(OH)_2 + 2NH_4Cl \rightarrow MgCl_2 + 2NH_4OH$
(D) $Zn + 2AgCN \rightarrow 2Ag + Zn(CN)_2$


Q.19 The product of oxidation of I^- and MnO_4^- in alkaline medium is **(2004)**

- (A) IO_3^- (B) I_2 (C) IO^- (D) IO_4^-

Q.20 For H_3PO_3 and H_3PO_4 the correct choice is **(2003)**

- (A) H_3PO_3 is dibasic and reducing
(B) H_3PO_3 is dibasic and non-reducing
(C) H_3PO_4 is tribasic and reducing
(D) H_3PO_3 is tribasic and non-reducing

Q.21 Consider the following reaction:



The values of X, Y and Z in the reaction are, respectively: **(2013)**

- (A) 5, 2 and 16 (B) 2, 5 and 8
(C) 2, 5 and 16 (D) 5, 2 and 8

Q.22 In which of the following reaction H_2O_2 acts as a reducing agent? **(2014)**

- (A) $H_2O_2 + 2H^+ + 2e^- \rightarrow 2H_2O$
(B) $H_2O_2 - 2e^- \rightarrow O_2 + 2H^+$
(C) $H_2O_2 - 2e^- \rightarrow 2OH^-$
(D) $H_2O_2 + 2OH^- - 2e^- \rightarrow O_2 + 2H_2O$

- (A) (a), (b) (B) (c), (d) (C) (a), (c) (D) (b), (d)

Q.23 The equation which is balanced and represents the correct product(s) is **(2014)**

- (A) $Li_2O + 2KCl \rightarrow 2LiCl + K_2O$
(B) $[CoCl(NH_3)_5]^{+} + 5H^+ \rightarrow Co^{2+} + 5NH_4^+ + Cl^-$
(C) $[Mg(H_2O)_6]^{2+} (EDTA)^{4-} \xrightarrow{\text{excess NaOH}} [Mg(EDTA)]^{2-} + 6H_2O$
(D) $CuSO_4 + 4KCN \rightarrow K_2[Cu(CN)_4] + K_2SO_4$

Q.24 From the following statements regarding H_2O_2 , choose the incorrect statement: **(2015)**

- (A) It can act only as an oxidizing agent
(B) It decomposed on exposure to light
(C) It has to be stored in plastic or wax lined glass bottles in dark
(D) It has to be kept away from dust

JEE Advanced/Boards


Exercise 1

Mole Concept

Q.1 How many gm of HCl is needed for complete reaction with 69.6 gm MnO_2 ?




Q.2 Titanium, which is used to make air plane engines and frames, can be obtained from titanium tetrachloride, which in turn is obtained from titanium oxide by the following process:



A vessel contains 4.32 g TiO_2 , 5.76 g C and 6.82 g Cl_2 , suppose the reaction goes to completion as written, how many gram of TiCl_4 can be produced? (Ti = 48).

Q.3 Sulphuric acid is produced when sulphur dioxide reacts with oxygen and water in the presence of a catalyst:



If 5.6 mol of SO_2 reacts with 4.8 mole of O_2 and a large excess of water, what is the maximum number of moles of H_2SO_4 that can be obtained?

Q.4 What weight of Na_2CO_3 of 95% purity would be required to neutralize 45.6 mL of 0.235 N acid?

Q.5 How much $\text{BaCl}_2 \cdot 2\text{H}_2\text{O}$ and pure water to be mixed to prepare 50g of 12.0% (by wt.) BaCl_2 solution.


Q.6 To 50 litre of 0.2 N NaOH, 5 litre of 1N HCl and 15 litre of 0.1 N FeCl_3 solution are added. What weight of Fe_2O_3 can be obtained from the precipitate? Also report the normality of NaOH left in the resultant solution.

Q.7 0.5 g fuming H_2SO_4 (oleum) is diluted with water. The solution requires 26.7 mL of 0.4 N NaOH for complete neutralization. Find the percentage of free SO_3 in the sample of oleum.

Q.8 200 mL of a solution of mixture of NaOH and Na_2CO_3 was first titrated with phenolphthalein and N/10 HCl. 17.5 mL of HCl was required for the end point. After this methyl orange was added and 2.5 mL of same HCl

was again required for next end point. Find out amount of NaOH and Na_2CO_3 in mixture.

Q.9 Potassium superoxide, KO_2 , is used in rebreathing gas masks to generate oxygen:



If a reaction vessel contains 0.158 mol KO_2 and 0.10 mol H_2O , how many moles of O_2 can be produced?

Q.10 A sample of mixture of CaCl_2 and NaCl weighing 4.22 gm was treated to precipitate all the Ca as CaCO_3 which was then heated and quantitatively converted to 0.959 gm of CaO. Calculate the percentage of CaCl_2 in the mixture.

Q.11 Cyclohexanol is dehydrated to cyclohexene on heating with conc. H_2SO_4 . If the yield of this reaction is 75%, how much cyclohexene will be obtained from 100 g of cyclohexanol? $\text{C}_6\text{H}_{12}\text{O} \xrightarrow{\text{con. H}_2\text{SO}_4} \text{C}_6\text{H}_{10}$

Q.12 How many grams of 90% pure Na_2SO_4 can be produced from 250 gm of 95% pure NaCl?

Q.13 A precipitate of AgCl and AgBr weighs 0.4066 g. On heating in a current of chlorine, the AgBr is converted to AgCl and the mixture loses 0.0725 g in weight. Find the percentage of Cl in original mixture.


Q.14 How many milli-litre of 0.5 M H_2SO_4 are needed to dissolve 0.5 g of copper II carbonate?


Q.15 What is the strength in g per litre of a solution of H_2SO_4 , 12 mL of which neutralized 15 mL of N/10 NaOH solution.

Q.16 n-butane is produced by the monobromination of ethane followed by Wurtz reaction. Calculate the volume of ethane at NTP required to produce 55 g n-butane if the bromination takes place with 90% yield and the Wurtz reaction with 85% yield.

Q.17 0.50 g of a mixture of K_2CO_3 and Li_2CO_3 required 30 mL of 0.25 N HCl solution for neutralization. What is percentage composition of mixture?

Q.18 Sodium chlorate, NaClO_3 , can be prepared by the following series of reactions:





What mass of NaClO_3 can be prepared from 100 mL of concentrated HCl (density 1.18 gm/mL and 36% by mass)? Assume all other substance are present in excess amounts.

Q.19 In a determination of P an aqueous solution of NaH_2PO_4 is treated with a mixture of ammonium and magnesium ions to precipitate magnesium ammonium phosphate $\text{Mg}(\text{NH}_4)\text{PO}_4 \cdot 6\text{H}_2\text{O}$. This is heated and decomposed to magnesium pyrophosphate, $\text{Mg}_2\text{P}_2\text{O}_7$ which is weighed. A solution of NaH_2PO_4 yielded 1.054g of $\text{Mg}_2\text{P}_2\text{O}_7$. What weight of NaH_2PO_4 was present originally?

Q.20 5 mL of 8 N HNO_3 , 4.8 mL of 5 N HCl and a certain volume of 17 M H_2SO_4 are mixed together and made upto 2 litre. 30 mL of this acid mixture exactly neutralizes 42.9 mL of Na_2CO_3 solution containing 1 g $\text{Na}_2\text{CO}_3 \cdot 10\text{H}_2\text{O}$ in 100 mL of water. Calculate the amount of sulphate ions in g present in solution.

Q.21 A sample of Mg was burnt in air to give a mixture of MgO and Mg_3N_2 . The ash was dissolved in 60 Meq of HCl and the resulting solution was back titrated with NaOH . 12 Meq of NaOH were required to reach the end point. As excess of NaOH was then added and the solution distilled. The ammonia released was then trapped in 10 Meq of second acid solution. Back titration of this solution required 6 Meq of the base. Calculate the percentage of Mg burnt to the nitride.

Q.22 A mixture of ethane (C_2H_6) and ethene occupies 40 litre at 1.00 atm and at 400 K. The mixture reacts completely with 130 g of O_2 to produce CO_2 and H_2O . Assuming ideal gas behaviour, calculate the mole fractions of C_2H_4 and C_2H_6 in the mixture.

Q.23 A solid mixture 5 g consists of lead nitrate and sodium nitrate was heated below 600°C until weight of residue was constant. If the loss in weight is 28%, find the amount of lead nitrate and sodium nitrate in mixture.

Q.24 Upon mixing 45.0 mL of 0.25 M lead nitrate solution with 25 mL of 0.10 M chromic sulphate, precipitation of lead sulphate takes place. How many moles of lead sulphate are formed? Also calculate the molar concentration of the species left behind in final solution. Assume that lead sulphate is completely insoluble.

Q.25 A 10 g sample of a mixture of calcium chloride and sodium chloride is treated with Na_2CO_3 to precipitate calcium as calcium carbonate. This CaCO_3 is heated to convert all the calcium to CaO and the final mass of CaO is 1.12 gm. Calculate % by mass of NaCl in the original mixture.

Q.26 A mixture of Ferric oxide (Fe_2O_3) and Al is used as solid rocket fuel which reacts to give Al_2O_3 and Fe. No other reactants and products are involved. On complete reaction of 1 mole of Fe_2O_3 , 200 units of energy is released?


(i) Write a balance reaction representing the above change.

(ii) What should be the ratio of masses of Fe_2O_3 and Al taken so that maximum energy per unit mass of fuel is released.

(iii) What would be energy released if 16 kg of Fe_2O_3 reacts with 2.7 kg of Al.



Q.27 A mixture of nitrogen and hydrogen. In the ratio of one mole of nitrogen to three moles of hydrogen, was partially converted into so that the final product was a mixture of all these three gases. The mixture was to have a density of 0.497 g per litre at 25°C and 1.00 atm. What would be the mass of gas in 22.4 litres at 1 atm and 273 K? Calculate the % composition of this gaseous mixture by volume.

Q.28 In one process for waterproofing, a fabric is exposed to $(\text{CH}_3)_3\text{SiCl}_2$ vapour. The vapour reacts with hydroxyl groups on the surface of the fabric or with traces of water to form the waterproofing film $[(\text{CH}_3)_2\text{SiO}]_n$ by the reaction



where n stands for a large integer. The waterproofing film is deposited on the fabric layer upon layer. Each layer is 6.0 Å thick [the thickness of the $(\text{CH}_3)_2\text{SiO}$ group]. How much $(\text{CH}_3)_3\text{SiCl}_2$ is needed to waterproof one side of a piece of fabric, 1.00 m by 3.00 m, with a film 300 layers thick? The density of the film is 1.0 g/cm³.

Q.29 Two substance P_4 and O_2 are allowed to react completely to form mixture of P_4O_6 and P_4O_{10} leaving none of the reactants. Using this information calculate the composition of final mixture when mentioned amount of P_4 and O_2 are taken.



(i) If 1 mole P_4 & 4 mole of O_2

(ii) If 3 mole P_4 & 11 mole of O_2

(iii) If 3 mole P_4 & 13 mole of O_2

Q.30 Chloride samples are prepared for analysis by using NaCl , KCl and NH_4Cl separately or as a mixture. What minimum volume of 5% by weight AgNO_3 solution (sp. gr., 1.04 g mL⁻¹) must be added to a sample of 0.3 g in order to ensure complete precipitation of chloride in every possible case?

Q.31 124 gm of mixture containing NaHCO_3 , AlCl_3 , and KNO_3 requires 500 mL, 8% w/w NaOH solution [$d_{\text{NaOH}} = 1.8 \text{ gm/mL}$] for complete neutralisation. On heating same amount of mixture, it known loss in weight of 18.6 gm. Calculate % composition of mixture by moles. Weak base formed doesn't interfere in reaction. Assume KNO_3 does not decompose under given conditions.

Q.32 If the yield of chloroform obtainable from acetone and bleaching powder is 75%. What is the weight of acetone required for producing 30 gm of chloroform?

Q.33 A sample of impure Cu_2O contains 66.67% of Cu . What is the percentage of pure Cu_2O in the sample?

Q.34 Equal weights of mercury and iodine are allowed to react completely to form a mixture of mercurous and mercuric iodide leaving none of the reactants. Calculate the ratio by weight of Hg_2I_2 and HgI_2 and formed. ($\text{Hg} = 200$, $\text{I} = 127$)

Redox Reactions

Q.1 Indicate the oxidation state of underlined in each case:

- | | | |
|---|--|--|
| (i) $\text{Na}\underline{\text{N}}\text{O}_2$ | (b) $\underline{\text{H}}$ | (c) $\underline{\text{Cl}}_2\text{O}_7$ |
| (ii) $\text{K}\underline{\text{Cr}}\text{O}_3\text{Cl}$ | (e) $\underline{\text{Ba}}\text{Cl}_2$ | (f) $\underline{\text{I}}\text{Cl}_3$ |
| (iii) $\text{K}_2\underline{\text{Cr}}_2\text{O}_7$ | (h) $\underline{\text{C}}\text{H}_2\text{O}$ | (i) $\underline{\text{Ni}}(\text{CO})_4$ |
| (iv) $\underline{\text{N}}\text{H}_2\text{OH}$ | | |

Q.2 Indicate the each reaction which of the reactant is oxidized or reduced if any:

- $\text{CuSO}_4 + 4\text{KI} \longrightarrow 2\text{CuI} + \text{I}_2 + 2\text{K}_2\text{SO}_4$
- $2\text{Na}_2\text{S} + 4\text{HCl} + \text{SO}_2 \longrightarrow 4\text{NaCl} + 3\text{S} + 2\text{H}_2\text{O}$
- $\text{NH}_4\text{NO}_2 \xrightarrow{\Delta} \text{N}_2 + 2\text{H}_2\text{O}$

Q.3 Calculate the number of electrons lost or gained during the changes:

- $3\text{Fe} + 4\text{H}_2\text{O} \longrightarrow \text{Fe}_3\text{O}_4 + 4\text{H}_2$
- $\text{AlCl}_3 + 3\text{K} \longrightarrow \text{Al} + 3\text{KCl}$

Q.4 Explain, why?

- H_2S acts as reductant whereas, SO_2 acts as reductant and oxidant both.
- H_2O_2 acts as reductant and oxidant both.

Q.5 MnO_4^- can oxidize NO_2^- to NO_3^- in basic medium. How many mol of NO_2^- are oxidized by 1 mol of MnO_4^- ?

Q.6 Which is stronger base in each pair?

- HSO_4^- ; HSO_3^-
- NO_2^- ; NO_3^-
- Cl^- ; ClO^-

Q.7 Fill in the blanks and balance the following equations:

- $\text{Zn} + \text{HNO}_3 \longrightarrow \dots + \text{N}_2\text{O} + \dots$
- $\text{HI} + \text{HNO}_3 \longrightarrow \dots + \text{NO} + \text{H}_2\text{O} \dots$

Q.8 What volume of 0.20 M H_2SO_4 is required to produce 34.0 g of H_2S by the reaction:



Q.9 20 mL of 0.2 M MnSO_4 are completely oxidized by 16 mL of KMnO_4 of unknown normality, each forming Mn^{4+} oxidation state. Find out the normality and molarity of KMnO_4 solution.

Q.10 KMnO_4 solution is to be standardized by titration against $\text{As}_2\text{O}_3(\text{s})$. A 0.1097 g sample of As_2O_3 requires 26.10 mL of the KMnO_4 solution for its titration. What are the molarity and normality of the KMnO_4 solution?

Q.11 0.518 g sample of limestone is dissolved and then Ca is precipitated as CaC_2O_4 . After filtering and washing the precipitate, it requires 40 mL of 0.25 N KMnO_4 solution to equivalence point. What is percentage of CaO in limestone?



Q.12 20 mL of a solution containing 0.2 g of impure sample of H_2O_2 reacts with 0.316 g of KMnO_4 (acidic). Calculate:

- Purity of H_2O_2 ,
- Volume of dry O_2 evolved at 27°C and 750 mm P.





Q.13 5.7 g of bleaching powder was suspended in 500 mL of water. 25 mL of this suspended on treatment with KI and HCl liberated iodine which reacted with 24.35 mL of N/10 $\text{Na}_2\text{S}_2\text{O}_3$. Calculate percentage of available Cl_2 in bleaching powder.

Q.14 Balance the following equation:



- $\text{C}_2\text{H}_5\text{OH} + \text{K}_2\text{Cr}_2\text{O}_7 + \text{H}_2\text{SO}_4 \longrightarrow \text{C}_2\text{H}_4\text{O}_2 + \text{Cr}_2(\text{SO}_4)_3 + \text{K}_2\text{SO}_4 + \text{H}_2\text{O}$
- $\text{As}_2\text{S}_5 + \text{HNO}_3 \longrightarrow \text{NO}_2 + \text{H}_2\text{O} + \text{H}_3\text{AsO}_4 + \text{H}_2\text{SO}_4$



Q.15 Balance the following equations:



Q.16 Mg can reduce NO_3^- to NH_3 in basic solution:




A 25.0 mL sample of NO_3^- solution was treated with Mg. The $\text{NH}_3(\text{g})$ was passed into 50 mL of 0.15 N HCl. The excess HCl required 32.10 mL of 0.10 M NaOH for its neutralization. What was the molarity of NO_3^- ions in the original sample?

Q.17 An acid solution of KReO_4 sample containing 26.83 mg of combined rhenium was reduced by passage through a column of granulated zinc. The effluent solution including the washing from the column, was then titrated with 0.05 N KMnO_4 . 11.45 mL of the standard KMnO_4 was required for the reoxidation of all the rhenium to the perrhenate ion ReO_4^- . Assuming that rhenium was the only element reduced, what is the oxidation state to which rhenium was reduced by the zinc column.


Q.18 100 mL solution of FeC_2O_4 and FeSO_4 is completely oxidized by 60 mL of 0.02 M in acid medium. The resulting solution is then reduced by Zn and dil.HCl. The reduced solution is again oxidized completely by 40 mL of 0.02 M KMnO_4 . Calculate normality of FeC_2O_4 and FeSO_4 in mixture.

Q.19 1 g of most sample of KCl and KClO_3 was dissolved in water to make 250 mL solution, 25 mL of this solution was treated with SO_2 to reduce chlorate to chloride and excess of SO_2 was removed by boiling. The total chloride was precipitated as silver chloride. The weight of precipitate was 0.1435 g. In another experiment, 25 mL of original solution was heated with 30 mL of 0.2 N ferrous sulphate solution and unreacted ferrous sulphate required 37.5 mL of 0.08 N solution of an oxidant for complete oxidation. Calculate the molar


ratio of chlorate to chloride in the given mixture. Fe^{2+} reacts with ClO_3^- according to equation.



Q.20 (i) CuSO_4 reacts with KI in acidic medium to liberate I_2



(ii) Mercuric per iodate $\text{Hg}_5(\text{IO}_6)_2$ reacts with a mixture of KI and HCl following the equation:



(iii) The liberated iodine is titrated against $\text{Na}_2\text{S}_2\text{O}_3$ solution. One mL of which is equivalent to 0.0499 g of $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$. What volume in mL of $\text{Na}_2\text{S}_2\text{O}_3$ solution will be required to react with I_2 liberated from 0.7245 g of $\text{Hg}_5(\text{IO}_6)_2$? M. wt. of $\text{Hg}_5(\text{IO}_6)_2 = 1448.5$ and M. wt. of $\text{CuSO}_4 \cdot 5\text{H}_2\text{O} = 249.5$.

Q.21 1.249 g of a sample of pure BaCO_3 and impure CaCO_3 containing some CaO was treated with dil. HCl and it evolved 168 mL of CO_2 at NTP. From this solution BaCrO_4 was precipitated, filtered and washed. The dry precipitate was dissolved in dilute H_2SO_4 and dilute to 100 mL. 10 mL of this solution when treated with KI solution liberated iodine which required exactly 20 mL of 0.05 N $\text{Na}_2\text{S}_2\text{O}_3$. Calculate percentage of CaO in the sample.

Q.22 A 10 g mixture of Cu_2S and CuS was treated with 200 mL of 0.75 M MnO_4^- in acid solution producing SO_2 , Cu^{2+} and Mn^{2+} . The SO_2 was boiled off and the excess of MnO_4^- was treated with 175 mL of 1 M Fe^{2+} solution. Calculate percentage of CuS in original mixture.

Q.23 For estimating ozone in the air, a certain volume of air is passed through an acidified or neutral KI solution when oxygen is evolved and iodide is oxidised to give iodine. When such a solution is acidified, free iodine is evolved which can be titrated with standard $\text{Na}_2\text{S}_2\text{O}_3$ solution. In an experiment 10 litre of air at 1 atm and 27°C were passed through an alkaline KI solution, at the end, the iodine entrapped in a solution on titration as above required 1.5 mL of 0.01 N $\text{Na}_2\text{S}_2\text{O}_3$ solution. Calculate volume percentage of O_3 in sample.

Q.24 30 mL of an acidified solution of 1.5 N MnO_4^- ions, 15 mL of 0.5 N oxalic acid and 15 mL of 0.4 N ferrous salt solution are added together. Find the molarities of MnO_4^- and Fe^{3+} ions in the final solution?

Q.25 (i) 25 mL of H_2O_2 solution were added to excess of acidified solution of KI. The iodine so liberated required 20 mL of 0.1 N $\text{Na}_2\text{S}_2\text{O}_3$ for titration. Calculate the strength of H_2O_2 in terms of normality, percentage and volume.

(ii) To a 25 mL H_2O_2 solution, excess of acidified solution of KI was added. The iodine liberated required 20 mL of 0.3 N sodium thiosulphate solution. Calculate the volume strength of H_2O_2 solution.

Q.26 An aqueous solution containing 0.10 g KIO_3 (formula weight = 214.0) was treated with an excess of KI solution. The solution was acidified with HCl. The liberated I_2 consumed 45 mL of thiosulphate solution to decolorise the blue starch-iodine complex. Calculate the molarity of the sodium thiosulphate solution.


Q.27 A sample of $\text{MnSO}_4 \cdot 4\text{H}_2\text{O}$ is strongly heated in air. The residue (Mn_3O_4) left was dissolved in 100 mL of 0.1 N FeSO_4 containing dil. H_2SO_4 . This solution was completely reacted with 50 mL of KMnO_4 solution. 25 mL of this KMnO_4 solution was completely reduced by 30 mL of 0.1 N FeSO_4 solution. Calculate the amount of $\text{MnSO}_4 \cdot 4\text{H}_2\text{O}$ in sample.

Q.28 Write complete balanced equation for the following in acidic medium by ion-electron method:

- $\text{ClO}_3^- + \text{Fe}^{2+} \rightarrow \text{Cl}^- + \text{Fe}^{+3} + \text{H}_2\text{O}$
- $\text{CuS} + \text{NO}_3^- \rightarrow \text{Cu}^{+2} + \text{S}_8 + \text{NO} + \text{H}_2\text{O}$
- $\text{S}_2\text{O}_3^{2-} + \text{Sb}_2\text{O}_3 \rightarrow \text{SbO} + \text{H}_2\text{SO}_3$
- $\text{HCl} + \text{KMnO}_4 \rightarrow \text{Cl}_2 + \text{KCl} + \text{MnCl}_2 + \text{H}_2\text{O}$
- $\text{KClO}_3 + \text{H}_2\text{SO}_4 \rightarrow \text{KHSO}_4 + \text{HClO}_4 + \text{ClO}_3 + \text{H}_2\text{O}$
- $\text{HNO}_3 + \text{HBr} \rightarrow \text{NO} + \text{Br}_2 + \text{H}_2\text{O}$
- $\text{IO}_4^- + \text{I}^- + \text{H}^+ \rightarrow \text{I}_2 + \text{H}_2\text{O}$

Q.29 Balance the following equations by oxidation method:

- $\text{Cu} + \text{NO}_3^- + \dots \rightarrow \text{Cu}^{+2} + \text{NO}_2 + \dots$
(Acid medium)
- $\text{Cl}_2 + \text{IO}_3^- + \text{OH}^- \rightarrow \text{IO}_4^- + \dots + \text{H}_2\text{O}$
(Basic medium)
- $\text{H}_2\text{S} + \text{K}_2\text{CrO}_4 + \text{H}_2\text{SO}_4 \rightarrow \dots$
(Acid medium)
- $\text{Fe}^{+2} + \text{MnO}_4^- \rightarrow \text{Fe}^{+3} + \text{Mn}^{+2} + \dots$
(Acid medium)
- $\text{KMnO}_4 + \text{H}_2\text{SO}_4 + \text{H}_2\text{O}_2 \rightarrow \dots$
 $\text{K}_2\text{SO}_4 + \text{MnSO}_4 + \text{H}_2\text{O} + \dots$
(Acid medium)



Q.30 Write complete balance equation for the following in basic medium by ion-electron method:

- $\text{S}_2\text{O}_4^{2-} + \text{Ag}_2\text{O} \rightarrow \text{Ag} + \text{SO}_3^{2-}$
- $\text{Cl}_2 + \text{OH}^- \rightarrow \text{Cl}^- + \text{ClO}^-$
- $\text{H}_2 + \text{ReO}_4^- \rightarrow \text{ClO}_2^- + \text{Sb}(\text{OH})_6^-$
- $\text{I}_2 + \text{OH}^- \rightarrow \text{I}^- + \text{IO}_3^-$
- $\text{MnO}_4^- + \text{Fe}^{+2} \rightarrow \text{Mn}^{+2} + \text{Fe}^{+3}$

Exercise 2

Mole Concept

Single Correct Choice Type

Q.1 'x' gms of an element 'A' on heating in a jar of chlorine give 'y' gms of ACl_2 the atomic weight of element A is

- (A) $\left(\frac{x}{y-71}\right) \times \frac{35.5}{2}$ (B) $\frac{71x}{y-71}$
(C) $\frac{35.5x}{y-71}$ (D) None of these

Q.2 The amount of H_2SO_4 present in 1200 mL of 0.2 N solution is

- (A) 10.76 gms (B) 11.76 gms
(C) 12.76 gms (D) 14.76 gms

Q.3 An iodized salt contains 0.5% of NaI. A person consumes 3 gm of salt everyday. The number of iodide ions going into his body everyday is

- (A) 10^{-4} (B) 6.02×10^{-4}
(C) 6.02×10^{19} (D) 6.02×10^{23}

Assertion Reasoning Type

- (A) If both statement-I and statement-II are true and statement-II is the correct explanation of statement-I, the mark (A).
(B) If both statement-I and statement-II are true and statement-II is not the correct explanation of statement -I, the mark (B).
(C) If statement-I is true but statement-II is false, then mark (C).
(D) If both statement-I and statement-II are false, then mark (D).

Q.4 Statement-I: 0.28 g of N_2 has equal volume as 0.44 g of another gas at same conditions of temperature and pressure.

Statement-II: Molecular mass of another gas is 44 g mol^{-1} .

Q.5 Statement-I: Boron has relative atomic mass 10.81.

Statement-II: Boron's two isotopes, $^{10}_5\text{B}$ and $^{11}_5\text{B}$ and their relative abundance is 19% and 81%.

Q.6 Statement-I: The percentage of nitrogen in urea is 46%.

Statement-II: Urea is ionic compound.

Q.7 Statement-I: The oxidation state of central sulfur of $\text{Na}_2\text{S}_2\text{O}_3$ is +6.

Statement-II: Oxidation state of an element should be determined from structure.

Q.8 Statement-I: Molarity of a solution and molality of a solution both change with density.




Statement-II: Density of the solution changes when percentage by mass of solution changes.

Q.9 Statement-I: $2A + 3B \rightarrow C$, $4/3$ moles of 'C' are always produced when 3 moles of 'A' and 4 moles of 'B' are added.

Statement-II: 'B' is the limiting reactant for the given data.

Multiple Correct Choice Type

Q.10 Given following series of reactions:



Select the correct option(s):

(A) Moles of HNO_3 obtained is half of moles of Ammonia used if HNO_2 is not used to produce HNO_3 by equation (iv)

(B) 100/6% more HNO_3 will be produced if HNO_2 is used to produce HNO_3 by reaction (iv) than if HNO_2 is not used to produce HNO_3 by reaction (iv)

(C) If HNO_2 is used to produce HNO_3 then 1/4th of total is produced by reaction (iv)

(D) Moles of NO produced in reaction (iv) is 50% of moles of total HNO_3 produced.

Comprehension Type

Paragraph 1: Normality is number of gram equivalents dissolved per litre of solution. It changes with change in temperature. In case of monobasic acid, normality and molarity are equal but in case of dibasic acid, normality is twice the molarity. In neutralization and redox reactions, number of mill equivalents of reactants as well as products are always equal.

Q.11 On heating a litre of a $\frac{N}{2}$ HCl solution, 2.750 g of HCl is lost and the volume of solution becomes 750 mL. The normality of resulting solution will be

- (A) 0.58 (B) 0.75 (C) 0.057 (D) 5.7

Q.12 The volume of 0.1 M $\text{Ca}(\text{OH})_2$ required to neutralize 10 mL of 0.1 N HCl will be

- (A) 10 mL (B) 20 mL (C) 5 mL (D) 40 mL

Q.13 Molarity of 0.5 N Na_2CO_3 is

- (A) 0.25 (B) 1.0 (C) 0.5 (D) 0.125

Q.14 6.90 N KOH solution in water contains 30% by weight of KOH. The density of solution will be

- (A) 1.288 (B) 2.88 (C) 0.1288 (D) 12.88

Q.15 The amount of ferrous ammonium sulphate required to prepare 250 mL of 0.1 N solution is

- (A) 1.96 g (B) 1.8 g (C) 9.8 g (D) 0.196 g

Paragraph 2: A 4.925 g sample of a mixture of CuCl_2 and CuBr_2 was dissolved in water and mixed thoroughly with a 5.74 g portion of AgCl. After the reaction and solid, a mixture of AgCl and AgBr, was filtered, washed, and dried. Its mass was found to be 6.63 g.

Q.16

(1) % By mass of CuBr_2 in original mixture is

- (A) 2.24 (B) 74.5 (C) 45.3 (D) None

(2) % By mass of Cu in original mixture is

- (A) 38.68 (B) 19.05 (C) 3.86 (D) None

(3) % by mole of AgBr in dried precipitate is

- (A) 25 (B) 50 (C) 75 (D) 60

(4) No. of moles of Cl^- ion present in the solution after precipitate ion are

- (A) 0.06 (B) 0.02 (C) 0.04 (D) None

Paragraph 3: Water is added to 3.52 grams of UF_6 . The products are 3.08 grams of a solid [containing only U, O and F] and 0.8 gram of a gas only. The gas [containing fluorine and hydrogen only], contains 95% by mass fluorine.

[Assume that the empirical formula is same as molecular formula.]

Q.17

(1) The empirical formula of the gas is

- (A) HF_2 (B) H_2F (C) HF (D) HF_3

(2) The empirical formula of the solid product is

- (A) UF_2O_2 (B) UFO_2 (C) UF_2O (D) UFO

(3) The percentage of fluorine of the original compound which is converted into gaseous compound is

- (A) 66.66% (B) 33.33% (C) 50% (D) 89.9%

Match the Columns

Q.18 One type of artificial diamond (commonly called YAG for yttrium aluminium garnet) can be represented by the formula $\text{Y}_3\text{Al}_5\text{O}_{12}$. [Y = 89, Al = 27]

Column I	Column II
(A) Y	(p) 22.73%
(B) Al	(q) 32.32%
(C) O	(r) 44.95%

Q.19 The recommended daily dose is 17.6 milligrams of vitamin C (ascorbic acid) having formula $\text{C}_6\text{H}_8\text{O}_6$. Match the following. Given: $N_A = 6 \times 10^{23}$

Column I	Column II
(A) O-atoms present	(p) 10^{-4} mole
(B) Moles of vitamin C in 1 gm of vitamin C	(q) 5.68×10^{-3}
(C) Moles of vitamin C in 1 gm should be consumed daily	(r) 3.6×10^{20}

Q.20 If volume strength of H_2O_2 solution is 'X-V' then its

Column I	Column II
(i) Strength in g/L	(p) $\frac{X}{11.2}$
(ii) Volume strength X	(q) $\frac{X}{5.6}$
(iii) Molarity	(r) $\frac{17X}{5.6}$
(iv) Normality	(s) $5.6 \times N$

(A) (i) - r, (ii) - p, (iii) - s, (iv) - q

(B) (i) - s, (ii) - p, (iii) - q, (iv) - p

(C) (i) - r, (ii) - s, (iii) - p, (iv) - q

(D) (i) - r, (ii) - q, (iii) - s, (iv) - p

Q.21 Match the entries in column I with entries in column II and then pick out correct options:

Column I	Column II
(i) M_R on mixing two acidic solutions	(p) $\frac{x \times d \times 10}{M_{\text{solute}}}$
(ii) M_R on mixing two basic solutions	(q) $n \times M \times V \text{ mL}$
(iii) M_R on mixing two acidic and basic solutions	(r) $\frac{M_1 V_1 - M_2 V_2}{V_1 + V_2}$
(iv) Milliequivalent	(s) $\frac{M_1 V_1}{V_2}$
(v) Molarity	(t) $\frac{M_1 V_1 + M_2 V_2}{V_1 + V_2}$

(A) (i) - p, (ii) - r, (iii) - p, (iv) - q, (v) - s

(B) (i) - t, (ii) - t, (iii) - r, (iv) - q, (v) - p, s

(C) (i) - q, (ii) - p, (iii) - q, (iv) - r, (v) - q

(D) (i) - p, (ii) - q, (iii) - q, (iv) - r, (v) - r

Redox Reactions

Single Correct Choice Type

Q.1 One mole of N_2H_4 loses ten moles of electrons to form a new compound Y. Assuming that all the nitrogen appears in the new compound, what is the oxidation state of nitrogen in Y? (There is no change in the oxidation state of hydrogen)

- (A) -1 (B) -3 (C) +3 (D) +5

Q.2 Which is best reducing agent

- (A) F^- (B) Cl^- (C) Br^- (D) I^-


Q.3 In the aluminothermite process, aluminium acts as

- (A) An oxidizing agent (B) A flux
(C) Reducing agent (D) A solder

Q.4 Zinc-copper couple that can be used as a reducing agent is obtained by

- (A) Mixing zinc dust and copper gas
- (B) Zinc coated with copper
- (C) Copper coated with zinc
- (D) Zinc and copper wires welded together

Q.5 In the following equations value of X is



- (A) 4e^-
- (B) 5e^-
- (C) 6e^-
- (D) 7e^-


Q.6 The brown ring complex compound is formulated as $[\text{Fe}(\text{H}_2\text{O})_5(\text{NO})^+]\text{SO}_4^-$. The oxidation state of iron is

- (A) 1
- (B) 2
- (C) 3
- (D) 0

Q.7 Oxidation state of oxygen atom in potassium superoxide is

- (A) $-1/2$
- (B) -1
- (C) -2
- (D) 0

Q.8 In the following reaction



- (A) Bromine is both reduced and oxidised
- (B) Bromine is neither reduced nor oxidised
- (C) Bromine is oxidised and carbonate is reduced
- (D) Bromine is reduced and water is oxidised

Comprehension Type

Paragraph 1: The redox titration involve the chemical reaction between the oxidising agent and reducing agent in aqueous solutions under suitable conditions. Titrations involve the direct use of iodine as oxidising agent are known as iodimetric titrations while those titrations involving indirect use of iodine are known as iodometric titrations. These titrations are used for the estimation of oxidising agents like KMnO_4 , $\text{K}_2\text{Cr}_2\text{O}_7$, CuSO_4 etc.

Q.9 50 mL of an aqueous solution of H_2O_2 was treated with excess of KI solution and the iodine so liberated quantitatively required 20 mL of 0.1 N solution of hypo. This titration is known as:

- (A) Iodometric titration
- (B) Iodimetric titration
- (C) Potassium iodide titration
- (D) All of these

Q.10 In the above problem, concentration of H_2O_2 in gm/litre is:

- (A) 6.8
- (B) 0.68
- (C) 0.068
- (D) 0.34

Q.11 0.5 gm sample of pyrolusite (MnO_2) is treated with HCl, the Cl_2 gas evolved is treated with KI, the violet vapours evolved are absorbed in 30 mL 0.1 N $\text{Na}_2\text{S}_2\text{O}_3$ solution percentage purity of pyrolusite sample is

- (A) 30%
- (B) 50%
- (C) 36%
- (D) 26.1%

Q.12 Arsenite gets converted into arsenate by using iodine, valency factor for Arsenite and Iodine are respectively

- (A) 2 and 2
- (B) 2 and 1
- (C) 1 and 2
- (D) 5 and 2

Paragraph 2: Oxidation and reduction process involves the transaction of electrons. Loss of electrons is oxidation and the gain of electrons is reduction. It is thus obvious that in a redox reaction, the oxidant is reduced by accepting the electrons and the reductant is oxidised by losing electrons. The reactions in which a species disproportionate into two oxidation states (lower and higher) are called disproportionation reactions. In electrochemical cells, redox reaction is involved, i.e., oxidation takes place at anode and reduction at cathode.

Q.13 The reaction: $\text{Cl}_2 \rightarrow \text{Cl}^- + \text{ClO}_3^-$ is

- (A) Oxidation
- (B) Reduction
- (C) Disproportionation
- (D) Neither oxidation nor reduction

Q.14 In the reaction: $\text{I}_2 + 2\text{S}_2\text{O}_3^{2-} \rightarrow 2\text{I}^- + \text{S}_4\text{O}_6^{2-}$

- (A) I_2 is reducing agent
- (B) I_2 is oxidising agent
- (C) $\text{S}_2\text{O}_3^{2-}$ is reducing agent
- (D) $\text{S}_2\text{O}_3^{2-}$ is oxidising agent

Q.15 Determine the change in oxidation number of sulphur is H_2S and SO_2 respectively in the following reaction: $2\text{H}_2\text{S} + \text{SO}_2 \rightarrow 2\text{H}_2\text{O} + 3\text{S}$

- (A) 0, +2
- (B) +2, -4
- (C) -2, +2
- (D) +4, 0

Multiple Correct Choice Type

Q.16 Which of the following reactions is/are correctly indicated?

Oxidant Reductant

- (A) $\text{HNO}_3 + \text{Cu} \longrightarrow \text{Cu}^{2+} + \text{NO}_2$
- (B) $2\text{Zn} + \text{O}_2 \longrightarrow \text{ZnO}$
- (C) $\text{Cl}_2 + 2\text{Br}^- \longrightarrow 2\text{Cl}^- + \text{Br}_2$
- (D) $4\text{Cl}_2 + \text{CH}_4 \longrightarrow \text{CCl}_4 + 4\text{HCl}$


Assertion and Reasoning Type

Each of the questions given below consist of statement-I and statement-II. Use the following Key to choose the appropriate answer.

- (A) If both statement-I and statement-II are true, and statement-II is the correct explanation of statement-I.
(B) If both statement-I and statement-II are true, and statement-II is not the correct explanation of statement-I.
(C) If statement-I is true but statement-II is false.
(D) If statement-I is false but statement-II is true.

Q.17 Statement-I: In CrO_5 oxidation number of Cr is +6.

Statement-II: CrO_5 has butterfly structure in which



peroxide bonds are present.

Q.18 Statement-I: In PbO_4 all Pb has +8/3 oxidation number.

Statement-II: PbO_4 is mixed oxide of PbO and PbO_2


Q.19 Statement-I: HClO_4 is only oxidising agent.

Statement-II: Cl is most electro-negative element in H, Cl and O.

Q.20 Statement-I: In FeS_2 oxidation number of iron is +4.

Statement-II: In FeS_2 ($\text{S}^- - \text{S}^-$) linkage is present.

Q.21 Statement-I: In given reaction H_2O_2 is oxidising & reducing agent



Statement-II: In H_2O_2 is a bleaching reagent.

Q.22 Statement-I: In basic medium colour of $\text{K}_2\text{Cr}_2\text{O}_7$ is changed from orange to yellow.

Statement-II: In basic medium $\text{K}_2\text{Cr}_2\text{O}_7$ is changed in chromate ion.

Q.23 Statement-I: $\text{I}_2 \longrightarrow \text{IO}_3^- + \text{I}^-$.

This reaction is disproportionation reaction.

Statement-II: Oxidation number of I can vary from -1 to +7.

Match the Columns

Q.24 Match the entries in column I with entries in column II and then pick out correct options:

Column I	Column II
(A) Increase in oxidation number	(p) Loss of electrons
(B) Decrease in oxidation number	(q) Redox reaction
(C) Oxidising agent	(r) Fractional oxidation number
(D) Reducing agent	(s) Zero oxidation number
(E) $2\text{Cu}^+ \rightarrow \text{Cu}^{2+} + \text{Cu}$	(t) Simple neutralisation reaction
(F) $\text{MnO}_2 + 4\text{HCl} \rightarrow \text{MnCl}_2$	(u) Gain of electrons + $\text{Cl}_2 + 2\text{H}_2\text{O}$
(G) Mn_3O_4	(v) Disproportionation
(H) CH_2Cl_2	(w) Oxidation
(I) $\text{NaOH} + \text{HCl} \rightarrow \text{NaCl} + \text{H}_2\text{O}$	(x) Reduction

Q.25 Match the reactions in column I with nature of the reactions/type of the products in Column II.

Column I	Column II
(A) $\text{O}_2^- \rightarrow \text{O}_2 + \text{O}_2^{2-}$	(p) Redox reaction
(B) $\text{CrO}_4^{2-} + \text{H}^+ \rightarrow$	(q) One of the products has trigonal planar structure
(C) $\text{MnO}_4^- + \text{NO}_2^- + \text{H}^+ \rightarrow$	(r) Dimeric bridged tetrahedral metal ion
(D) $\text{NO}_3^- + \text{H}_2\text{SO}_4 + \text{Fe}^{2+} \rightarrow$	(s) Disproportionation

Previous Years' Questions

Mole Concept

Q.1 Naturally occurring boron consists of two isotopes whose atomic weights are 10.01 and 11.01. The atomic weight of natural boron is 10.81. Calculate the percentage of each isotope in natural boron. **(1978)**

Q.2 The vapour density (hydrogen = 1) of a mixture consisting of NO_2 and N_2O_4 is 38.3 at 26.7°C . Calculate the number of moles of NO_2 in 100 g of the mixture. **(1979)**

Q.3 A solid mixture (5.0 g) consisting of lead nitrate and sodium nitrate was heated below 600°C until the weight

of the residue was constant. If the loss in weight is 28.0 percent, find the amount of lead nitrate and sodium nitrate in the mixture. **(1990)**

Q.4 'A' is a binary compound of a univalent metal. 1.422 g of A reacts completely with 0.321 g of sulphur in an evacuated and sealed tube to give 1.743 g of a white crystalline solid B, that forms a hydrated double salt, C with $\text{Al}_2(\text{SO}_4)_3$. Identify A, B and C. **(1994)**

Q.5 Calculate the molality of 1.0 L solution of 93% H_2SO_4 (weight/volume). The density of the solution is 1.84 g/mL. **(1990)**

Q.6 20% surface sites have adsorbed N_2 . On heating N_2 gas evolved from sites and were collected at 0.001 atm and 298 K in a container of volume is 2.46 cm^3 . Density of surface sites is $6.023 \times 10^{14} / \text{cm}^2$ and surface area is 1000 cm^2 , find out the number of surface sites occupied per molecule of N_2 . **(2005)**

Q.7 If 0.50 mole of BaCl_2 is mixed with 0.20 mole of Na_3PO_4 the maximum number of moles of $\text{Ba}_3(\text{PO}_4)_2$ that can be formed is **(1981)**

- (A) 0.70 (B) 0.50 (C) 0.20 (D) 0.10

Q.8 In the standardization of $\text{Na}_2\text{S}_2\text{O}_3$ using $\text{K}_2\text{Cr}_2\text{O}_7$ is **(2001)**

(A) $\left(\frac{\text{Molecular Weight}}{2} \right)$

(B) $\left(\frac{\text{Molecular Weight}}{6} \right)$

(C) $\left(\frac{\text{Molecular Weight}}{3} \right)$

(D) Same as molecular weight

Q.9 The difference in the oxidation numbers of the two types of sulphur atoms in $\text{Na}_2\text{S}_4\text{O}_6$ is. **(2011)**

Read the following questions and answer as per the direction given below:

(A) Statement-I is true; statement-II is true; statement-II is the correct explanation of statement-I.

(B) Statement-I is true; statement-II is true; statement-II is not the correct explanation of statement-I.

(C) Statement-I is true; statement-II is false.

(D) Statement-I is false; statement-II is true.

Q.10 Statement-I: In the titration of Na_2CO_3 with HCl using methyl orange indicator, the volume required at the equivalent point is twice that of the acid required using phenolphthalein indicator.

Statement-II: Two moles of HCl are required for the complete neutralization of one mole of Na_2CO_3 . **(1991)**

Q.11 2.68×10^{-3} moles of a solution containing an ion A^{n+} require 1.61×10^{-3} moles of MnO_4^- for the oxidation of A^{n+} to AO_3^- in acidic medium. What is the value of n? **(1984)**

Q.12 A 5.0 cm^3 solution of H_2O_2 liberates 0.508 g of iodine from an acidified KI solution. Calculate the strength of H_2O_2 solution in terms of volume strength at STP. **(1995)**

Q.13 A solution of 0.2 g of a compound containing Cu^{2+} and $\text{C}_2\text{O}_4^{2-}$ ions on titration with 0.02 M KMnO_4 in presence of H_2SO_4 consumes 22.6 mL of the oxidant. The resultant solution is neutralized with NaCO_3 , acidified with dilute acetic acid and treated with excess KI. The liberated iodine requires 11.3 mL of 0.05 M $\text{Na}_2\text{S}_2\text{O}_3$ solution for complete reduction. Find out the mole ratio of Cu^{2+} to $\text{C}_2\text{O}_4^{2-}$ in the compound. Write down the balanced redox reactions involved in the above titrations. **(1991)**

Q.14 A mixture of $\text{H}_2\text{C}_2\text{O}_4$ (oxalic acid) and NaHC_2O_4 weighing 2.02 g was dissolved in water and the solution made up to one litre. Ten millilitres of the solution required 3.0 mL of 0.1 N sodium hydroxide solution for complete neutralization. In another experiment, 10.0 mL of the same solution, in hot dilute sulphuric acid medium, required 4.0 mL of 0.1 N potassium permanganate solution for complete reaction. Calculate the amount of $\text{H}_2\text{C}_2\text{O}_4$ and NaHC_2O_4 in the mixture. **(1990)**

Q.15 The unbalanced chemical reactions given in list I show missing or condition which are provided in list II. Match list I with list II and select the correct answer using the code given below the lists: **(2013)**

	List I		List II
(i)	$\text{PbO}_2 + \text{H}_2\text{SO}_4 \xrightarrow{?}$ $\text{PbSO}_4 + \text{O}_2 + \text{other product}$	(p)	NO
(ii)	$\text{Na}_2\text{S}_2\text{O}_3 + \text{H}_2\text{O} \xrightarrow{?}$ $\text{NaHSO}_4 + \text{other product}$	(q)	I_2
(iii)	$\text{N}_2\text{H}_4 \xrightarrow{?}$ $\text{N}_4 + \text{other product}$	(r)	Warm
(iv)	$\text{XeF}_2 \xrightarrow{?}$ $\text{Xe} + \text{other product}$	(s)	Cl_2

Codes:

	(i)	(ii)	(iii)	(iv)
(p)	4	2	3	1
(q)	3	2	1	4
(r)	1	4	2	3
(s)	3	4	2	1

Q.16 For the reaction $\text{I}^- + \text{ClO}_3^- + \text{H}_2\text{SO}_4 \rightarrow \text{Cl}^- + \text{HSO}_4^- + \text{I}_2$

The correct statement(s) in the balanced equation is/are:

(2014)

- (A) Stoichiometric coefficient of HSO_4^- is 6.
- (B) Iodide is oxidized.
- (C) Sulphur is reduced.
- (D) H_2O is one of the products

Q.17 Hydrogen peroxide in its reaction with KIO_4 and NH_2OH respectively, is acting as a

(2014)

- (A) Reducing agent, oxidising agent
- (B) Reducing agent, reducing agent
- (C) Oxidising agent, oxidising agent
- (D) Oxidising agent, reducing agent

PlancEssential Questions

JEE Main/Boards

Exercise 1

Mole Concept

Q.1	Q.3	Q.7
Q.11	Q.13	Q.18
Q.21	Q.29	

Redox

Q.3 (C)	Q.4 (F)
---------	---------

Exercise 2

Mole Concept

Q.1	Q.6	Q.10	Q.15
Q.19	Q.23	Q.29	Q.33

Redox

Q.1	Q.8	Q.15
Q.24	Q.25	

Previous Years' Questions

Mole Concept and Redox

Q.1	Q.5	Q.14
-----	-----	------

JEE Advanced/Boards

Exercise 1

Mole Concept

Q.3	Q.7	Q.14
Q.15	Q.22	Q.26
Q.28		

Redox

Q.2	Q.5	Q.13
Q.23	Q.17	

Exercise 2

Mole Concept

Q.2	Q.7	Q.13
Q.16		

Redox

Q.1	Q.6	Q.9
Q.16	Q.19	Q.21

Previous Years' Questions

Mole Concept and Redox

Q.3	Q.14
-----	------

Answer Key

JEE Main/Boards

Exercise 1

Mole Concept

Q.1 (i) 68.125 Kg (ii) 7.63 Kg/m² (iii) 1.72 m

Q.2 7.818×10^{22} atoms

Q.3 4.82×10^{22} atoms

Q.4 0.437

Q.5 6.02×10^{10} g / cm³

Q.6 0.1 M HNO₃

Q.7 14.0 tablets

Q.8 0.1 M

Q.9 (C₃O₄) = C₁₂O₁₆

Q.10 0.44 g

Q.11 35.5: 1, 35.5: 1, 1: 1

Q.12 0.7985, 0.798

Q.13 M₂O₃

Q.14 K₂MnO₄

Q.15 (i) 55.5 M

Q.16 (i) 3.24×10^{-22} g/molecule

(ii) 4.09×10^{22}

Q.17 55.56 moles

Q.18 7.098×10^7 g mol⁻¹

Q.29 2.03 kg

Q.20 10 mol

Q.21 260 mL

Q.22 $\frac{x}{y} = 5$

Q.23 24

Q.24 0.394 g

Q.25 79.714 gm

Q.26 0.744

Q.27 1.125, 1.99, 2.00

Q.28 0.302 M

Q.29 1800 mL

Redox Reaction

Q.1 (i) 5/2 (ii) +2 (iii) +3 (iv) +6 (v) +2 (vi) +6
(vii) +2 (viii) -2 (ix) +5/2 (x) +1 (xi) +3 (xii) +8/3
(xiii) +7 (xiv) 0 (xv) +5 (xvi) +5

Q.2 (i) $5\text{Br}^- + \text{BrO}_3^- + 6\text{H}^+ \longrightarrow 3\text{Br}_2 + 3\text{H}_2\text{O}$,

(iii) $\text{Au} + 2\text{NO}_3^- + 4\text{Cl}^- + 4\text{H}^+ \longrightarrow \text{AuCl}_4^- + 2\text{NO}_2 + 2\text{H}_2\text{O}$

(v) $3\text{MnO}_4^- + 4\text{H}^+ \longrightarrow 2\text{MnO}_4^{1-} + \text{MnO}_2 + 2\text{H}_2\text{O}$

(vii) $5\text{Cl}_2 + \text{I}_2 + 3\text{H}_2\text{O} \longrightarrow 2\text{IO}_3^- + 10\text{Cl}^- + 6\text{H}^+$












(viii) $5\text{Fe}(\text{CN})_6^{4-} + 188\text{H}^+ + 61\text{MnO}_4^- \longrightarrow 5\text{Fe}^{3+} + 30\text{CO}_2 + 30\text{NO}_3^- + 61\text{Mn}^{+2} + 94\text{H}_2\text{O}$

(ix) $6\text{Cu}_3\text{P} + 124\text{H}^+ + 11\text{Cr}_2\text{O}_7^{2-} \longrightarrow 18\text{Cu}^{+2} + 6\text{H}_3\text{PO}_4 + 22\text{Cr}^{+3} + 53\text{H}_2\text{O}$

(ii) $8\text{Cr}_2\text{O}_7^{2-} + 24\text{H}_2\text{S} + 16\text{H}^+ \longrightarrow 8\text{Cr}_2\text{O}_3 + 3\text{S}_8 + 32\text{H}_2\text{O}$

(iv) $3\text{Cu}_2\text{O} + 14\text{H}^+ + 2\text{NO}_3^- \longrightarrow 6\text{Cu}^{+2} + 2\text{NO} + 7\text{H}_2\text{O}$

(vi) $2\text{Cu}^{+2} + \text{SO}_2 + 2\text{H}_2\text{O} \longrightarrow 2\text{Cu}^+ + 4\text{H}^+ + \text{SO}_4^{2-}$



Exercise 2

Mole Concept

Single Correct Choice Type

Q.1 B	Q.2 C	Q.3 B	Q.4 B	Q.5 B	Q.6 B	Q.7 D
Q.8 C	Q.9 A	Q.10 A	Q.11 C	Q.12 A	Q.13 B	Q.14 C
Q.15 A	Q.16 C	Q.17 D	Q.18 A	Q.19 B	Q.20 C	Q.21 A
Q.22 C	Q.23 B	Q.24 C	Q.25 B	Q.26 C	Q.27 D	Q.28 A
Q.29 C						

Redox Reaction

Single Correct Choice Type

Q.1 A	Q.2 D	Q.3 A	Q.4 C	Q.5 C	Q.6 C	Q.7 B
Q.8 C	Q.9 B	Q.10 C	Q.11 D	Q.12 C	Q.13 C	Q.14 B
Q.15 C	Q.16 D	Q.17 C	Q.18 D	Q.19 A	Q.20 A	Q.21 C
Q.22 A	Q.23 D	Q.24 B	Q.25 C	Q.26 A	Q.27 A	Q.28 D
Q.29 C	Q.30 D	Q.31 C	Q.32 C			

Previous Year's Questions

Q.1 A	Q.2 C	Q.3 D	Q.4 B	Q.5 A	Q.6 C	Q.7 A
Q.8 D	Q.9 B	Q.10 B	Q.11 D	Q.12 A	Q.13 C	Q.14 A
Q.15 C	Q.16 D	Q.17 C	Q.18 D	Q.19 A	Q.20 A	Q.21 C
Q.22 D	Q.23 B	Q.24 A				

JEE Advanced/Boards

Exercise 1

Mole Concept

- Q.1 116.8 gm Q.2 9.12 Q.3 5.6 Q.4 0.597 g
Q.5 $\text{BaCl}_2 \cdot 2\text{H}_2\text{O} = 7.038 \text{ g}$, $\text{H}_2\text{O} = 42.962 \text{ g}$ Q.6 120 g Q.7 20.78%
Q.8 $\text{NaOH} = 0.06 \text{ g per } 200 \text{ mL}$, $\text{Na}_2\text{CO}_3 = 0.0265 \text{ g per } 200 \text{ mL}$
Q.9 0.1185 Q.10 45% Q.11 61.5 gm Q.12 320.3 gm
Q.13 6% Q.14 8.097 mL Q.15 6.125 g/litre Q.16 55.53 litre
Q.17 $\text{K}_2\text{CO}_3 = 96\%$, $\text{Li}_2\text{CO}_3 = 4\%$ Q.18 12.9 gm Q.19 1.14 gm
Q.20 SO_4^{2-} ion concentration = 6.528 Q.21 27.27% Q.22 $\text{C}_2\text{H}_6 = 0.66$, $\text{C}_2\text{H}_4 = 0.34$
Q.23 $\text{Pb}(\text{NO}_3)_2 = 3.32 \text{ g}$, $\text{NaNO}_3 = 1.68 \text{ g}$
Q.24 0.0075, $[\text{Pb}^{2+}] = 0.0536 \text{ M}$, $[\text{NO}_3^-] = 0.32 \text{ M}$, $[\text{Cr}^{3+}] = 0.0714 \text{ M}$ Q.25 %NaCl = 77.8%
Q.26 (i) $\text{Fe}_2\text{O}_3 + 2\text{Al} \longrightarrow \text{Al}_2\text{O}_3 + 2\text{Fe}$; (ii) 80: 27; (iii) 10,000 units
Q.27 12.15 gm, $\text{N}_2 = 14.28\%$, $\text{H}_2 = 42.86\%$, $\text{NH}_3 = 42.86\%$
Q.28 0.9413 gram Q.29 (i) 0.5, 0.5; (ii) 0.66, 0.33; (iii) 1, 2 Q.30 13.4 mL
Q.31 $\text{AlCl}_3 = 33.33\%$; $\text{NaHCO}_3 = 50$; $\text{KNO}_3 = 16.67$
Q.32 9.4 gm Q.33 75% Q.34 0.532: 1.00

Redox Reaction

- Q.1 (i) +3 (ii) 0 (iii) +7 (iv) +6 (v) +2 (vi) +3 (vii) +6 (viii) 0 (ix) 0 (x) -1
Q.2 Oxidized: KI, Na_2S , NH_4^+ ; Reduced: CuSO_4 , SO_2 , NO_2^-
Q.3 (i) 8 electrons, (ii) electrons
Q.4 (i) Oxidation number of sulphur in H_2S and SO_2 are respectively -2 and +4.
Q.5 NO_2^- is oxidized to NO_3^- by MnO_4^- (in basic medium) which is reduced to MnO_2 .




Thus, $\text{MnO}_4^- \longrightarrow \text{MnO}_2$ oxidation number decreases by 3-units

$\text{NO}_2^- \longrightarrow \text{NO}_3^-$ oxidation number increases by 2 units

Thus, $2\text{MnO}_4^- \equiv 3\text{NO}_2^-$ $\text{MnO}_4^- \equiv \frac{3}{2}\text{NO}_2^- = 1.5 \text{ mol NO}_2^-$

- Q.6 (i) HSO_3^- ; (ii) NO_2^- ; (ii) Cl^-



Q.8 25 litre

Q.9 0.5 N, 0.167 M





Q.10 0.085 M, 0.042 N

Q.11 54%

Q.12 (i) 85%; (ii) 124.79 mL

Q.13 30.33%

Q.14	(i)	3	2	8	3	1	2	11
	(ii)	1	40	40	12	2	5	
	(iii)	2	27	64	6	2	54	32
	(iv)	3	14	18	14	6	9	



Q.16 0.1716

Q.17 +3

Q.18 $\text{FeC}_2\text{O}_4 = 0.03 \text{ N}$, $\text{FeSO}_4 = 0.03 \text{ N}$

Q.19 Molar ratio = 1: 1

Q.20 40 mL

Q.21 14%

Q.22 57.4%










Q.23 $1.847 \times 10^{-3}\%$






Q.24 $[\text{Fe}^{3+}] = 0.1\text{M}$, $[\text{MnO}_4^-] = 0.105\text{M}$

Q.25 (i) 0.08 N, 0.136%, 0.448 volume; (ii) 1.344

Q.26 0.062 M

Q.27 1.338 g





Exercise 2

Mole Concept

Single Correct Choice Type

Q.1 D

Q.2 B

Q.3 C

Assertion Reasoning Type

Q.4 B

Q.5 A

Q.6 C

Q.7 B

Q.8 A

Q.9 C

Multiple Correct Choice Type

Q.10 A, C, D

Comprehension Type

Paragraph 1: **Q.11** A

Q.12 C

Q.13 A

Q.14 A

Q.15 C

Paragraph 2: **Q.16** (1) C; (2) A; (3) B; (4) A

Paragraph 3: **Q.17** (1) C; (2) A; (3) A

Match the Columns

Q.18 A \rightarrow r; B \rightarrow p; C \rightarrow q

Q.19 A \rightarrow r; B \rightarrow q; C \rightarrow p

Q.20 C

Q.21 B

Redox Reaction

Single Correct Choice Type

Q.1 C

Q.2 D

Q.3 C

Q.4 D

Q.5 C

Q.6 A

Q.7 A

Q.8 B

Comprehension Type

Paragraph 1: **Q.9** D

Q.10 D

Q.11 D

Q.12 D

Paragraph 2: **Q.13** C

Q.14 B, C

Q.15 B

Q.16 A, B, D

Assertion Reasoning Type

Q.17 A

Q.18 D

Q.19 C

Q.20 D

Q.21 B

Q.22 A

Q.23 B

Match the Columns

Q.24 A → w; B → x; C → u; D → p; E → v; F → q; G → r; H → s; I → t

Q.25 A → p, s; B → r; C → p, q; D → p

Previous Year's Questions

Q.1 20%

Q.2 0.437

Q.3 1.7 g

Q.4 A = KO₂

Q.5 10.43

Q.6 2

Q.7 D

Q.8 B

Q.9 5

Q.10 B

Q.11 2

Q.12 4.48 V

Q.13 Moles of Cu²⁺; Moles of C₂O₄²⁻ = 1:2

Q.14 0.9 g, 1.12 g

Q.15 D

Q.16 A, B, D

Q.17 A

Solutions

JEE Main/Boards

Exercise 1

Mole Concept

Sol 1: (i) 125 pound

1 pound = lb = 545 gm

125 pound = 125 × 545 gm

= 125 × 545 × 10⁻³ kg = 68.125 kg

(ii) 14 lb/m²

1 lb = 545 gm

In SI units = 14 × 545 × 10⁻³ kg/m²

= 7.63 kg/m²

(iii) 5'8"

(1' = 12")

5'8" = (12" × 5) + 8" = 68"

= 68 × 2.54 cm = 1.72 m

Sol 2: M_{avg.} = (0.932) 39 + (0.068) 41 = 39.136

Mass of 2g-atoms = 2 × 39.136 gm

Mass of "41K" in 2g-atoms

= 2 × 39.136 × (0.068)

Number of atoms = $\frac{2 \times 39.136}{41} \times (0.068) \times 6.023 \times 10^{23}$

= 7.818 × 10²²

Sol 3: Barium phosphate = Ba₃(PO₄)₂

No. of oxygen atoms = $\frac{6.025}{602.5} \times 8 \times 6.023 \times 10^{23}$

= 4.82 × 10²² atoms

Sol 4: Molecular weight = Vapour density × 2 = 76.6

Let's suppose x % mole of NO₂ is there

76.6 = x (46) + (1 - x) 92

46x = 15.4

x = 0.3347 = 33.47% = mole fraction of NO₂

Total mole = $\frac{100}{76.6} = 1.305$ mole

Mole of NO₂ = (0.3347) × (1.305) = 0.437 mole.

Sol 5: Correction: radius of fluorine

$$= 5 \times 10^{-3} \text{ \AA}$$

$$\text{Mass of nucleus} = 19 \times 1.67 \times 10^{-27} \text{ kg}$$

Volume of nucleus

$$= \frac{4}{3} \times \pi \times (5 \times 10^{-3})^3 \text{ cm}^3$$

Density

$$= \frac{3 \times 19 \times 1.67 \times 10^{-27} \times 10^3}{4 \times \pi \times (5 \times 10^{-3})^3 \times (10^{-10})^3} \frac{\text{gm}}{\text{cm}^3}$$

$$= \frac{95.19 \times 10^{-27} \times 10^{30}}{1570.79 \times 10^{-9}} \text{ gm/cm}^3$$

$$= 6.02 \times 10^{13} \text{ gm/cm}^3$$

Sol 6: mole = $M_1 V_1$

For neutralisation $\text{H}_{\text{HNO}_3} = \text{H}_{\text{NaOH}}$

$$M_1 V_1 = M_2 V_2$$

$$20 \times M_1 = 0.08 \text{ M} \times 25$$

$$M_1 = \frac{0.08 \times 25}{20} = 0.1 \text{ M}$$

Sol 7: HCl produced per day

$$= 3.0 \text{ gm} \times 2.5 = 7.5 \text{ gm} = \frac{7.5}{36.5} \text{ mole}$$

Moles of $\text{Al}(\text{OH})_3$ in an antacid tablet

$$= \frac{400 \times 10^{-3}}{(27 + 51)} = \frac{0.4}{78} \text{ mole}$$

$3 \times \text{mole of } \text{Al}(\text{OH})_3 = \text{moles of HCl}$

$$3 \times n \times \frac{0.4}{78} = \frac{7.5}{36.5}$$

$$n \approx 14$$

Sol 8: $\text{HCl} + \text{AgNO}_3 \rightarrow \text{AgCl} + \text{HNO}_3$

$$M \times 10 \times 10^{-3} = \frac{0.1435}{108 + 35.5}$$

$$M = 0.1 \text{ M}$$

Sol 9: Let's it is C_xO_y

$$x(12) + y(16) = 400$$

$$\frac{x(12)}{400} = 0.36$$

$$x = 12; y = 16$$

$$\text{the formula is } \text{C}_{12}\text{O}_{16} = (\text{C}_3\text{O}_4)_4$$

Sol 10: $\text{C}_x\text{H}_y + \text{O}_2 \rightarrow \text{H}_2\text{O} + \text{CO}_2$

$$0.1 \text{ mole} \quad 0.2 \text{ mole}$$

$$\text{Mole of } \text{H}_2\text{O} = \frac{0.18}{18} = 0.1 \text{ mole}$$

$$\text{Mole of } \text{CO}_2 = 0.02 \text{ mole}$$

$$\text{Mole of } \text{O}_2 \text{ required} = \frac{1}{2}(0.1) + 0.02 = 0.07 \text{ mole}$$

$$y = \text{Mole of H} = (0.1)2 = 0.2 \text{ mole}$$

$$x = \text{Mole of C} = 0.02 \text{ mole}$$

$$\begin{aligned} \text{Mass of hydro is} &= (0.02)12 + (0.2)1 \\ &= 0.24 + 0.2 = 0.44 \text{ gm.} \end{aligned}$$

Sol 11: PCl_3 and PH_3

$$2257\% \quad 91.18\%$$

$$\text{Mass of Cl in } \text{PCl}_3 = 3 \times 35.5 = 106.5$$

$$\text{Mass of H in } \text{PH}_3 = 3 \times 1 = 3$$


$$\left. \begin{aligned} \text{ratio} &= \frac{106.5}{3} = 35.5 \\ \text{Ratio of Cl : H} &= 35.5 \end{aligned} \right\} \text{equal.}$$

Hence prove.

Sol 12: Exp. (I) $\text{CuO} \rightarrow \text{Cu} + \text{X}$

Ratio of mass of

$$\text{Cu : CuO} = \frac{1.098}{1.375} = 0.7985$$



$$\text{Ratio of mass of Cu : CuO} = \frac{1.179}{1.479} = 0.798$$

both ratio are same. Hence prove.

Sol 13: M_xO_y

$$x \times \text{atomic mass of M} = 0.540 \quad \dots (i)$$

$$y \times 16 = 1.020 - 0.540$$

$$y = 0.03$$

Dulong-Petit law (atomic mass of M)

$$\times 0.216 = 5.83$$

$$\text{Atomic mass of M} \approx 27 \quad \dots (ii)$$

$$\text{Petit } x \times (27) = 0.540$$

$$x = 0.02$$

$$\text{Formula of metal oxide} = \text{M}_2\text{O}_3$$

Sol 14: Let's say substance is 100 gm

$$\text{Moles of K} = \frac{39.7}{39} = 1.017$$

$$\text{Moles of Mn} = \frac{29.9}{55} = 0.5436$$

$$\text{Moles of O} = \frac{100 - 39.7 - 29.9}{16} = 1.9$$

so by seeing on ratio of K : Mn : O

empirical formula is K_2MnO_4 .

Sol 15: Molarity = $\frac{\text{no. of moles}}{\text{volume (in litre)}}$

$$\text{No. of moles} = \frac{\text{mass}}{18} = \frac{1000 \times 0.997}{18} \text{ per litre.}$$

$$\text{Molarity} = 55.38 \text{ M}$$

Sol 16: (a) Mass = $(8 \times 12) + (10 \times 1) + (4 \times 14) + (2 \times 16)$
 $= 96 + 10 + 56 + 32$

$$\text{Mass} = 194 \text{ amu}$$

$$= 194 \times 1.66 \times 10^{-24} \text{ gm/molecule}$$

$$= 3.24 \times 10^{-22} \text{ gm/molecule}$$

(b) Molecular mass of $\text{Cl}_2 = 71$

$$\text{Total no. of electrons in one molecule of } \text{Cl}_2 = 34$$

$$\text{So no. of electrons} = \frac{0.142}{71} \times 34 \times 6.023 \times 10^{23}$$

$$= 4.029 \times 10^{22}$$

Sol 17: Molarity = moles per litre

$$= \frac{\text{Mass}}{18 \times V} = \frac{\text{Density} \times \text{Volume}}{18 \times \text{Volume}} = \frac{1000}{18} = 55.55 \text{ M}$$

Sol 18: Volume of plant virus = $(\pi r^2 h)$

$$= \pi \times (75 \times 10^{-10})^2 \times (5000 \times 10^{-10})$$

$$= 8.835 \times 10^{-23} \text{ m}^3$$

$$= 8.835 \times 10^{-23} \times (10^{+2})^3 \text{ cm}^3$$

$$= 8.835 \times 10^{-17} \text{ cm}^3$$

$$\text{Mass} = \frac{8.835 \times 10^{-17}}{0.75} \text{ gm}$$

$$= 11.78 \times 10^{-17} \text{ gm}$$

$$= 11.78 \times 6.023 \times 10^{+23} \times 10^{-17}$$

$$= 7.098 \times 10^7 \text{ g mol}^{-1}$$

Sol 19: 25% of heavy water = 0.5 litre.

Mass of heavy water

$$= 0.5 \times 10^3 \times 1.06 \text{ gm/cm}^3 = 530 \text{ gm}$$

Mass of normal water

$$= 1.5 \times 10^3 \times 1 \text{ gm/cm}^3 = 1500 \text{ gm}$$

$$\text{Total mass} = 2030 \text{ gm} = 2.030 \text{ kg}$$

Sol 20 $\text{SO}_2\text{Cl}_2 + 2\text{H}_2\text{O} \rightarrow \text{H}_2\text{SO}_4 + 2\text{HCl}$

$$\text{Initially} \quad 2.5 \quad 0 \quad 0$$

$$\text{after} \quad 0 \quad 2.5 \quad 5$$

For 2.5 moles of H_2SO_4 , $\text{KOH} = 5$ mole

For 5 mole of HCl , $\text{KOH} = 5$ mole

$$\text{Total KOH} = 5 + 5 = 10 \text{ mole}$$

Sol 21: $\text{NH}_4\text{Cl} + \text{MgCl}_2 + \text{AgNO}_3 \rightarrow$

2% by mass 5% by mass



$$d = 1040 \text{ gm/lit.}$$

Moles of Cl^- in

$$\text{NH}_4\text{Cl} = \frac{2}{(14 + 4 + 35.5)} = 0.0373 \text{ mole}$$

Moles of Cl^- in

$$\text{MgCl}_2 = 2 \times \frac{2}{(24 + 71)} = 0.0421 \text{ mole}$$

Total mole of Cl^-

$$= \text{Total mole of } \text{AgNO}_3 \text{ required}$$

$$= 0.0421 + 0.0373 = 0.07940 \text{ mole}$$

$$\text{Mass of } \text{AgNO}_3 = 0.07940 \times 170 = 13.49 \text{ gm}$$



Mass of solution of

$$\text{AgNO}_3 \times 13.49 \times \frac{100}{5} = 269.97 \text{ gm}$$

$$\text{Volume required} = \frac{269.97}{1.04} \text{ cm}^3 = 259.59 \text{ cm}^3$$

Sol 22: Oxalic acid = $\text{H}_2\text{C}_2\text{O}_4$

Formic acid = HCOOH



Sol 23: $\text{CH}_4 \rightarrow x \text{ mole} \rightarrow \text{Molecular weight} = 16$

$(\text{C}_2\text{H}_4) \rightarrow y \text{ mole} \rightarrow \text{Molecular weight} = 28$

Mean molecular weight

$$= 20 = \frac{x(16) + y(28)}{x + y}$$

$$20 = 16 + \frac{12y}{x + y}$$

$$0.33 = \frac{y}{x + y}$$


$$\frac{x}{x + y} = 0.66$$

$$x : y = 2 : 1$$


$$\text{If } x : y = 1 : 2$$

$$\text{Then, mean molecular weight} = \frac{1(16) + 2(28)}{3} = 24$$

Sol 24:



$$4.369 \times 10^{-3} \quad 6.55 \times 10^{-3}$$



$$3.794 \times 10^{-3} \quad 2.845 \times 10^{-3}$$

Moles of oxygen produced

$$= \frac{146.8 \times 10^{-3}}{22.4} = 6.55 \times 10^{-3} \text{ mole}$$

$$\text{Total mole of KClO}_3 = \frac{1}{39 + 35.5 + 48}$$

$$= 8.163 \times 10^{-3} \text{ mole}$$

Moles of KClO_3 in IInd reaction = 3.794×10^{-3} moles

Moles of KClO_4 produced in IInd reaction

$$= 2.8496 \times 10^{-3} \text{ mole}$$

$$\text{Mass of KClO}_4 = 2.8456 \times 10^{-3} \times (39 + 35.5 + 64)$$

$$= 0.394 \text{ gm}$$

Sol 25: Let's say we have 100 gm mix.




$$x \text{ gram} \quad 0.310 \text{ gram} \quad 0.680x \text{ gram}$$

Initially $\text{FeO} \rightarrow (100 - x) \text{ gram}$

Total $(\text{FeO}) \rightarrow (100 - x + 0.310x) \text{ gm}$

$$= (100 - 0.690x) \text{ gm}$$



$$(105 - 0.690x) \text{ gm}$$

$$2 \times \left(\frac{100 - 0.690x}{72} \right) = \frac{(105 - 0.690x)}{160}$$

$$(100 \times 0.690x) \times \frac{160}{36} = 105 - 0.690x$$

$$4000 - 27.6x = 945 - 6.21 \times 3055 = 21.39x$$

$$x = \text{oxygen external} = 5 \text{ gm} = \frac{5}{32} \text{ mole}$$

So moles of FeO that was present

$$\frac{(100 - 0.690x)}{56 + 16} = \frac{4 \times 5}{32}$$

$$\text{Fe}_3\text{O}_4 + x = 79.71 \text{ gm}$$

$$\text{FeO} = 100 - x = 20.29 \text{ gm}$$

Sol 26: $\text{Zn} + 2\text{I} \rightarrow \text{ZnI}_2$

$$m \quad m$$

$$2x \text{ moles of Zn} = \text{moles of I}$$

(to complete reaction)

$$2 \times \frac{m}{65} = 2x \text{ moles of Zn}$$

$$\frac{m}{127} = \text{moles of I}$$

Since moles of I < 2x moles of Zn

So Zn will be left unreacted

$$\text{Zn unreacted} = \frac{m}{65} - \frac{m}{127 \times 2} \text{ mole}$$

$$\text{Mass Zn unreacted} = m - \frac{65}{254}m = 0.744m$$

$$\text{Sol 27: Mole of P}_4 = \frac{2}{4 \times 31} = \frac{1}{62} \text{ mole}$$

$$\text{Moles of O}_2 = \frac{2}{2 \times 16} = \frac{1}{16} \text{ mole}$$

P_4	+	3O_2	\rightarrow	P_4O_6	
x		3x		0	Initial
0		0		x	After

P_4	+	5O_2	\rightarrow	P_4O_{10}	
y		5y		0	Initial
0		0		y	After

$$\left. \begin{aligned} x + y &= \frac{1}{62} = 0.0161 \\ 3x + 5y &= \frac{1}{16} = 0.0625 \end{aligned} \right\} \text{ by solving}$$

$$y = 7.056 \times 10^{-3}$$

$$x = 9.0435 \times 10^{-3}$$

$$\text{Mass of } P_4O_6 = 9.0435 \times 10^{-3}$$

$$[(4 \times 31) + (6 \times 16)] = 1.9895 \text{ gm}$$


$$\text{Mass of } P_4O_{10} = 7.056 \times 10^{-3}$$

$$[(4 \times 31) + (10 \times 16)] = 2.003 \text{ gm}$$

Sol 28: Moles of aluminium = $\frac{2.7}{27} = 0.1 \text{ mole}$

Moles of H_2SO_4 in solution

$$= \frac{(1.18 \times 100) \times 0.25}{98} = 0.3010 \text{ mole}$$








For consumption of Al, required mole of

$$H_2SO_4 = (0.10) \times \frac{3}{2} = 0.15 \text{ mole}$$

$$\text{remaining mole of } H_2SO_4 = 0.3010 - 0.15 = 0.151 \text{ mole}$$




$$\text{Molarity} = \frac{0.151}{500} \times 1000 = 0.302M$$



$$(0.5)V = \frac{(1.5)}{5} \times 3$$


$$V = 1.8 \text{ lit.} = 1800 \text{ mL}$$

Redox Reactions






$$\therefore \text{Oxidation number of N} = -\frac{5}{2}$$

$$-2x + 5 = 0; x = \frac{5}{2}$$




$$3x - 6 = 0$$



$$x = 2 \quad Mg^{2+}$$




$$\therefore x - 1 = +2 \quad x = +3$$



$$+2 + x - 8 = 0; x = +6$$




$$2 + x - 4 = 0; x = +2$$



$$+ 2 + x - 8 = 0; x = +6$$



$$-4 + 2x = 0; x = +2$$



$$+2 + 4x - 12 = 0$$



$$x = +\frac{5}{2}$$



$$+2x - 2 = 0; x = +1$$



(k) RNO_2

$$+1 + x - 4 = 0; x = 3$$

N^{+3}

(l) Pb_3O_4

$$+3x - 8 = 0$$

$$x = +\frac{8}{3} \quad \text{Pb}^{+8/3}$$

(m) $\text{S}_2\text{O}_8^{2-}$

$$2x - 16 = -2$$

$$2x = 14; x = +7$$

S^{+7}

(n) $\text{C}_6\text{H}_{12}\text{O}_6$

$$+6x + 12 - 12 = 0; x = 0$$

C^0

(o) $\text{Mg}_2\text{P}_2\text{O}_7$

$$+2(2) + 2x - 14 = 0$$



$$x = +5$$

P^{+5}





(p) KClO_3

$$+1 + x - 6 = 0; x = +5$$




Cl^{+5}






$$((1) \times 5) + (2)$$




















$$((1) \times 3) + (2)$$






























$$(1) + (2 \times (2))$$




$$(3 \times (1)) + (2 \times (2))$$









By the oxidation number method,






\therefore To balance the electrons transferred,




To balance charges on both sides,




To balance the electrons transferred to balance oxygen





To balance the number of electrons transferred,











To balance charges on both sides,




We observe that the number of oxygen atoms are simultaneously balanced



To balance charges on both sides,



Sol 5: Disproportionation is a specific type of redox reaction in which a species is simultaneously reduced and oxidized to form two different products.



Sol 6: Ion-electron method :-

(I) Divide the complete equations into two half reactions.

(II) Balance the atoms in each half reaction separately according to the following steps :-

(a) Balance all atoms other than O and H.

(b) For O and H.

1. Acidic Medium:

(i) Add H_2O to the side which is oxygen deficient.

(ii) Add H^+ to the side which is hydrogen deficient.

2. Basic Medium:

(i) Add OH^- to the side which has less -ve charge.

(ii) Add H_2O to the side which is oxygen deficient.

(iii) Add H^+ to the side which is hydrogen deficient.


3. Oxidation State Method: This method is based on the fact that the number of electrons gained during reduction must be equal to the number of e⁻s lost during oxidation.

Sol 7: Definition of Redox Reaction: Reaction which involves change in oxidation state of their atom, generally involve the transfer of electron between species. So, the most essential conditions that must be satisfied is the exchange of electron change in oxidation state.


Sol 8: No, oxidation state term is just introduced to easily calculate the exchange of electron in redox reaction.

So, oxidation no. of an element in a particular compound represents the no. of e⁻s lost or gained by an element during its change from free state into that compound or it represents the extent of oxidation or reduction of an element during its change from free state into that compound.

Sol 9: Redox Couple: Oxidation half reaction and reduction half reaction contributes to redox couple



Sol 10: (1) Combination of half cells (a) and (b)




E.M.F. of the cell, E

$= E_{\text{right}} - E_{\text{left}} = 0.34 - (-0.76)$

$= 1.10 \text{ V}$

(2) Combination of half cells (b) and (d)




E.M.F. of the cell, E



$= E_{\text{right}} - E_{\text{left}} = 0.80 - (+0.34)$

$= 0.46 \text{ V}$






Sol 11: (a)



(i) Ion-electron method :-







(ii) Oxidation number method :-




So, no need to multiply this equation with any co-efficients.

Sol 12 (a) $\text{Fe}^{3+} + e^- \rightarrow \text{Fe}^{2+}$




Sol 13: Oxidation: Increase in oxidation number




Reduction: Decrease in oxidation number



Sol 14: (a) Combination reaction: Reaction in which two or more elements or compounds combine together to form a single compound




(b) Decomposition reaction: Reaction is the separation of a chemical compound into elements or simpler compounds



(c) Displacement reaction:

Reaction in which one element or ion moves out of one compound and into another




Sol 15: Oxidation No.: No. of e⁻s lost or gained by an element during its change from free state into compound or represent the extent of oxidation or reduction of an element during its change from free state into that compound.

Valence: Number of valence bonds a given atom has formed or can form with one or more than one with other atoms.

Sol 16: S in SO_2 has oxidation state +4. It lies between the minimum oxidation state (-2) and maximum oxidation state (+6) of S. Thus, S in SO_2 can show an increase in its ox. no. (i.e., act as reductant) or can show a decrease in its ox. no. (i.e. acts as oxidant). On the other hand in H_2S , S is in - 2 oxidation state and can only increase its oxidation state to act as reductant.

Sol 17: Half Reaction: This is either the oxidation or reduction reaction component of a redox reaction. This is obtained by considering the change in oxidation states of individual substances involved in the redox reaction



oxidation half reaction



Reduction half reaction.

Sol 18: (i) Oxidation-Oxidation is the loss of electrons or an increase in oxidation state by a molecule, atom, or ion.

(ii) Reduction-Reduction is the gain of electrons or a decrease in oxidation state by a molecule, atom, or ion.

(iii) Oxidizing agent an oxidizing agent is a chemical species that removes an electron from another species.

(iv) Reducing agent-Reducing agent is an element or compound that loses an electron to another chemical species in a redox chemical reaction.

Exercise 2

Mole Concept

Single Correct Choice Type

Sol 1: (B) $A + O_2 \rightarrow$

$$M_1V_1 = N_2V_2$$

Equivalent of A = Equivalent of O_2

$$\frac{x}{\text{Equivalent weight of A}} = \left(\frac{16}{16}\right) \times 2$$

$$\frac{x}{2} = \text{equivalent weight of A}$$

Sol 2: (C) Mass O_2 in 88 gm

$$CO_2 = \frac{88 \times 32}{44} = 64 \text{ gm}$$

$$\text{Mole of O} = \frac{64}{16} = 4 \text{ mole}$$

So, mass of CO is = $4 \times (12 + 16) = 112 \text{ gm}$

Sol 3: (B) $Mg + \frac{1}{2}O_2 \rightarrow MgO$

0.25 mole

0.5 mole

$$\text{Mass of MgO} = 0.5 \times (24 + 16) = 20 \text{ gm}$$

Sol 4: (B) Let's diacidic acid is $C_xH_yO_z$

$$\text{Weight of C} = \frac{x(12)}{M}$$

$$\text{Weight} = H = \frac{y}{M}$$


$$\text{Weight of O} = \frac{z(16)}{M}$$

$$x(12) = 8 \times y \Rightarrow 3x = 2y$$

$$x(12) = \frac{1}{2} \times 16(z) \Rightarrow 3x = 2z$$

$$y = z = \frac{3}{2}x$$

\Rightarrow Empirical formula $C_2H_3O_3$



$$\frac{0.5934}{108}$$

$$\text{Mole of salt} = \frac{0.5934}{2 \times 108} = \frac{1}{[216 + (24 + 3 + 48)x]}$$

$$216 + 75x = 364$$

$$x \sim z$$

So the formula would be = $C_4H_6O_6$

Sol 5: (B) $12C(s) + 11H_2(g) + \frac{11}{2}O_2(g) \rightarrow C_{12}H_{22}O_{11}(s)$

$\frac{84}{12}$	$\frac{12}{1}$	$\frac{56}{22.4}$
7	12	2.5


Here O_2 is limiting reagent

$$\text{Moles of } C_{12}H_{22}O_{11} \text{ formed} = \frac{2.5}{11} \times 2 = \frac{5}{11} \text{ mole}$$

$$\text{Mass} = \frac{5}{11} \times [(12 \times 12) + 22 + (11 \times 16)] = 155.45 \text{ gm}$$

Sol 6: (B) $M(CO_3) \rightarrow CO_2 + MO$

or



$$\text{Mass of } CO_3 = 12 + 48 = 60$$

$$\text{Mole of } CO_2 = \frac{12.315}{(PV)} (RT) = \frac{12.315}{1 \times (12.315)} \times 0.0821 \times$$

$$300 = 0.5 \text{ mole}$$

$$\text{Mole of } M(CO_3) \text{ or } M_2CO_3 = 0.5 \text{ mole}$$


$$\text{So, mass of } CO_3 \text{ in carbonate} = 0.5 \times 60 = 30 \text{ gram}$$

Checking all options one by one

(B) is correct.

Sol 7: (D) Empirical formula would be NH_2 because ratio of N and H is given 1 : 2. By stability we see N_2H_4 is correct answer.

Sol 8: (C) $\text{C}_x\text{H}_y + \text{O}_2 \rightarrow \text{CO}_2 + \text{H}_2\text{O}$



By oxygen atom balance $\text{H}_2\text{O} = 10V$

By hydrogen atom balance $y(5) = 2(10) \therefore y = 4$

By carbon balance $= x(5) = 5$

the molecule is CH_4

Sol 9: (A) Molecular weight of $\text{NO}_2 = 32 + 14 = 46$

Molecular weight of $\text{NO} = 16 + 14 = 30$

let's $x = \text{NO}$ $1 - x = \text{NO}_2$

$34 = x(30) + (1 - x) 46$

$16x = 12$

$x = \frac{3}{4}$ so $\text{NO}_2 \% = 25\%$

Sol 10: (A) $5\text{A}_2 + 2\text{B}_4 \rightarrow 2\text{AB}_2 + 4\text{A}_2\text{B}$

4 mole

Molecular mass of $\text{AB}_2 = 250$

Molecular mass of $\text{A}_2\text{B} = 140$

$\text{B}_4 = 480$

$\text{A}_2 = 20$

Moles of AB_2 to be produced $= \frac{1000}{250} = 4$ mole

Moles of A_2B to be produced $= \frac{1000}{140} = 7.14$ mole

So, mass of A_2 , B_4 would be according to AB_2

So, mass of A_2 required $= \frac{5}{2} \times 4 \times 20 = 10 \times 20 = 200$ gm

Mass of B_4 required $= 4 \times 480 = 2 \times 960$ gm $= 1920$ gm

Total mass $= 1920 + 200 = 2120$ gm

Sol 11: (C) $\text{C}_x\text{H}_y\text{O}_z + \text{O}_2 \rightarrow \text{CO}_2 + \text{H}_2\text{O}$

$$\begin{array}{ccc} 132 \text{ gm} & & 54 \text{ gm} \\ \downarrow & & \downarrow \\ 3 \text{ mole} & & 3 \text{ mole} \end{array}$$

$y = 6$ mole

$x = 3$ mole

Sol 12: (A) $\text{Zn} + \frac{1}{2}\text{O}_2 \rightarrow \text{ZnO}$

x gm

$$\frac{1}{2} \times \frac{x}{65} = \frac{v}{22.4}$$

$$v = \frac{x}{65} \times 11.2 = \frac{2x}{65} \times 5.6 \text{ lit.}$$

Sol 13: (B) Let's say 100 gm of clay is given initially

12 gm water

x gm silica

y gm other

After that $(100 - A)$ gm of clay

$(12 - A)$ gm water $\left(\frac{100 - A}{2} \right)$ silica

$$(12 - A) = \frac{(100 - A) \times 7}{100}$$

$$1200 - 100A = 700 - 7A$$

$$93A = 500$$

$$A = \frac{500}{93}$$

By conservation of silica

$$\frac{100 - \frac{500}{93}}{2} = x$$

$$x = 47.31$$

Sol 14: (C) $\text{C}_2\text{H}_4\text{O}_2 + 2\text{O}_2 \rightarrow 2\text{CO}_2 + 2\text{H}_2\text{O}$

x gm 620 - x gm

11 11

$\frac{x}{60}$ mole $\frac{620 - x}{32}$

To produce maximum energy $\text{C}_2\text{H}_4\text{O}_2$ and O_2 will be fully consumed.

$$x \frac{x}{60} = \frac{620 - x}{32} \times \frac{1}{2}$$

$$64x = 37200 - 60x$$

$$x = 300 \text{ gm}$$

$$\text{Weight of } \text{CO}_2 = 2 \times \frac{300 \times 44}{60} = 440 \text{ gm}$$

Sol 15: (A) (Organic compound) + $\text{H}_2\text{O} \rightarrow \text{N}_2$
0.42 gm

$$\text{Moles of N}_2 = \frac{PV}{RT} = \frac{860}{760} \times \frac{100}{11} \times \frac{10^{-3}}{0.08 \times 250}$$

$$= \frac{86}{167200} = 5.143 \times 10^{-4}$$

$$\text{Mass of N}_2 = 5.143 \times 10^{-4} \times 28 = 0.0144 \text{ gm}$$

$$\text{Fraction} = \frac{0.0144}{0.42} = 0.034 = \frac{10}{3} \%$$

Sol 16: (C) Moles of H^+ = (0.1) (0.3) + (0.2) (0.3) \times 2
= 0.15 mole

$$\text{Normality} = \frac{0.15}{500} \times 1000 = 0.3\text{N}$$


Sol 17: (D) Moles of NaOH = (0.300) (0.5) = 0.15 moles

$$\text{For molarity} = 0.2 \text{ M} = \frac{0.15}{V}$$

$$V = 750 \text{ mL}$$

$$\text{Volume to be added} = 750 - 300 = 450 \text{ mL}$$

Sol 18: (A) Moles of water = $\frac{250}{18} = 13.888$ mole



$$\text{Moles urea} = \frac{3}{60} = 0.05 \text{ mole}$$

$$\text{Mole fraction} = 0.0036$$

Sol 19: (B) $\text{P}_4\text{S}_3 + 8\text{O}_2 \rightarrow \text{P}_4\text{O}_{10} + 3\text{SO}_2$

$$\text{Moles of O}_2 = \frac{384}{32} = 12 \text{ mole}$$

$$\text{Moles of P}_4\text{S}_3 = \frac{440}{124 + 96} = 2 \text{ mole}$$

L. R. = O_2

So mass of P_4O_{10} produced

$$= \frac{12}{8} \times [124 + 160] = 426 \text{ gm}$$

Sol 20: (C) $\text{PCl}_5 \rightarrow \text{PCl}_3 + \text{Cl}_2$

Initially 1 mole

$$\text{After } \frac{1}{2} \text{ mole} \quad \frac{1}{2} \text{ mole} \quad \frac{1}{2} \text{ mole}$$

$$\text{Initially } M_{\text{avg.}} = M_{\text{PCl}_5} = 31 + 5(35.5) = 208.5$$

$$\text{After } M_{\text{avg.}} = \frac{M}{3/2} = \frac{208.5}{3} \times 2 = 208.5 \times \frac{2}{3}$$

$$\text{So change in } M_{\text{avg.}} = 33.33\%$$

Sol 21: (A) $3\text{Mg} + 2\text{NH}_3 \rightarrow \text{Mg}_3\text{N}_2 + 3\text{H}_2$

2 mole 2 mole

L. R. = Mg

$$\text{Mass of Mg}_3\text{N}_2 \text{ produced} = \frac{2}{3} \times (72 + 28) = \frac{200}{3} \text{ gm}$$

Sol 22: (C) Let's say solution is in 100 gm.

HCl mole = 1 mole

$$\text{Molality} = \frac{\text{moles of solute}}{\text{mass of solvent (in kg)}}$$

$$\text{Molality} = \frac{1}{(100 - 36.5)} \times 1000 = 15.75$$

Sol 23: (B) Weight of $\text{Na}_2\text{CO}_3 \cdot x\text{H}_2\text{O}$ in 10 mL solution

$$= \frac{0.025}{5} = 0.07$$

$$2 \times \frac{0.07}{46 + 12 + 48 + x(18)} = \frac{9.9}{10} \times 10^{-3}$$

$$\frac{0.14}{106 + 18x} = 0.99 \times 10^{-3}$$

$$0.07 = [104.94 + 17.82x] \times 10^{-3}$$

$$35.06 = 17.82x$$

$$x \sim 2$$

Sol 24: (C) Washing soda (Na_2CO_3) in 25 cc = 0.12 gm

$$2 \times \frac{0.12}{106 + 18x} = 1.7 \times 10^{-3}$$

$$240 = 180.2 + 30.6x$$

$$x \sim 2$$

$$\text{Percentage of carbonate} = \frac{106}{106 + 36} \sim 76\%$$

Sol 25: (B) No. of carbon atoms



$$= \frac{1.2 \times 10^{-3}}{12} \times 6.023 \times 10^{23} = 6.02 \times 10^{19}$$

Sol 26: (C) $21.31 = (0.79)(24) + (0.21 - x)(25) + (x)(26)$

$$24.31 = 18.96 + 5.25 + x$$



$$x = 0.1 = 10\%$$

Sol 27: (D) Using HPh



Using

MeOH



Moles of HCl used in HPh = 4m mole

Moles of HCl used in MeOH = 4.5 m mole

It means that for $\text{NaHCO}_3 \rightarrow \text{H}_2\text{CO}_3$ required mole of HCl 0.5 m mole

Moles of $\text{Na}_2\text{CO}_3 = 0.5$ m mole

$$= 0.5 \times 10^{-3} \times 106 \text{ g/mol} / 250 \text{ mL}$$

$$= 5.3 \text{ mg/25 mL}$$

$$= 2.12 \text{ g/L}$$

NaOH moles = 3.5 m mole

$$\text{Mass of NaOH} = 40 \times 3.5$$

$$= 140 \text{ mg/25 mL}$$

$$= 5.6 \text{ g/L}$$

None of these

Sol 28: (A) Moles of HCl = 0.25×30 m mole = 7.5 m mole

$$\frac{7.5}{2} = 10^{-3} = \frac{x(0.5)}{138} + \frac{(1-x)0.5}{74}$$

$$7.5 \times 10^{-3}$$

Solving this $x \sim 96\%$

$\text{K}_2\text{CO}_3 \sim 96\%$

$\text{Li}_2\text{CO}_3 \sim 4\%$

Sol 29: (C) $\text{KMnO}_4 + 5\text{FeSO}_4 \rightarrow \text{Mn}^{2+} + \text{Fe}^{3+}$

$$\frac{2.0}{152} \text{ moles of KMnO}_4$$

$$= \frac{2.0}{5 \times 152} = 2.631 \times 10^{-3}$$

Volume \times molarity = moles


$$\text{Volume} = \frac{2.631 \times 10^{-3}}{0.05} = 52.63 \text{ mL}$$

Redox Reactions

Single Correct Choice Type


Sol 1: (A) Equivalent weight = Molecular weight $\times n_{\text{factor}}$

$$= (M_0)_{\text{FeSO}_4} \times 1$$




Sol 2: (D) Equivalent weight = Molecular weight $\times n_{\text{factor}}$

$$= (M)_{\text{K}_2\text{Cr}_2\text{O}_7} \times 6$$



Sol 3: (A) $\text{H}_2\text{S} + \text{KMnO}_4 \rightarrow \text{S} + \text{Mn}^{2+}$



$$\frac{m}{34} \times 2 = \frac{1.58}{158} \times 5 = 0.85$$

Sol 4: (C) $\text{HNO}_3 + \text{I}_2 \rightarrow \text{I}_2\text{O}_5 + \text{NO}_2$

$$\frac{m}{63} \times 1 = \frac{127}{127 \times 2} \times 10 = 315$$

Sol 5: (C) $N_1V_1 = N_2V_2$

Oxalic acid KMnO_4

$$10 \times 10^{-3} \times N = 20 \times 10^{-3} \times 0.02 \times (5)$$

$$n\text{-factor} = 0.2N$$

Sol 6: (C) $\frac{m}{34} \times 2 = 10 \times 10^{-3} \times 1$; $m = 0.17$

$$\therefore \text{Purity} = \frac{0.17}{0.2} \times 100 = 85\%$$

Sol 7: (B) $\text{KMnO}_4 + \text{FeC}_2\text{O}_4 \rightarrow \text{Mn}^{2+} + \text{Fe}^{3+} + \text{CO}_2$

$$n_{\text{KMnO}_4} \times 5 = 1 \times (1 + 2) = \frac{3}{5}$$

Sol 8: (C) $\text{M}_2\text{X}_2 + x\text{H}_2 \rightarrow 2\text{M} + x\text{H}_2\text{O}$

1 mole M_2X_2 gives 2 moles M

$$\therefore \frac{3.15 \text{ g}}{(\text{MW})} \rightarrow \frac{1.05}{\text{M}}$$

$$\therefore 6\text{M} = 2\text{M} + x(16)$$

$$\therefore 4\text{M} = x(16)$$

$$\therefore M = 4x$$

$$\text{Now, (EW)}_M = \frac{(\text{MW})_M}{x} = \frac{M}{x}$$

$$\therefore (\text{EW})_M = \frac{M}{x} = 4.$$

Sol 9: (B) Oxidation means increase of oxidation number

\therefore So, there is loss of electrons.

Sol 10: (C) $\text{K}_2\text{Cr}_2\text{O}_7$



$$+2 + 2x - 14 = 0$$

$$x = +6$$

Sol 11: (D) $\text{K}_2\text{Cr}_2\text{O}_7 \rightarrow \text{K}_2\text{Cr}_2\text{O}_4$

$$+6 \rightarrow +3$$

\therefore Change in oxidation No. = [3].

Sol 12: (C) $\text{P} + \text{NaOH} \rightarrow \text{PH}_3 + \text{NaH}_2\text{PO}_2$

P is getting oxidised and also reduced

\therefore It is oxidation and reduction

(D is disproportionation)





Sol 13: (C) CH_2O



$$x + 2 - 2 = 0$$


$$x = 0$$

Sol 14: (B) $\text{CH}_4 \rightarrow \text{C}^{x+} + 4\text{H}^+ \quad x = -4$



Sol 15: (C) Redox: Exchange of electrons

\therefore Change in oxidation state.



No change in oxidation

Sol 16: (D) (a) $+4 - 6 + x = 0; \quad x = +2$

$$(b) +2 + x - 8 = 0; \quad x = +6$$

$$(c) +2x - 2 = 0; \quad x = +1$$

$$(d) x + 5(0) = 0; \quad x = 0$$

Sol 17: (C) (a) $\text{HClO}^{-2} + 1 + x - 2 = 0; \quad x = +1$

$$(b) \text{HClO}_2 + 1 + x - 4 = 0; \quad x = +3$$

$$(c) \text{HClO}_3 + 1 + x - 6 = 0; \quad x = 5$$

$$(d) \text{HClO}_4 + 1 + x - 8 = 0; \quad x = +7$$

Sol 18: (D) $\text{M}^{3+} \rightarrow \text{M}^{6+} + 3\text{e}^-$

Sol 19: (A) $\text{MnO}_4^{x-} + x - 8 = -1; \quad x = +7$


Sol 20: (A) The oxidation number of carbon in CHCl_3 is +4

Sol 21: (C) $\text{Pb}^{2+} \rightarrow \text{Pb}^{4+} + 2\text{e}^-$

Sol 22: (A) $\text{C}_{12}\text{H}_{22}\text{O}_{11}$


$$12x + 22 + 11(-2) = 0; \quad x = 0$$

Sol 23: (D) SO_4^{2-}



$$x - 8 = -2; \quad x = +6$$

Sol 24: (B)



Sol 25: (C) $\text{SO}_3^{2-} \xrightarrow{-2\text{e}^-} \text{SO}_4^{2-}$

Sol 26: (A) $\text{S}_2\text{O}_7^{2-} + 2x - 14 = -2$

$$2x = +12; \quad x = +6$$

Sol 27: (A) $\text{MnO}_2 + 4\text{H}^+ + \text{X} \rightarrow \text{Mn}^{2+} + \text{H}_2\text{O}$



$$8x = 0; \quad x = 0$$

$$\text{Covalency} = 2$$

Sol 28: (D) N_3H

$$3x + 1 = 0; \quad x = -\frac{1}{3}$$

Sol 29: (C) $K_3[Fe(CN)_6]$

$$+3 - (x - 6) = 0; x = +3$$

Sol 30: (D) MH_2

$$+x + 2 = 0; x = -2$$

Sol 31: (C) PH_3^{-3} to $H_3PO_4^{+5}$

Sol 32: (C) In chlorine atom had +5 oxidation state.

$$HClO_3 + 1 + x - 6 = 0; x = 5$$

Previous Years' Questions


Mole Concept

Sol 1: (A) If we consider that $\frac{1}{6}$, in place of $\frac{1}{12}$, mass

of carbon atom is taken to be the relative atomic mass unit, the mass of one mole of a substance will decrease twice.

Sol 2: (C) According to definition of molar solution is one that contains one mole of a solute in one litre of the solution.

Sol 3: (D) $2Al_{(s)} + 6HCl_{(s)} \longrightarrow$



For each mole of HCl reacted, 0.5 mole of H_2 is formed at STP.

1 mole of an ideal gas occupies 22.4 lit at STP.

Volume of H_2 gas formed at STP per mole of HCl reacted is 22.4×0.5 litre.

Sol 4: (B) From molecular formula of magnesium phosphate, it is evident that each mole of $Mg(PO_4)_2$ contains 8 mole of oxygen atoms.

Therefore, 0.25 mole of oxygen atom will remain present in $\frac{0.25}{8}$ mole i.e. 3.125×10^{-2} mole of $Mg(PO_4)_2$.

Sol 5: (A) $200 \text{ mg of } CO_2 = 200 \times 10^{-3} = 0.2 \text{ g}$

$44 \text{ g of } CO_2 = 6 \times 10^{23} \text{ molecules (approx.)}$

$$0.2 \text{ g of } CO_2 = \frac{6 \times 10^{23}}{44} \times 0.2$$

$$= 0.0272 \times 10^{23} = 2.72 \times 10^{21} \text{ molecule}$$

Now, 10^{21} molecule are removed.

$$\text{So remaining molecules} = 2.72 \times 10^{21} - 10^{21}$$

$$= 10^{21}(2.72 - 1) = 1.72 \times 10^{21} \text{ molecule}$$

Now, 6.023×10^{23} molecules = 1 mol

$$1.72 \times 10^{21} \text{ molecules} = \frac{1 \times 1.72 \times 10^{21}}{6.023 \times 10^{23}}$$

$$= 0.285 \times 10^{-2} = 2.85 \times 10^{-3}$$

Sol 6: (C) $K_2Cr_2O_7 + 4H_2SO_4 \rightarrow K_2SO_4 + Cr_2(SO_4)_3$
 $\begin{matrix} +12/\text{two atom} & & & & +6/\text{two atom} \end{matrix}$
Change by 6

$$\text{Eq. wt.} = \frac{\text{Mol. wt.}}{6}$$

$$\begin{aligned} \text{Sol 7: (A)} \quad M_f &= \frac{M_1 V_1 + M_2 V_2}{V_1 + V_2} \\ &= \frac{0.5 \times \frac{3}{4} + 2 \times \frac{1}{4}}{1} = 0.875 \text{ M} \end{aligned}$$

Sol 8: (D) 18 g H_2O contains 2 g H

$\therefore 0.72 \text{ g } H_2O \text{ contains } 0.08 \text{ g H.}$

44 g CO_2 contains 12 g C

$\therefore 3.08 \text{ g } CO_2 \text{ contains } 0.84 \text{ g C}$

$$\therefore C : H = \frac{0.84}{12} : \frac{0.08}{1} = 0.07 : 0.08 = 7 : 8$$

\therefore Empirical formula = C_7H_8

Sol 9: (B) $M_{0.98}O$

Consider one mole of the oxide.

Moles of M = 0.98, Moles of $O^{2-} = 1$

Let moles of $M^{3+} = x$

$$\Rightarrow \text{Mole of } M^{2+} = 0.98 - x$$

\Rightarrow Doing charge balance

$$(0.98 - x) \times 2 + 3x - 2 = 0$$

$$\Rightarrow 1.96 - 2x + 3x - 2 = 0$$

$$\Rightarrow x = 0.04$$

$$\Rightarrow \% \text{ of } M^{3+} = \frac{0.04}{0.98} \times 100 = 4.08\%$$

Sol 10: (B) Let the mass of $O_2 = x$


Mass of $N_2 = 4x$

Number of moles of $O_2 = \frac{x}{32}$

Number of moles of $N_2 = \frac{4x}{28} = \frac{x}{7}$

\therefore Ratio = $\frac{x}{32} : \frac{x}{7} = 7 : 32$

Sol 11: (D) 1 g of $C_8H_7SO_3Na = \frac{1}{206}$ mole



Sol 12: (A) $C_xH_y(g) + \left(x + \frac{y}{4}\right)O_2(g) \rightarrow xCO_2(g) + \frac{y}{2}H_2O(l)$


15 mL

Volume of O_2 used = $\frac{20}{100} \times 375 = 75$ mL

Volume of air remaining = 300 mL

Total volume of gas left after combustion = 330 mL

Volume of CO_2 gases after combustion = 330 - 300 = 30 mL



$$\frac{x}{1} = \frac{30}{15} \Rightarrow x = 2$$


$$\frac{x + \frac{y}{4}}{1} = \frac{75}{15} \Rightarrow x + \frac{y}{4} = 5$$

$$\Rightarrow y = 12$$

$$\Rightarrow C_2H_{12}$$

Redox Reactions



Sol 13: (C) Prevent action of water and salt.




Sol 15: (C) CH_2O

$$x + 2 - 2 = 0; x = 0$$


Sol 16: (D) $K_2Cr_2O_7 + 6KI + 7H_2SO_4 \longrightarrow$



Sol 17: (C) Number of e^- transferred in each case is 1, 3, 4, 5.




Sol 19: (A) $6MnO_4^- + I^- + 6OH^- \longrightarrow$




Sol 20: (A) $H-O-\overset{\overset{H}{|}}{\underset{\underset{O}{|}}{P}}-OH$, hence it is dibasic. It acts as a reducing agent also.

Sol 21: (C)







\therefore Balanced Equation:



So, $x = 2$, $y = 5$ & $z = 16$.

Sol 22: (D) The reducing agent oxidises itself:



Note: Powers of 'O' are oxidation number of 'O' in the compound.

Sol 23: (B)


The complex $[\text{CoCl}(\text{NH}_3)_5]^+$ decomposes under acidic medium, so $[\text{CoCl}(\text{NH}_3)_5]^+ + 5\text{H}^+ \rightarrow \text{Co}^{2+} + 5\text{NH}_4^+ + \text{Cl}^-$

Sol 24: (A) H_2O_2 can undergo reduction as well as oxidation because oxidation number of oxygen in H_2O_2 is -1. So, it can act both as reducing agent and oxidising agent.

JEE Advanced/Boards

Exercise 1

Mole Concept



69.6 gm

$$69.6 \text{ gm of MnO}_2 = \frac{69.6}{87} \text{ mole}$$

$$\text{Mole of HCl} = \frac{69.6}{87} \times 4 \text{ mole}$$

$$\text{Weight of HCl} = \frac{69.6}{87} \times 4 \times 35.5 = 116 \text{ gm}$$



4.32 gm 5.76 gm 6.82 gm

0.054 mole 0.48 mole 0.0960 mole




L. R. = Cl_2

So TiCl_4 mole produced

$$= \frac{1}{2} \times 0.0960 = 0.048 \text{ mole}$$

$$\text{Weight of TiCl}_4 \text{ produced} = 0.048 \times 190 = 9.12 \text{ gm}$$




5.6 moles 4.8 moles

L. R. = SO_2

So H_2SO_4 mole obtained in maximum = 5.6 mole



$$\text{Pure Na}_2\text{CO}_3 = (0.95)x \text{ gm}$$



$$\text{Mole of acid} = (45.6 \text{ mL}) \times (0.235) = 10.716 \text{ m mole}$$

$$\text{Moles of Na}_2\text{CO}_3 \text{ required} = 5.358 \text{ m mole}$$

$$\text{Weight of Na}_2\text{CO}_3 \text{ required} = (0.95) \times 5.358 (106) \times 10^{-3}$$

$$x = 0.597 \text{ gm}$$




$$\text{Molecular weight of BaCl}_2 \cdot 2\text{H}_2\text{O} = 208 + 36 = 244$$

$$\text{BaCl}_2 = 6 \text{ gm}$$


$$\text{BaCl}_2 \cdot 2\text{H}_2\text{O} = 6 \times \frac{244}{208} = 7.038 \text{ gm}$$

$$\text{H}_2\text{O} = 42.962 \text{ gm.}$$




$$\text{HCl mole} = 5 \text{ mole}$$

$$\text{FeCl}_3 \text{ mole} = 1.5 \text{ mole (acidic)}$$



After this reaction NaOH left = 5 mole




$$1.5$$



After this reaction NaOH left = $5 - (1.5)3 = 0.5 \text{ mole}$

$$\text{Volume after reaction} = 15 + 5 + 50 = 70 \text{ litre}$$

$$\text{Normality} = \frac{0.5}{70} = 7.142 \times 10^{-3} \text{ N}$$




$$\text{Weight of Fe}_2\text{O}_3 = \frac{1.5}{2} \times 160 = 120 \text{ gm}$$




$$\text{Mole of NaOH} = (26.7) \times (0.4) \text{ m mole} = 10.68 \text{ m mole}$$

$$\text{Mole of H}_2\text{SO}_4 = 5.34 \text{ m mole}$$

$$\text{Weight of H}_2\text{SO}_4 = 0.523 \text{ gm}$$



$$x \text{ gram} \quad (0.5 - x) \text{ gm}$$





$$= \frac{(0.5 - x)}{80} \times 98 \text{ gm}$$


$$\text{Total H}_2\text{SO}_4 = x + \frac{(0.5 - x)98}{80} \times 0.523$$

$$x = \frac{0.0895}{0.225} \sim 0.3977 \text{ gm}$$

$$\% \text{SO}_3 = \frac{0.5 - 0.3977}{0.5} \sim 20.4\%$$



after MeOH:



Mole of HCl (when HPh) = 1.75 m mole

Mole of HCl (when MeOH) = 0.25 m mole (extra added)

Amount of NaHCO_3 = 0.25 m mole

Amount of HCl required in (2) and (3)


$$= (0.25)_2 = 0.5 \text{ m mole}$$

Amount of Na_2CO_3 = 0.25 m mole

Amount of NaOH = $1.75 - (0.25) = 1.5$ m mole

NaOH (in gram) = $1.5 \times 10^{-3} \times 40 = 0.06$ gm per 200 mL


Na_2CO_3 (in gram) = $0.25 \times 10^{-3} \times 106$
= 0.0265 gm/200 mL



0.158 mole 0.1 mole

L. R = KO_2

Moles of O_2 formed = $\frac{3}{4} \times 0.158 = 0.1185$




0.959 gm

Moles of CaO = 0.017125 mole

Moles of CaCl_2 = 0.017125 mole

Mass of CaCl_2 = $(0.017125) \times 111$
= 1.9 gm

% of CaCl_2 = $\frac{1.9}{4.22} = 45\%$



100 gm

Moles of cyclohexanol = $\frac{100}{100} = 1$ mole

Mole of cyclohexene = 0.75 mole



Mass of cyclohexene = $(0.75) \times 89 = 66.75$ gm



Pure NaCl mole = $\frac{(0.95)250}{23 + 35.5} = 4.059$ mole

Pure Na_2SO_4 = $\frac{4.059}{2} \times (46 + 96) = 288.24$ gm

Na_2SO_4 (90% pure) = $\frac{288.24 \times 100}{90} = 320.27$ gm



AgCl formed = $\frac{x}{188} \times (1435)$ gm = $0.763 x$

Total weight after reaction = $0.4066 - x + 0.763 x$


Weight lost = $(1 - 0.763)x = 0.0725$

$$x = 0.306 \text{ gm} = 30.6\%$$

Weight of Cl in initial mixture

$$= (0.4066 - 0.306) \times \frac{35.5}{143.5} = 0.0248 \text{ gm}$$

$$\% \text{ of Cl} = \frac{0.0248}{0.4066} = 6.1\%$$



Moles of CaCO_3 = moles of H_2SO_4

required = $\frac{0.5}{63.5 + 60} = 4.048 \times 10^{-3}$ mole

m litre of 0.5M H_2SO_4 required $\frac{4.048}{0.5} = \text{mL} = 8.096$ mL



Moles of NaOH = $15 \times \frac{1}{10} = 1.5$ m mole


Moles of H_2SO_4 required = $\frac{1.5}{2} = 0.75$ m mole

In 12 mL, mole of H_2SO_4 = 0.75 m mole

In 1 L, mole of H_2SO_4 = $\frac{0.75}{12}$ mole

In 1 L, weight of H_2SO_4 required

$$= \frac{0.75}{12} \times 98 \text{ gram} = 6.125 \text{ gm/L}$$



XV

Let's say volume of ethane = x l

$$\text{Weight of } C_4H_{10} = \frac{x}{2 \times 22.4} \times \frac{90}{100} \times \frac{85}{100} \times 58 = 55.53 \text{ l}$$

$$x = 55.53 \text{ l}$$

Sol 17: Mole of HCl = 30×0.25 m mole = 7.5 m mole

let's say x fraction is K_2CO_3 so

$$\frac{7.5}{2} \times 10^{-3} = \frac{x(0.5)}{138} + \frac{(1-x)(0.5)}{74}$$




$$x \sim 96\%$$

Sol 18: Mass of solution of HCl

$$= 100 \times 1.18 \text{ gm} = 118 \text{ gm}$$


$$\text{Mass of HCl in solution} = (0.36) (118) = 42.48 \text{ gm}$$

$$n_{HCl} = \text{mole of HCl} = \frac{42.48}{365} = 1.163 \text{ mole}$$



Mass prepared of $NaClO_3$

$$= 0.1212 \times \text{molecular weight} = 12.911 \text{ gm}$$



$$\text{Mole of } Mg_2P_2O_7 = \frac{1.054}{224} = 4.747 \times 10^{-3}$$

$$\text{Weight of } NaH_2PO_4 = 2 \times 4.747 \times 10^{-3} \times 119.98$$

$$(\text{Molecular weight}) = 1.139 \text{ gm}$$

Sol 20: Moles of HNO_3 = 8×5 m mole = 40 m mole

Mole of HCl = 4.8×5 m mole = 24 m mole

Let's say volume of H_2SO_4 is V mL

So mole of H_2SO_4 = 17 V m mole

Moles of HNO_3 in 30 mL (picked up from 2 l sol)

$$= \frac{40}{2000} \times 30 = \frac{120}{200} \text{ m mole}$$

Moles of HCl in 30 mL (picked up from 2 l sol)

$$= \frac{24}{2000} \times 30 = \frac{72}{200} \text{ m mole}$$

Moles of H_2SO_4 is 30 mL (picked up from 2 lt. sol)

$$= \frac{17V}{2000} \times 30 = \frac{51V}{200} \text{ m mole}$$

$$\text{Total moles of } H^+ \text{ from 30 mL solution} = \frac{120}{200} + \frac{72}{200} + \frac{102V}{200}$$

$$= \left(\frac{192 + 102V}{200} \right) \text{ m mole}$$

$$\text{Mole } Na_2CO_3 \cdot 10H_2O = \frac{1}{286} \text{ mole}$$

$$\text{Mole of OH} = \frac{2}{286} \text{ mole (in 100 mL)}$$

$$\text{Mole of OH in 42.9 mL} = \frac{2 \times 0.429}{2.86}$$






$$= \frac{0.858}{286} = 0.003 \text{ mole}$$

$$10^{-3} \times \left(\frac{192 + 102V}{200} \right) = 0.003 ; V = 4 \text{ mL}$$

$$\text{Amount of sulphate ion in gm} = \frac{51 \times 4}{200} \times (96) \times 10^{-3}$$


$$= 0.09792 \text{ gm/30 mL}$$

$$= 6.528 \text{ gm/L}$$



initially 10 Meq. x mole

after (10 - x) m mole





$$\frac{x}{2} + \frac{60-x}{2} = \frac{12}{2}$$

$$x = 27.27\%$$

Sol 22: $PV = n_T RT$

$$(1) (40) = n_T (0.0821) (400)$$

$$n_T = \text{Total mole} = 1.2180$$





Mole of O_2 required

$$= \frac{7}{2} x + 3(1.218 - x) = \frac{130}{32}$$


$$x = 0.817 \text{ mole}$$

$$\text{Mole fraction of } C_2H_4 = \frac{1.218 - 0.817}{1.218} = 0.33$$

$$\text{Mole fraction of } C_2H_6 = 0.67$$



$$\left[\frac{(5-x)}{85} \right] + \frac{1}{4} \left(\frac{5-x}{85} \right) + 2(x)$$



$$L. R. = CrSO_4$$

So moles of $PbSO_4$ formed

$$= 2.5 \text{ m mole} \times 3$$



$$= 7.5 \text{ m mole}$$

$$\text{Molar conc. of } [Pb^{2+}] = \frac{11.25 - 7.5}{70} = 0.0536 \text{ M}$$

$$\text{Molar conc. of } [NO_3^-] = \frac{(2 \times 11.25)}{70} = 0.32 \text{ M}$$

$$[Cr^{3+}] = \frac{2 \times 2.5}{70} = 0.0714 \text{ M}$$

Sol 25: $NaCl$




$$\text{Mole of } CaO = \frac{1.12}{56} = 0.02 \text{ mole}$$

$$\text{Moles of } CaCl_2 = 0.02 \text{ mole}$$

$$\text{Weight of } CaCl_2 = 2.22 \text{ gm}$$

$$NaCl = 10 - 2.22 = 7.78 \text{ gm}$$

$$\% NaCl = 77.8\%$$



(ii) Mole ratio (to complete reaction) = 1 : 2



$$\text{mass ratio} = 1 \times (112 + 48) : (2 \times 27) = 80 : 27$$

$$(iii) 2.7 \text{ kg of } Al = \frac{2700}{27} \text{ mole} = 100 \text{ mole}$$

$$16 \text{ kg of } Fe_2O_3 = \frac{16000}{160} \text{ mole} = 100 \text{ mole}$$

$$L. R. = Al$$

$$\text{So energy released} = 200 \times \frac{100}{2} = 10000 \text{ unit}$$



$$\text{Initially } 1 \quad 3a$$

$$\text{after } 1-x \quad 3-3x \quad 2x$$

$$P(\text{Molecular weight}) = SRT$$

$$1(M.W.) = (0.497) \times (0.0821) (298)$$

$$\text{Molecular weight} = 12.15 \text{ gm}$$

$$\frac{(2x)(17) + (3-3x)2 + (1-x)28}{4-2x} = 12.15$$

$$34x + 6 - 6x + 28 - 28x = 48.63 - 24.31x$$

$$24.31x = 14.63$$

$$x = 0.602$$

% composition by volume

$$N_2 = \frac{1-0.602}{4-2(0.602)} = \frac{0.398}{2.795} = 14.21\%$$

$$H_2 = 3(N_2\%) = 42.86\%$$


$$NH_3 = \frac{2(0.602)}{2.795} = 42.86\%$$

We know average molecular weight = 12.15

$$\text{So } (1) (12.15)$$

$$= \frac{\text{Mass}}{22.4} \times (0.0821) \times (273)$$

$$\text{Mass} = 12.14 \text{ gm.}$$





$$\text{Volume of film} = 6 \times 10^{-10} \times 300 \times 1 \times 3 \text{ m}^3$$

$$= 54 \times 10^{-8} \text{ m}^3 = 0.54 \text{ cm}^3$$

$$\text{Mass of the film} = 0.54 \text{ gm}$$

$$\text{Mole of } [(\text{CH}_3)_2\text{SiO}]_n = \frac{0.54}{n[30 + 28 + 16]}$$

$$\text{Mass of } [(\text{CH}_3)_2\text{SiCl}_2] = \frac{0.54}{71} \times (58 + 71) = 0.941 \text{ gm.}$$



$$5x = y$$



$$3(1-x) = 4-y$$

$$3-3x = 4-5x$$

$$x = \frac{1}{2}$$

$$y = \frac{5}{2}$$

$$\text{P}_4\text{O}_6 = \text{P}_4\text{O}_{10} = 50\%$$





$$5x = y$$

$$3(3-x) = 11-y$$

$$9-3x = 11-5x$$

$$x = 1$$

$$\text{P}_4\text{O}_{10} = \frac{1}{3}; \quad \text{P}_4\text{O}_6 = \frac{2}{3}$$



$$5x = y$$

$$3(3-x) = 13-y$$

$$9-3x = 13-5x$$

$$x = 2$$

$$\text{P}_4\text{O}_6 = 2$$

$$\text{P}_4\text{O}_{10} = 1$$



Let's say V mL must be added

$$\text{Weight of solution} = (1.04 V) \text{ gm}$$

$$\text{Weight of AgNO}_3 = 0.05 \times (1.04 V) \text{ gm}$$

$$\text{Moles of AgNO}_3 = \frac{(0.05)(1.04V)}{173}$$

Minimum moles of Cl^- (it will be case of more molecular weight i. e. KCl)

$$= \frac{0.3}{39+35.5} = \frac{0.3}{74.5}$$

$$\frac{(0.05)(1.04V)}{173} = \frac{0.3}{74.5}$$

$$V = 13.4 \text{ mL.}$$

Sol 31: In 500 mL of NaOH

$$\text{Weight of solution} = 1.8 \times 500 = 900 \text{ gm}$$

$$\text{So, weight of NaOH} = (0.08) (900) = 72 \text{ gm}$$

$$\text{Mole of NaOH} = \frac{72}{40} = 1.8 \text{ mole}$$

$$\text{Moles of H}^+ = 1.8 \text{ mole}$$



$$\text{On C-balance } n_{\text{CO}_2} = n_{\text{NaHCO}_3}$$


$$\text{Mass of NaHCO}_3 = \frac{18.6}{44} \times (84) = 33.50 \text{ gm}$$

$$\text{H}^+ = 1.8 = \frac{18.6}{44} + 3 \left[\frac{x}{27 + 3(35.5)} \right] + 0$$

$$1.8 = \frac{x}{44.5} + 0.418 \Rightarrow x = 61.5 \text{ gm} = \text{mass of AlCl}_3$$

$$\text{Mass of KNO}_3 = 124 - 97 = 27 \text{ gm}$$

$$\text{Total mole} = 0.267 + 0.460 + 0.422 = 1.149 \text{ mole}$$



$$\text{Mole of } \text{CHCl}_3 = \frac{30}{119.5}$$

By carbon balance

Mole of acetone (ideally)

$$= \frac{1}{2} \times \frac{30}{119.5}$$

As the yield is 75%

So, weight required

$$= \frac{30}{2 \times 119.5} \times \frac{100}{75} \times (58) = 9.7 \text{ gm}$$

Sol 33: $\text{Cu}_2\text{O} + x$

Let's assume total 100 gm is given

$\text{Cu} = 66.67 \text{ gm}$

$$0 = \frac{66.67}{63.5} \times \frac{1}{2} \times 16$$

Oxygen (O) = 8.4

% $\text{Cu}_2\text{O} = 66.67 + 8.4 = 75\%$

Sol 34: $\text{Hg} + \text{I}_2 \rightarrow \text{HgI}_2$

$$\left(\frac{M}{200} - x \right) \left(\frac{M}{254} - \frac{x}{2} \right)$$



$$x \text{ mole } \frac{x}{2}$$

Let's say M gm is initially taken

$$\frac{M}{200} - x = \frac{M}{254} - \frac{x}{2}$$

$$M \left(\frac{54}{200} \right) \frac{1}{254} = \frac{x}{2}$$

$$M = \left(\frac{254}{0.54} \right) x$$

$$(\text{gm}) \text{Hg}_2\text{I}_2 = \frac{x}{2} \times (200 + 127) \times 2 = 327 x \text{ HgI}_2(\text{gm})$$

$$= \left(\frac{M}{200} - x \right) = \left[\frac{254}{(0.54)(200)} - 1 \right]$$

x Molecular weight = $(1.351 \times 454) \times \text{HgI}_2$

$\text{HgI}_2 : \text{Hg}_2\text{I}_2 = 0.532 : 1.$

Redox Reactions

Sol 1: (a) NaNO_2

$$\text{Na}^+ \text{N}^{x+} 2\text{O}_2^- + 1 + x - 4 = 0; x = +3$$

(b) H_2

$$2x = 0; x = 0$$

(c) Cl_2O_7

$$2x - 14 = 0; x = +7$$

(d) KCrO_3Cl

$$\text{K}^+ \text{Cr}^{x+} 3\text{O}_2^- \text{Cl}^-; +1 + x - 6 - 1 = 0; x = +6$$

(e) BaCl_2

$$+x - 2 = 0; x = +2$$

(f) ICl_3

$$+x - 3 = 0; x = +3$$

(g) $\text{K}_2\text{Cr}_2\text{O}_7$

$$+2 + 2x - 14 = 0; x = +6$$


(h) CH_2O

$$+x + 2 - 2 = 0; x = 0$$




(i) $\text{Ni}(\text{CO})_4$

$$+x + 0 = 0; x = 0$$

(j) NH_2OH



$$+3 + x - 2 = 0; x = -1$$



Sol 3: (a) $3\text{Fe} + 4\text{H}_2\text{O} \longrightarrow \text{Fe}_3\text{O}_4 + 4\text{H}_2 = 8 \text{ electrons}$

(b) $\text{AlCl}_3 + 3\text{K} \longrightarrow \text{Al} + 3\text{KCl} = 3 \text{ electrons}$

Sol 4: (a) Sulphur shows various oxidation states such as -2, 0, +2, +4, +6


In H_2S , oxidation no. of S is '-2'

So now it can only act as a reducing agent because it can't get more electrons since S^{2-} is in its lowest oxidation state.

But in SO_2 , oxidation state of S is +4 which is an intermediate oxidation state. So, it can gain or lose electrons and can go to -2, 0, +2 or higher +6 oxidation state.


(b) Oxidation state of oxygen in H_2O_2 is '-1' so it can get oxidised or reduced because O have multiple oxidation state like -2, -1, 0. By losing electrons it can form O_2 and act as a reducing agent and by gaining e^- s, it can form H_2O and behave as an oxidising agent.

Sol 5: NO_2^- is oxidized to NO_3^- by MnO_4^- (in basic medium) which is reduced to MnO_2



Thus, $\text{MnO}_4^- \longrightarrow \text{MnO}_2$ oxidation number decreases by 3-units

$\text{NO}_2^- \longrightarrow \text{NO}_3^-$ oxidation number increases by 2 units



Sol 6: (a) HSO_3^- (b) NO_2^- (c) Cl^-

Sol 7: (a) $4\text{Zn} + 10\text{HNO}_3 \longrightarrow 4\text{Zn}(\text{NO}_3)_2 + \text{N}_2\text{O} + 5\text{H}_2\text{O}$

(b) $6\text{HI} + 2\text{HNO}_3 \longrightarrow 3\text{I}_2 + 2\text{NO} + 4\text{H}_2\text{O}$

Sol 8: 5 moles of H_2SO_4 can produce

1 mole of H_2S

$$0.2 \times V \times 10^{-3} = n\text{H}_2\text{SO}_4$$

(equating equivalents)

\therefore Volume = 25 lit.

Sol 9: $\therefore 20 \times 0.2 \times 2 = 0.167 \text{ M}$

$$\text{Normality} = n_f \times M = 0.5 \text{ N}$$

Sol 10: mole of $\text{As}_2\text{O}_3 = 5.54 \times 10^{-4}$

equating equivalents,


$$(5.54 \times 10^{-4}) \times (2) = (26.1 \times 10^{-3}) \times M \times 5$$

$$\therefore \text{Molarity} = 8.49 \times 10^{-3},$$

$$\text{Normality} = \text{molarity} \times n\text{-factor}$$

$$= (8.49 \times 10^{-3}) \times (5) = 4.24 \times 10^{-2}$$

Sol 11: $\text{CaO} \longrightarrow \text{CaC}_2\text{O}_4$



Equating equivalents

$$(\text{equivalent})_{\text{CaC}_2\text{O}_4} = (\text{equivalent})_{\text{KMnO}_4}$$

$$n_{\text{CaC}_2\text{O}_4} \times 2 = 40 \times 0.25 \times 10^{-3}$$

$$\text{Moles of CaC}_2\text{O}_4 = 5 \times 10^{-3}$$


$$\therefore \text{Mole of CaO} = \text{CaC}_2\text{O}_4 = 5 \times 10^{-3}$$

$$\therefore \text{Mass of CaO} = 0.28$$

\therefore % composition

$$= \frac{0.28}{0.518} \times 100 = 54\%$$

Sol 12: Reaction



Assume mass of $\text{H}_2\text{O}_2 = x \text{ gm}$

\therefore Molarity of

$$\text{H}_2\text{O}_2 = \frac{\frac{x}{34} \times 1000}{20} = 147 \times \text{M}$$

$$\text{Moles of KMnO}_4 = \frac{0.316}{158} = 2 \times 10^{-3}$$

Now equating equivalents,

$$\text{Equivalents of H}_2\text{O}_2 = \text{Equivalents of KMnO}_4$$

$$1.47x \times 20 \times 10^{-3} \times 2 = 2 \times 10^{-3} \times 5$$

$$\therefore x = 0.17 \text{ gm}$$

$$\therefore \text{Purity of H}_2\text{O}_2 = 85\% \quad (\text{i})$$

moles O_2 evolved = moles of H_2O_2 consumed.

$$\therefore \text{Moles of O}_2 = 5 \times 10^{-3}$$

$$\therefore \text{Volume} = \frac{nRT}{P} = 124.8 \text{ ml} \quad (\text{ii})$$



Sol 13: $(\text{CaOCl}) + \text{Cl}^-$ 5.7 gm is taken

\therefore Lets take moles of $\text{CaOCl}_2 = x$

molarity of

$$\text{CaOCl}_2 = \frac{x}{500} \times 1000 = 2x$$

Now on treatment with KI + HCl



\therefore Equivalents of I_2 = equivalents of $\text{Na}_2\text{S}_2\text{O}_3$

$$= 24.35 \times \frac{1}{10} \times 10^{-3} = 2.435 \text{ milliequ.}$$






Now, equiv. of I_2 = equiv. of Bleaching powder

$$2.435 \times 10^{-3} = 2x \times 25 \times 10^{-3}$$

$$\therefore x = 4.87 \times 10^{-2}$$


\therefore Mass of bleaching powder = 1.73 gm

$$\therefore \% \text{ availability} = \frac{1.73}{5.7} \times 100 = 30.33\%$$





Sol 15:



(i)



(ii)



Sol 16:



Say molarity of NO_3^- ions = x M

$$\therefore \text{Moles of } \text{NO}_3^- = x \times 25 \times 10^{-3}$$

Equivalents of NO_3^-

$$= \text{Equivalents of } \text{NH}_3 = 8 \times x + 25 \times 10^{-3} = 0.2x$$

$$\therefore \text{Moles of } \text{NH}_3 = 0.2x$$


$$\text{Moles of NaOH} = 32.10 \times 10^{-3} \times 0.1 = 3.21 \times 10^{-3}$$

Now, moles of HCl = (moles of NH_3) + (moles of NaOH)

$$50 \times 0.15 \times 10^{-3} = 0.2x + 3.21 \times 10^{-3}$$

$$x = 2.145 \times 10^{-2}$$

$$\therefore \text{Molarity} = 8x = 0.1716 \text{ M}$$



$$\therefore \text{Moles of } \text{KReO}_4 = 9.28 \times 10^{-5}$$

$$\therefore \text{Moles of } \text{KMnO}_4 = 0.05 \times 11.45 \times 10^{-3} \times 5$$

Now equating equivalents of

$$\text{Re}^{x+} = \text{KMnO}_4$$

$$(x_{\text{Re}^{x+}})(7-x) = 5.725 \times 10^{-4} \quad \dots\dots (1)$$

Now equating equivalents of


$$\text{KReO}_4 = \text{Re}^{x+}$$

$$9.28 \times 10^{-5} = \text{Re}^{x+} = x_{\text{Re}^{x+}}$$

$$\therefore 7-x = \frac{5.725 \times 10^{-4}}{9.28 \times 10^{-5}}; x = +1$$

Sol 18: Let moles of $\text{FeC}_2\text{O}_4 = x$

$\text{FeSO}_4 = y$



$$\text{Now, } (2x + x + y) 0.1 = 60 \times 0.02 \times 5$$

$$3x + y = 60$$

$$x + y = 40 \times 0.02 \times 5 = 4$$

$$\therefore 4 - x = 6 - 3x$$

$$2x = 2$$

$$x = 1$$

$$y = 3$$

$$\therefore \text{Normality} = 1 \times 3 \times 10^{-2} = 0.03\text{N of FeC}_2\text{O}_4$$


$$= 3 \times 10^{-2} = 0.03 \text{ M of FeSO}_4$$

Sol 19: Mass of KCl = x gm

$$\text{H}_2\text{O} = 1 - x - y \text{ gm}$$

$$\text{KClO}_3 = y \text{ gm}$$


Treating with SO_2




Then silver chloride formed

$$\therefore \text{Total moles of chloride} = 10^{-3} = \frac{x}{74.5} + \frac{y}{122.5}$$

Now for another experiment



$$\therefore \text{Equivalents reacted} = 3 \times 10^{-3}$$



Equivalents of Fe^{2+} = Equivalent of ClO_3^-

$$3 \times 10^{-3}$$

$$= \frac{y}{122.5} \times 10^3 \times 25 \times 10^{-3} \times 6$$

$$y = 0.6125 \text{ gm}$$

$$\text{Moles of ClO}_3^- = 0.005$$


$$\text{Molar ratio} = 1 : 1$$

Sol 20: (iii) One mL of $\text{Na}_2\text{S}_2\text{O}_3$ is equivalent to

$$0.0499 \text{ gm of CuSO}_4$$


$$0.2 \text{ millimole of CuSO}_4$$

Since reaction is balanced



$$1 \quad \quad \quad 8$$

$$5 \times 10^{-4} \quad \quad 4 \text{ millimole}$$



$$2 \quad \quad \quad 1$$

$$0.2 \text{ millimole} \quad 0.1 \text{ millimole}$$


$$\therefore 0.1 \text{ millimole of I}_2 \text{ equivalent to } 0.2 \text{ millimole of CuSO}_4$$

But we have 4 millimole


$$\text{So, } 1 \text{ mL} \longrightarrow 0.1 \text{ millimole}$$

$$40 \text{ mL} \longleftarrow 4 \text{ millimole}$$

Sol 21: $\text{BaCO}_3 + \text{CaCO}_3 + \text{CaO}$



$$x + y = 7.5 \times 10^{-3}$$



$$0.1 \text{ x}$$

Equating equivalents of $\text{BaCrO}_4 = \text{I}^-$

$$0.1 \text{ x} - 3 = 20 \times 0.05 \times 10^{-3}$$

$$x = \frac{10^{-2}}{3} = 3.33 \times 10^{-3}$$

$$\therefore y = 4.167 \times 10^{-3}$$

$$\text{Mass of BaCO}_3 = 0.659 \text{ gm}$$

$$\text{CaCO}_3 = 0.4167 \text{ gm}$$

$$\therefore \text{Mass of CaO} = 0.1733$$

$$\% \text{ CaO} = 13.87\%$$

Sol 22: $\text{Cu}_2\text{S} + \text{CuS} + \text{MnO}_4^- \longrightarrow$



$$200 \times 0.75 \times 10^{-3} = \frac{175 \times 10^{-3}}{5} + \text{moles of required}$$

$$\therefore \text{Moles required of MnO}_4^- = 0.115$$

$$\text{Let moles of Cu}_2\text{S}^{-2} = x$$

$$\text{CuS} = y$$

$$\therefore 0.115 \times 5 = 8x + 6y$$

Let m = mass of Cu_2S


$$8x + 6y = 0.575$$

$$\frac{8m}{159} + \frac{6(10-m)}{95.5} = 0.575$$

$$[(5.03 \times 10^{-2}) - (6.28 \times 10^{-2})] m = -5.327 \times 10^{-2}$$

$$m = 4.26 \text{ gm}$$

$$\% \text{ CuS} = \frac{5.74}{10} \times 100 = 57.4\%$$



$$\begin{array}{c} \text{moles of air} \\ = 0.406 = \frac{PV}{RT} \end{array} \quad \downarrow \text{Na}_2\text{S}_2\text{O}_3$$

$$\text{Equivalents of } \text{I}_2 = 1.5 \times 10^{-3} \times 0.01 = 1.5 \times 10^{-5}$$


$$\text{Moles of } \text{O}_3 = x$$

$$\therefore x \times 6 = 1.5 \times 10^{-5}$$

$$x = 2.5 \times 10^{-6}$$

$$\therefore \text{Volume of } \text{O}_3 = 1.847 \times 10^{-4} \text{ lit.}$$

$$\% \text{ O}_3 = 1.847 \times 10^{-3} = \frac{V_{\text{O}_3}}{10} \times 100$$



$$30 \times 1.5 \quad 15 \times 0.5 \quad 15 \times 0.4$$



$$(30 \times 1.5) - (15 \times 0.5 + 15 \times 0.4) = 31.5 \text{ milliequivalents}$$


$$\therefore \text{Final of } \text{MnO}_4^- = \frac{31.5 \times 10^{-3}}{60 \times 10^{-3}} = 0.525 \text{ M}$$

Final molarity of

$$\text{Fe}^{3+} = \frac{15 \times 0.4 \times 10^{-3}}{60 \times 10^{-3}} = 0.1 \text{ M}$$

$$\therefore \text{Final normalities } \text{MnO}_4^- = 0.105 \text{ M}$$

$$\text{Fe}^{3+} = 0.1 \text{ M}$$



$$\therefore \text{Equivalents of } \text{H}_2\text{O}_2 = 2 \times 10^{-3}$$

$$\therefore \text{Moles of } \text{H}_2\text{O}_2 = \frac{2 \times 10^{-3}}{2} = 10^{-3}$$

$$\text{Now, molarity} = \frac{10^{-3}}{25} \times 10^3 = 0.04 \text{ M}$$

$$\therefore \text{Normality} = 0.04 \times 2 = 0.08 \text{ N}$$

$$(ii) 20 \times 0.3 \times 10^{-3} = \text{equivalent of } \text{I}^-$$

$$\therefore \text{Normality of } \text{H}_2\text{O}_2 = \frac{20 \times 0.3 \times 10^{-3}}{25 \times 10^{-3}} = 0.24$$

$$\therefore \text{Strength} = 1.344$$





\therefore Equivalent of thiosulphate

$$= \text{Eq. of } \text{I}^- = \text{Eq. of } \text{I}_2 = x \times 45 \times 10^{-3}$$

$$\frac{0.1}{214} \times 5 = x \times 45 \times 10^{-3}$$

$$x = 0.062 \text{ M}$$



Let assume no. of moles of

$$\text{MnSO}_4 \cdot 5\text{H}_2\text{O} = x$$

$$\therefore \text{Moles of } \text{Mn}_3\text{O}_4 = 3x$$

$$3x(6) + 100 \times 0.1 \times 10^{-3} = 0.12 \times 100 \times 10^{-3}$$







$$25 \times N = 30 \times 0.1$$

















$$N = 0.12$$

$$\therefore x = 1.11 \times 10^{-4}$$

$$\therefore \text{Mass of } \text{MnSO}_4 \cdot 4\text{H}_2\text{O} = 1.338 \text{ gm.}$$

Sol 28:





$$5x = y$$



$$3(1-x) = 4-y$$

$$3-3x = 4-y$$

$$x = \frac{1}{2}$$

$$y = \frac{5}{2}$$

$$P_4O_6 = P_4O_{10} = 50\%$$





$$5x = y$$

$$3(3-x) = 11-y$$

$$9-3x = 11-y$$

$$x = 1$$

$$P_4O_{10} = \frac{1}{3}; \quad P_4O_6 = \frac{2}{3}$$



$$5x = y$$

$$3(3-x) = 13-y$$











$$9-3x = 13-y$$



$$x = 2$$

$$P_4O_6 = 2$$

$$P_4O_{10} = 1$$

Sol 30:





Exercise 2

Mole Concept

Single Correct Choice Type




$$\frac{x}{M} = \frac{y}{71+M}$$

$$\frac{x}{M} = \frac{y}{71+M}$$

$$71x + Mx = My$$

$$M = \frac{71x}{x-y}$$



$$\text{Moles of } H_2SO_4 = 0.12$$

$$\text{Mass of } H_2SO_4 = 0.12 \times 98 = 11.76 \text{ gm}$$



$$= \frac{0.5}{100} \times 3 \text{ gm} = 0.015 \text{ gm}$$

$$\text{Number of } I^- = \frac{0.015}{127+23} \times 6.023 \times 10^{23} = 6.023 \times 10^{19}$$

Assertion Reasoning Type



$$PV = nRT$$

At same P and T, $V \propto n$

If M. W. = 44 gm of gas

$$n = 0.01 \text{ mole}$$


$$V \propto n$$

So, volume will be same as moles are also same.



$$M_{\text{avg.}} = x(M_1) + (1-x)M_2$$

So, statement-II is explaining statement-I and both are correct.



$$\text{Mass of nitrogen} = 28$$

$$\% = \frac{28}{60} = 46.66$$


Statement-II: Urea not ionic.



$$2x + 3(-2) = -2$$

$$x = +2$$

Statement: Yes, Because these may be per-oxide bond.




$$= \text{molality} = \frac{\text{moles of solute}}{\text{mass of solvent}}$$

Density increases = moles of solute \uparrow

Molality and molarity both changes.

Statement-II: Density results in change in mass thus increases moles.



Statement-II: Correct.

Multiple Correct Choice Type



by nitrogen balance

$$n_{HNO_3} = \frac{1}{2} n_{NH_3}$$



Let's say 1 mole of NH_3 is initially taken.

$$\text{It makes } \frac{1}{2} - \frac{1}{2} \text{ mole of } HNO_2 \text{ and } HNO_3 \text{ till}$$

reaction-III $\frac{1}{2}$ mole HNO_2 make $\frac{1}{6}$ mole of HNO_3 in
reaction-IV so HNO_3 made

$$= \frac{1}{2} + \frac{1}{6} \text{ mole} = \frac{2}{3} \text{ mole}$$

$$\% \text{ increase} = \frac{\frac{1}{6}}{\frac{1}{2}} = \frac{100}{3} \%$$

(C) By above data, it is correct

$$(D) \text{ Mole of NO produced} = \frac{1}{2} \times \frac{2}{3} = 50 \% \text{ of } \text{HNO}_3$$

Comprehension Type

Paragraph 1

Sol 11: (A) Initially mole of $\text{HCl} = \frac{1}{2}$ mole

$$= \frac{1}{2} \times 36.5 \text{ gm} = 18.25 \text{ gm}$$


So, after heating mole of HCl

$$= \frac{18.25 - 2.75}{36.5} = \frac{15.5}{36.5} = 0.424 \text{ mole}$$

$$\text{Normality} = \frac{0.424}{0.750} = 0.5662 \sim 0.58$$

Sol 12: (C) Please note that, there is a small typo in questions,

Instead of Ca(OH) , it should be Ca(OH)_2



Moles of $\text{HCl} = 0.1 \times 10 = 1 \text{m mole}$

Moles of Ca(OH)_2 required = 0.5 m mole

$$\text{Volume} = \frac{0.5}{0.1} \text{ mL} = 5 \text{ mL}$$

Sol 13: (A) We know valency factor for Na_2CO_3 is 2

$$\text{So, molarity will be} = \frac{0.5}{2} = 0.25 \text{M}$$

Sol 14: (A) 6.90 N means in 1 lit. solution

$\text{KOH} = 6.90 \text{ moles}$


Weight of $\text{KOH} = 6.90 \times (56) = 386.4 \text{ gm}$

given 30% by weight is KOH

So, weight of solution = 12.88 gm

$$\text{Density} = \frac{12.88}{1} = 12.88$$

Sol 15: (C) Ferrous ammonium sulphate



Molecular weight = 390

$$\text{Moles in 0.1 N, 250 mL} = \frac{(0.1)(0.250)}{\text{Valency factor}}$$




Mass of ferrous ammonium sulphate required

$$= (0.1)(0.250)(390) = 9.8 \text{ gm}$$

Paragraph 2

Sol 16: $\text{CuCl}_2 + \text{AgCl} \rightarrow \text{unreacted}$
(4.925-x)gm (5.74-y)gm



Let's say initially $\text{CuBr}_2 = x \text{ gm}$

$$\text{CuCl}_2 = 4.925 - x \text{ gm}$$

$\text{AgCl} = y \text{ gram}$ (reacts with reacted)

$\text{AgCl} = 5.74 - y \text{ gram}$ (in reacted)

Finally same $\text{AgCl} \rightarrow \text{AgBr}$ and CuBr_2

$\rightarrow \text{CuCl}_2$ (completely)

$$\text{Moles of AgCl in reaction} = \frac{y}{143.5}$$

= Mole of AgBr produced

$$\text{Finally AgCl} = (5.74 - y) \text{ gm}$$

$$\text{AgBr} = \frac{y}{143.5} \times (80 + 108) = y(1.310)$$

$$\text{AgCl} + \text{AgBr} = 6.63 = 5.74 + y(0.310)$$

$$y = 2.87 \text{ gm}$$

$$\text{So moles of CuBr}_2 = \frac{2.87}{2 \times 143.5} = \frac{x}{223.5}$$

$$x = 2.235 \text{ gm}$$

$$(1) \text{ (C) CuBr}_2 \text{ mass \%} = \frac{2.235}{4.925} = 45.38\%$$

(2) (B)% mass of $\text{Cu} =$

$$\left[\left(\frac{2.235}{223.5} \right) + \left(\frac{4.925 - 2.235}{63.5 + 71} \right) \right] \times \frac{63.5}{4.925}$$

$$= \frac{0.03 \times 63.5}{4.925} = 38.68\%$$

(3) (B) Mole % of AgBr =

$$\frac{\left(\frac{2.87}{143.5}\right)}{\left(\frac{2.87}{143.5}\right) + \left(\frac{9.74 - 2.87}{143.5}\right)} = 50\%$$

(4) (A) Moles of CuBr₂ = Moles of CuCl₂ produced

= 0.01 mole

Moles of CuCl₂ initially take

$$= \frac{4.925 - 2.235}{134.5} = 0.02 \text{ mole}$$

Mole of Cl⁻ in final solution = (0.01 + 0.02) × 2 = 0.06

Paragraph 3

Sol 17: UF₆ + xH₂O → UO_xF_y + gas (F_{6-y}, H_{2x})

3.52gm 3.08 gm 0.8 gm

0.01 mole

Gas contains 95% fluorine by mass

$$= 0.8 \times \frac{95}{100} = (6 - y) \times 19$$

y = 5.96

$$0.8 \times \frac{5}{100} = (2x)$$

x = 0.02


(1) (C) So empirical formula F_{6-5.96} H_{2(0.02)}

$$= F_{0.04} H_{0.04}$$

= HF

(2) (A) Empirical formula of solid = UO_xF_y

final reaction



0.01 0.01

A = 2B (H-balance)

6 = Y + A (F-balance)

$$B = X = \frac{A}{2} \text{ (O-balance)}$$

Y = 6 - A

$$UO_{\frac{A}{2}}F_{\frac{6-A}{2}} \text{ molecular weight} = \frac{3.08}{0.01} = 308 \text{ gm}$$

$$238 + \frac{A}{2} (16) + (6 - A) \times 19 = 308$$

$$8A + 114 - 19A = 70$$

$$11A = 44$$

$$A = 4$$

So UO₂F₂

$$(3) (A) \% \text{ of F converted} = \frac{A}{6} = 66.66\%$$

Match the Columns

Sol 18: A → r; B → p; C → q

$$\frac{1}{3} Al_2O_3 \text{ molecular weight} = 267 + 135 + 196 = 598$$

$$(A) Y = \frac{267}{598} = 44.95\%$$

$$(B) Al = \frac{135}{598} = 22.57\%$$

$$(C) O = \frac{196}{598} = 32.32\%$$

Sol 19: A → r; B → q; C → p

$$C_6H_8O_6 \text{ molecular weight} = 72 + 8 + 96 = 176$$


$$\text{Moles of } C_6H_8O_6 = \frac{17.6 \text{ mg}}{176} = 0.1 \text{ m mole}$$

$$(A) O - \text{atom} = 6 \times n_{C_6H_8O_6} \times N_A = 3.6 \times 10^{20}$$

$$(B) \text{Mole} = \frac{1}{176} = 5.68 \times 10^{-3}$$

$$(C) \text{Moles of } C_6H_8O_6 = 0.1 \text{ m mole}$$

Sol 20: (C) Volume strength



1 lt. of H₂O₂ gives x lt. of O₂ gas then X is said to be volume strength of H₂O₂

It X - V is given at S. T. P. then

$$\text{Mole of } O_2 \text{ produced} = \frac{x}{22.4}$$

$$\text{Mole of } H_2O_2 \text{ required} = \frac{x}{11.2} \text{ (in litre)}$$

$$\text{Molarity} = \frac{x}{11.2}$$

$$\text{Normality} = \frac{x}{11.2} \times (\text{valency factor}) = \frac{x}{5.6}$$

$$\text{Strength in g/L} = \frac{x}{11.2} \times 34 = \frac{17x}{5.6}$$

$$\text{Volume strength} = \text{Normality} \times 5.6$$

Sol 21: (B) (A) acid + acid \rightarrow No reaction

$$M_{\text{avg.}} = \frac{M_1 V_1 - M_2 V_2}{V_1 + V_2} = \frac{\text{Total no. of moles}}{\text{Total volume}}$$

Similarly (B)

(C) acid + basic \rightarrow

$$M_{\text{avg.}} = \frac{M_1 V_1 - M_2 V_2}{V_1 + V_2}$$

(D) Mili equivalent = $x \times M \times \text{V mL}$

$$(E) \text{ Molarity} = \frac{\text{moles}}{\text{volume}} = \frac{M_1 V_1}{\text{Volume}} = \frac{M_1 V_1}{V_2}$$

$$\text{or } = \frac{\text{moles}}{\text{volume}} = \frac{\text{mass (gm)}}{M_{\text{solute}} \times \text{Volume (lt.)}}$$

Redox Reactions

Single Correct Choice Type

Sol 1: (C) $\text{N}_2\text{H}_4 \rightarrow y = 10e^-$

Each nitrogen loses $5e^-$


\therefore Oxidation no. of N in

$$y = -2 + 5 = +3$$

Sol 2: (D) The ore which gets easily oxidised is best reducing agent

$\text{I}^- \rightarrow \frac{1}{2} \text{I}_2$ is most feasible because.

Sol 3: (C) Alumino thermite process :-



\downarrow

Reducing agent

Sol 4: (D) (a) Oxidation number of S in $\text{H}_2\text{S} = +2$

Oxidation number of S in $\text{SO}_2 = +4$


(b) H_2O_2 can undergo reduction as well as oxidation because oxidation number of oxygen in H_2O_2 is -1. So, it can act both as reducing agent and oxidising agent.

Sol 5: (C) $\text{ClO}_3^- + 6\text{H}^+ + x \rightarrow \text{Cl}^- + 3\text{H}_2\text{O}$

\downarrow


$$6e^- + 5 \rightarrow -1$$

Sol 6: (A) $[\text{Fe}(\text{H}_2\text{O})_5(\text{NO})^+]^{-1} \text{SO}_4^{2-}$



$$x + 1 = +2; x = +1$$

Sol 7: (A) KO_2^-



$$\therefore x = -\frac{1}{2}$$

Sol 8: (B) $3\text{Br}_2^0 + 6\text{CO}_3^{2-} + 3\text{H}_2\text{O} \rightarrow 5\text{Br}^- + 6\text{HCO}_3^- + 5\text{BrO}_3^{+5}$

$\text{BrO} \rightarrow \text{Br}^{-1}$ Reduction

$\text{BrO} \rightarrow \text{Br}^{+5}$ Oxidation

Comprehension Type

Paragraph 1

Sol 9: (D) $\text{H}_2\text{O}_2 + \text{KI} \rightarrow \text{I}_2$
($\times 5$)

\downarrow
hypo

Sol 10: (D) Eq. of hypo solution eq. of I_2

$$20 \times 0.1 \times 10^{-3} = 50 \times 10^{-3} \times N_{\text{H}_2\text{O}_2}$$

$$\therefore N_{\text{H}_2\text{O}_2} = 0.04$$

\therefore Concentration of H_2O_2


$$\text{in gm/lit.} = \frac{0.04}{4} \times 34 = 0.34$$

Sol 11: (D) \therefore Eq. of MnO_2 + Eq. of hypo solution

$$\frac{m}{87} \times 2 = 30 \times 0.1 \times 10^{-3}$$

$$m = 0.1305$$


$$\therefore \% = \frac{0.1305}{0.5} \times 100 = 26.1\%$$



\therefore Valence factor = 5 for As




2 for I

Paragraph 2



Disproportionation

(oxidation as well as reduction)



Multiple Correct Choice Type


Sol 16: (A, B, D) Meq. of formed = Meq. of HCl used for NH_3

$$= 50 \times 0.15 - 32.10 \times 0.10$$

$$= 4.29$$

These Meq. of NH_3 are derived using valance factor of $\text{NH}_3 = 1$ (an acid base reaction)

In redox change valance factor of NH_3 is 8;



\therefore Meq. of NH_3 for valance factor


Also, Meq. of $\text{NO}_3^- = \text{Meq. of } \text{NH}_3$

$$= 8 \times 4.29 = 34.32$$

$$\therefore N_{\text{NO}_3^-} = \frac{34.32}{25} = 1.37$$


($N \times V$ in mL = Meq.)

Assertion Reasoning Type



Valency of Cr is 6 all O have higher electronegativity than Cr

\therefore Cr's oxidation no. = +6



	BaCl_2	+	$\text{Al}_2(\text{SO}_4)_3$	\longrightarrow	BaSO_4	+	AlCl_3
Meq. before	30×0.1		40×0.2		0		0
reaction	= 3		= 8		= 0		= 0
Meq. after reaction	0		5		3		3

Sol 18: (D) Avg. oxidation no. of Pb_3O_4 is $+\frac{8}{3}$. But in reality, Pb_3O_4 is made up of $\text{PbO} + \text{PbO}_2$. So, actively, Pb have oxidation state +2, +4.

Sol 19: (C) Oxidation no. of Cl = +7 it can not be greater than this

\therefore It can get only reduced

$\therefore \text{HClO}_4$ is an oxidising agent

\therefore In HClO_3 , oxidation no. of chlorine = + 5

E.N. order $\text{O} > \text{Cl} > \text{H}$




Sol 20: (D) Since S_2^{2-} has S – S⁻ linkage structure

$\therefore \text{FeS}_2^{2-} \rightarrow \text{Fe}^{2+}$ (S – S) oxidation no. = +2

Sol 21: (B) Yes, the given reaction is an example of disproportionation

$\therefore \text{H}_2\text{O}_2$ is a reducing as well as an oxidising agent

So it is not only bleaching (oxidising agent)




These reactions show $E^\circ > 0$

∴ It is not feasible because iodine can show multiple oxidation state.

Match the Columns

Sol 24: A → w; B → x; C → u; D → p; E → v; F → q; G → r; H → s; I → t

- (1) Increase in oxidation no:- Loss of electrons (oxidation)
- (2) Decrease in oxidation no:- Gain of only e⁻s (reduction)
- (3) Oxidation agent:- Gain of e⁻s
- (4) Reducing agent:- Loss of e⁻s
- (5) $2\text{Cu}^+ \longrightarrow \text{Cu}^{2+} + \text{Cu}$



Disproportionation reaction

(6) Redox reaction

(7) Mn_3O_4 oxidation no:-

$$+\frac{8}{3} \text{ fractional}$$

(8) CH_2Cl_2


$x + 2 - 2 = 0$ zero oxidation no.

$x = 0$

(9) $\text{NaOH} + \text{HCl} \rightarrow \text{NaCl} + \text{H}_2\text{O}$




Simple neutralisation reaction

Sol 25: A → p, s; B → r; C → p, q; D → p





Disproportionation

Redox reaction



Redox Reaction



Redox reaction

Previous Years' Questions

Mole Concept

Sol 1: Average atomic weight

$$= \frac{\text{Percentage of an isotope} \times \text{atomic weight}}{100}$$

$$\Rightarrow 10.81 = \frac{10.01x + 11.01(100 - x)}{100}$$

$$\Rightarrow x = 20\%$$

Therefore, natural boron contain 20% (10.01) isotope and 80% other isotope.

Sol 2: From the vapour density information,

Molar mass = Vapour density × 2

(∵ Molar mass of $\text{H}_2 = 2$)

$$= 38.3 \times 2 = 76.6$$

Now, let us consider 1.0 mole of mixture and it contains x mole of N_2 .

$$\Rightarrow 46x + 92(1 - x) = 76.6$$


$$\Rightarrow x = 0.3348$$

$$\text{Also, in 100 g mixture, number of moles} = \frac{100}{76.6}$$


$$\Rightarrow \text{Moles of } \text{N}_2 \text{ in mixture}$$

$$= \frac{100}{76.6} \times 0.3348 = 0.437$$

Sol 3: Heating below 600°C converts $\text{Pb}(\text{NO}_3)_2$ into PbO but to NaNO_3 into NaNO_2 as:



MW. 330 222



MW.85 69

$$\text{Weight loss} = 5 \times \frac{28}{100} = 1.4 \text{ g}$$

$$\Rightarrow \text{Weight of residue left} = 5 - 1.4 = 3.6 \text{ g}$$

Now, let the original mixture contain x g of $\text{Pb}(\text{NO}_3)_2$

\therefore 330 g gives 222 g PbO

\therefore x g $\text{Pb}(\text{NO}_3)_2$ will give $\frac{222x}{330}$ g PbO

Similarly, 85 g NaNO_3 gives 69 g

\Rightarrow (5 - x)g will give


$\frac{69(5-x)}{85}$ g NaNO_2

\Rightarrow Residue: $\frac{222x}{330} + \frac{69(5-x)}{85} = 3.6\text{g}$

x = 3.3 g $\text{Pb}(\text{NO}_3)_2$

\Rightarrow $\text{NaNO}_3 = 1.7$ g

Sol 4: Compound B forms hydrated crystals with $\text{Al}_2(\text{SO}_4)_3$. Also, B is formed with univalent metal on heating with sulphur. Hence, compound B must have the molecular formula M_2SO_4 and compound A must be an oxide of M which reacts with sulphur to give metal sulphate as



\therefore 0.321 g sulphur gives 1.743 g of M_2SO_4


\therefore 32.1 g S (one mole) will give 174.3 g M_2SO_4

Therefore, molar mass of $\text{M}_2\text{SO}_4 = 174.3$ g


$\Rightarrow 174.3 = 2 \times \text{Atomic weight of M} + 32.1 + 64$

\Rightarrow Atomic weight of M = 39, metal is potassium (K)

K_2SO_4 on treatment with aqueous $\text{Al}_2(\text{SO}_4)_3$ gives potash-alum.



If the metal oxide A has molecular formula MO_x , two moles of it combine with one mole of sulphur to give one mole of metal sulphate as



$\Rightarrow x = 2$, i.e., A is KO_2 .

Sol 5: 93% H_2SO_4 solution weight by volume indicates that there is 93 g H_2SO_4 in 100 mL of solution.

If we consider 100 mL solution, weight of solution = 184 g

Weight of H_2O in 100 mL solution

= 184 - 93 = 91 g

\Rightarrow Molality = $\frac{\text{Moles of solute}}{\text{Weight of solvent (g)}} \times 1000$

$$= \frac{93}{98} \times \frac{1000}{91} = 10.43$$

Sol 6: Partial pressure of $\text{N}_2 = 0.001$ atm,

T = 298 K, V = 2.46 dm³.

From Ideal Gas law : $pV = nRT$

$$n(\text{N}_2) = \frac{pV}{RT} = \frac{0.001 \times 2.46}{0.082 \times 298} = 10^{-7}$$

\Rightarrow No. of molecules of

$$= 6.023 \times 10^{23} \times 10^{-7}$$

$$= 6.023 \times 10^{17}$$


Surface sites used in adsorption

$$= \frac{20}{100} \times 6.023 \times 10^{17} = 2 \times 6.023 \times 10^{16}$$

\Rightarrow Sites occupied per molecules

$$= \frac{\text{Number of sites}}{\text{Number of molecules}} = \frac{2 \times 6.023 \times 10^{16}}{6.023 \times 10^{16}} = 2$$

Sol 7: (D) The balanced chemical reaction is



In this reaction, 3 moles of BaCl_2 combined with 2 moles of Na_3PO_4 . Hence, 0.5 mole of BaCl_2 requires

$$\frac{2}{3} \times 0.5 = 0.33 \text{ mole of } \text{Na}_3\text{PO}_4.$$

Since available Na_3PO_4 (0.2 mole) is less than required mole (0.33), it is the limiting reactant and would determine the amount of product $\text{Ba}_3(\text{PO}_4)_2$.

\therefore 2 moles of Na_3PO_4 gives 1 mole $\text{Ba}_3(\text{PO}_4)_2$

\therefore 0.2 mole of Na_3PO_4 would give

$$\frac{1}{2} \times 0.2 = 0.1 \text{ mole } \text{Ba}_3(\text{PO}_4)_2$$

Sol 8: (B) The following reaction occur between

$\text{S}_2\text{O}_3^{2-}$ and $\text{Cr}_2\text{O}_7^{2-}$:




Change in oxidation number of $\text{Cr}_2\text{O}_7^{2-}$ per formula unit is 6 (it is always fixed for $\text{Cr}_2\text{O}_7^{2-}$).

Hence, equivalent weight of $\text{K}_2\text{Cr}_2\text{O}_7$

$$= \frac{\text{Molecular weight}}{6}$$


Sol 9: $\text{Na}_2\text{S}_4\text{O}_6$ is a salt of $\text{H}_2\text{S}_4\text{O}_6$ which has the following structure



⇒ Difference in oxidation number of two types of sulphur = 5.

Sol 10: (B) 1. Both assertion and reason are factually true but the reason does not exactly explain the assertion. The correct explanation is, methyl orange and phenolphthalein changes their colours at different pH.

Sol 11: For the oxidation of A^{n+} as:




Gram equivalent of $\text{A}^{n+} = 2.68 \times 10^{-3} (5 - n)$

Now equating the above gram equivalent with gram equivalent of KMnO_4 :

$$2.68 \times 10^{-3} (5 - n) = 1.61 \times 10^{-3} \times 5$$

⇒ $n = +2$

Sol 12: The redox reaction involved is :



If M is molarity of H_2O_2 solution, then

$$5M = \frac{0.508 \times 1000}{254}$$

(∵ 1 mole $\text{H}_2\text{O}_2 \equiv 1$ mole I_2)

⇒ $M = 0.4$

Also, n-factor of H_2O_2 is 2, therefore normality of H_2O_2 solution is 0.8 N.

⇒ Volume strength = Normality $\times 5.6$

= $0.8 \times 5.6 = 4.48 \text{ V}$

Sol 13: With KMnO_4 oxalate ion is oxidized only as:



Let, in the given mass of compound, x millimol of $\text{C}_2\text{O}_4^{2-}$ ion is present, then

Meq. of $\text{C}_2\text{O}_4^{2-} = \text{Meq of MnO}_4^-$

⇒ $2x = 0.02 \times 5 \times 22.6$; ⇒ $x = 1.13$

At the later stage, with I^- , Cu^{2+} is reduced as :



and $\text{I}_2 + 2\text{S}_2\text{O}_3^{2-} \longrightarrow 2\text{I}^- + \text{S}_4\text{O}_6^{2-}$

Let there be x millimole of Cu^{2+}

⇒ Meq of $\text{Cu}^{2+} = \text{Meq of I}_2 = \text{meq of hypo}$

⇒ $x = 11.3 + 0.05 = 0.565$

⇒ Meq of $\text{Cu}^{2+} : \text{Meq of C}_2\text{O}_4^{2-} = 0.565 : 1.13 = 1 : 2$

Sol 14: Let us consider 10 mL of the stock solution contain x millimol oxalic acid $\text{H}_2\text{C}_2\text{O}_4$ and y millimol of NaHC_2O_4 . When titrated against NaOH , basicity of oxalic acid is 2 while that of NaHC_2O_4 is 1.

⇒ $2x + y = 3 \times 0.1 = 0.3$... (i)

When titrated against acidic KMnO_4 , n-factor of both oxalic acid and NaHC_2O_4 would be 2.

$2x + 2y = 4 \times 0.1 = 0.4$... (ii)

Solving equations (i) and (ii) gives $y = 0.1$, $x = 0.1$.

⇒ In 1.0 L solution, mole of $\text{H}_2\text{C}_2\text{O}_4 = \frac{0.1}{1000} \times 100 = 0.01$

Mole of $\text{NaHC}_2\text{O}_4 = \frac{0.1}{1000} \times 100 = 0.01$

⇒ Mass of $\text{H}_2\text{C}_2\text{O}_4 = 90 \times 0.01 = 0.9 \text{ g}$

Mass of $\text{NaHC}_2\text{O}_4 = 112 \times 0.01 = 1.12 \text{ g}$


Sol 15: (D) (p) $\text{PbO}_2 + \text{H}_2\text{SO}_4 \xrightarrow{\Delta} \text{PbSO}_4 + \text{O}_2 + \frac{1}{2} \text{O}_2$

(q) $2\text{Na}_2\text{S}_2\text{O}_3 + \text{Cl}_2 + 2\text{H}_2\text{O} \longrightarrow 2\text{NaCl} + 2\text{NaHSO}_4 + 2\text{S}$


(r) $\text{N}_2\text{H}_4 + 2\text{I}_2 \longrightarrow \text{N}_2 + 4\text{HI}$

(s) $\text{XeF}_2 + 2\text{NO} \longrightarrow \text{Xe} + 2\text{NOF}$

Sol 16: (A, B, D) The balanced equation is



Sol 17: (A) $\text{KIO}_4 + \text{H}_2\text{O}_2 \longrightarrow \text{KIO}_3 + \text{H}_2\text{O} + \text{O}_2$



Empower your JEE Main & Advanced preparation with specially designed and structured study packages from PLANCESS

PlanceSS offers this must-have Study Package in Chemistry comprising seven books to meet the complete curriculum needs of engineering aspirants. The books offer complete and systematic coverage of JEE (Main) and JEE (Advanced) syllabi and aims to provide firm foundation in learning and problem solving skills.

The theory offers engaging explanation of concepts with important points highlighted through Facts, Notes, Tips & Tricks, Misconceptions and Key Take Away. The application of the concepts is explained by means of illustrative examples and the problem-solving skills are further enhanced through solved examples of JEE (Main) and JEE (Advanced) levels. Extensive and classified assessment (JEE Main, JEE Advanced and Boards) along with the questions from previous years papers of the target examinations, offers the aspirants a competitive edge in preparation. This study package thus provides a single solution for all their preparation requirements.

Salient Features

- Concept description sprinkled with PlanceSS Concepts and Illustrative Examples
- Problem-Solving Strategy (Tactics)
- Formula Sheet/Points to Remember
- Solved Examples - JEE Main
- Solved Examples - JEE Advanced
- Exercises - JEE Main and Boards
- Previous Years' Questions - JEE Main
- Exercises - JEE Advanced and Boards
- Previous Years Questions - JEE Advanced
- Answer Key and Solutions (Complete)

PlanceSS EduSolutions Pvt. Ltd.

A005/A006 Boomerang
Chandivali Farm Road,
Chandivali, Andheri (East)
Mumbai - 400072, India
Tel: 022 6604 3405
www.planceSS.com



Copyright © 2017