



PLANT KINGDOM

Contents

Topic	Page No.
Exercise	02 - 7
Notes	
Classification	08 - 09
Algae	10 - 11
Algae Classification	12 - 14
Bryophyte	15 - 17
Pteridophytes	18 - 19
Gymnosperms	20 - 22
Angiosperms	23 - 26

Syllabus

PLANT KINGDOM

Classification, Algae, Algae Classification, Bryophyte, Pteridophytes, Gymnosperms, Angiosperms

Name : _____ Contact No. _____

ETOOSINDIA.COM

India's No.1 Online Coaching for JEE Main & Advanced
3rd Floor, H.No.50 Rajeev Gandhi Nagar, Kota, Rajasthan, 324005
HelpDesk : 92-14-233303

PLANT KINGDOM

LEVEL - I

1. Colonial algae is -
(A) Spirogyra (B) Iamanaria (C) Fucus (D) None
2. Algae is shows -
(A) Isogamous (B) Ausogamous (C) Heterogamous (D) All
3. Largest size algae is -
(A) Kelp (B) Spirogyra (C) Chlamydomonas (D) Volvox
4. Anisogamy shows by -
(A) Chlamydomonas (B) Volvox (C) Fucus (D) Spirogyra
5. Heterogamy shows by -
(A) Fucus (B) Volvox (C) both 1 & 2 (D) Spirogyra
6. Filamentous algae is -
(A) Ulothrix (B) Volvox (C) Chlamydomonas (D) None
7. Red Algae is -
(A) Volvox (B) Polysiphonia (C) Chara (D) Ectocarpus
8. Flagella absent in gamete of -
(A) Chlorophyceae (B) Rhodophyceae (C) Phacophyceae (D) None
9. Phaeophyceae algae is -
(A) Chara (B) Sargassum (C) Ulothrix (D) Spirogyra
10. Rhodophyceae algae is not -
(A) Gracillaria (B) Porphira (C) Gellidium (D) Fucus
11. Storage food is starch is -
(A) Chlorophyceae (B) Phaeophyceae (C) Rhodophyceae (D) None
12. Kelp belong to -
(A) Phaeophyceae (B) Rhodophyceae (C) Chlorophyceae (D) None
13. Moss is -
(A) Marchantia (B) Sphagnum (C) Riccia (D) None
14. Fern belong with -
(A) Aglae (B) Bryophytes (C) Ptendophyte (D) Gymnosperm
15. Pterodophyte is NOT -
(A) Ginkago (B) Selaginella (C) Fern (D) Equisetum
16. Horse tail plant is -
(A) Equistitum (B) Fern (C) Selaginella (D) None

17. Protonemas formed is -
 (A) Bryophyte (B) Algae (C) Gymnosperm (D) None
18. Moss & liver wort belong with -
 (A) Algae (B) Lichen (C) Bryophyte (D) None
19. Michorrhiza shows -
 (A) Gymnosperm (B) Angiosperm (C) Bryophyte (D) None
20. Psilopsida members is -
 (A) lycopodium (B) Pteris (C) Selaginella (D) none

Answer Key (Level - I)

- | | | | | | | | | | | | |
|-----|---|-----|---|-----|---|-----|---|-----|---|-----|---|
| 1. | D | 2. | D | 3. | A | 4. | A | 5. | C | 6. | A |
| 7. | B | 8. | B | 9. | B | 10. | D | 11. | A | 12. | A |
| 13. | B | 14. | C | 15. | A | 16. | A | 17. | A | 18. | C |
| 19. | A | 20. | A | | | | | | | | |

LEVEL - II

1. An algae which can be employed as food for human being is -
(A) Ulothrix (B) Chlorella (C) Spirogyra (D) None
2. Lack of a all or all show -
(A) Cyanobacteria (B) Sea fan (C) Sacchromyces (D) BGA
3. The life cycle of algae such as spirogyra is -
(A) Haplontic (B) Diplontic (C) Haplo-diplontic (D) diplo
4. Chlorophyll of red algae is -
(A) Chl - a (B) Chl - a, Chl - b (C) Chl - a, Chl - d (D) Chl - a, Chl - e
5. Manmitol as a strong food in -
(A) Rhodophyceae (B) Phaeophyceae (C) Chlerophyceae (D) Poaceae
6. Ectocarpus shows life cycle -
(A) haptontic (B) diplontic (C) haplo-diplontic (D) diplo-haplontic
7. Presence of pyrenoid shows by -
(A) Phaeophyceae (B) Chlorophycar (C) Rhodophyceae (D) Poaceae
8. Alginic acid is found in the cell wall of -
(A) Gigartira (B) Laminaria (C) Gelidium (D) Sytonema
9. Algae which form motile colony is -
(A) Volvox (B) Nostoc (C) Spirogyra (D) None
10. Iodine is found in algae -
(A) Ulva (B) Ulothrix (C) dilorella (D) Lamanaria
11. Reserve food is floridean starch in -
(A) Rhodophyceae (B) Phaeophyceae (C) Chlorophyceae (D) Xantaophyceae
12. Non motile, greatly tuicked, asexual spoxe in chlomydomonas is -
(A) Carpospore (B) Aplanospore (C) Alcinete (D) Hypnospor
13. Parosite algae is -
(A) Volvox (B) Ulothrix (C) Porphyra (D) Cephaluros
14. Identify the alga, which exhibits diplontic life cycle -
(A) Spirogyra (B) Chlomydomonos (C) Fucus (D) Volvox
15. Botanical name of peat moss is -
(A) Sphagnum (B) Funaria (C) Antroceros (D) Polytrichum
16. Present in Archegoniophore -
(A) Chara (B) Adiantum (C) Funaria (D) Marchantia

17. Moss spore germinate to form -
 (A) Sporophyte (B) Protonema (C) Seta (D) capsule
18. Largest moss is -
 (A) Pogonatum (B) Funaria (C) Dawsonia (D) Polytrichum
19. Dominant generation in bryophytes is -
 (A) capsule (B) sporophyte (C) Gametophyte (D) Seta
20. The plants having vascular tissue but lacking seeds are present under -
 (A) Algae (B) bryophytes (C) pteridophytes (D) Gymnosperm

Answer Key (Level - II)

- | | | | | | |
|-------|-------|-------|-------|-------|-------|
| 1. C | 2. B | 3. A | 4. C | 5. B | 6. C |
| 7. B | 8. B | 9. A | 10. D | 11. A | 12. D |
| 13. D | 14. C | 15. A | 16. D | 17. B | 18. C |
| 19. A | 20. C | | | | |

LEVEL - III

1. Protuallus of fern produces ?
(A) spores (B) gametes (C) Both (D) 1 & 2
2. The site of photo synthesis in BGA is -
(A) Chromato phores (B) mitochondria (C) Chleroplast (D) root hair
3. Phycology is the study of -
(A) Algae (B) Fern (C) Fungi (D) Bryophytes
4. Fern spores are usually ?
(A) haploid (B) diploid (C) triploid (D) tetra ploid
5. Agar-Agar is obtained from -
(A) Chlorella (B) Spirogyra (C) Ulthrix (D) Gelidium
6. Multiflagellate gamete in :
(A) Equisetum (B) Riccia (C) Lycopodium (D) Anthoceros
7. Club moss belongs to :
(A) Algae (B) Pteridophytes (C) Fungi (D) Bryophytes
8. Diatoms belong to which class -
(A) Phawphyceae (B) Bacillariophyceae (C) Chlorophyceae (D) Xantnophyceae
9. Heterocyst are found in -
(A) Eyanophyceae (B) Chlorophyceae (C) Phaeophyceae (D) Rhodophyceae
10. Manmitol as a strong food in -
(A) Rhodhophyceae (B) Phaeophyceae (C) Chlerophyceae (D) Poaceae
11. Green algae contain -
(A) chl-a & chl-b (B) starch (C) carotenoid (D) All of these
12. Which is the following responsible for peat farmation -
(A) Marchentia (B) Riccia (C) Funaria (D) Splagnum
13. Endo sperm of gymnosperm is -
(A) diploid (B) Tetraploid (C) baploid (D) none
14. Number of archlgania in cycas is -
(A) 8 (B) 4 (C) 3 (D) 2
15. In cycas -
(A) Archegania are present (B) Antneridia are present
(C) Archegania are absent (D) Both a & b
16. Rhodophyceae algae is not -
(A) Gracillaria (B) Porphira (C) Gellidium (D) Fucus

17. Ovule of gymnosperm is -
 (A) bitegmic (B) Unitegmic (C) naked (D) both 2 & 3
18. In gymnosperm, the seeds are naked because they lack -
 (A) Integument (B) uncellus (C) pericarp (D) perianth
19. Resin and Turpentine are products of -
 (A) Teak (B) Oak (C) Eucalyptus (D) Pine
20. Monoecious plant is -
 (A) Marchantia (B) Pinus (C) cycas (D) Papaya

Answer Key (Level - III)

- | | | | | | | | | | | | |
|-----|---|-----|---|-----|---|-----|---|-----|---|-----|---|
| 1. | B | 2. | C | 3. | A | 4. | A | 5. | D | 6. | A |
| 7. | B | 8. | B | 9. | A | 10. | B | 11. | A | 12. | D |
| 13. | C | 14. | B | 15. | A | 16. | D | 17. | D | 18. | C |
| 19. | D | 20. | B | | | | | | | | |

CLASSIFICATION

Artificial System of Classification

This system is based on comparison of one or a few superficial characteristics, which are helpful in easy identification of organisms. This system remained in use for about two thousand years.

Advantage

Advantages of artificial system are given below

- (i) Artificial system is easy to remember as only one or few characters are used.
- (ii) The traits used are of interest to humans.

Disadvantages

Disadvantages of artificial system are given below

- (i) This system uses only few superficial characters (i.e., habits, numbers, colours and shapes of leaves, etc) which leads to many organisms grouped together.
- (ii) They considered mainly the vegetative characters or the androecium characters as given by Linnaeus.
- (iii) It does not demonstrate natural and phylogenetic relationships.
- (iv) This gave equal weightage to vegetative and reproductive/sexual characters this is not acceptable, as vegetative characters are more easily influenced by the environmental factors.
- (v) They separated the closely related species.

Natural System of Classification

It is also known as **phenetic system** of classification. The natural system of classification is based on natural affinities among the organisms. It considers both external and internal features like structure, anatomy, embryology and phytochemistry.

Advantage

Advantages of natural system are given below :

- (i) Only related organisms are kept in a group.
- (ii) Unrelated organisms are kept in separate groups.
- (iii) It shows natural relationships among the organisms.
- (iv) It shows possible origin of different taxa.

Disadvantage

Disadvantages of natural system are given below :

- (i) There is more emphasis given on natural character.
- (ii) In this system several related families are separate and unrelated families are put together.
- (iii) Evolutionary basis is neglected.

Phylogenetic System of Classification

The phylogenetic system of classification indicates the evolutionary as well as genetic relationship among organisms. This system is based on fossil records of biochemical, anatomical, morphological, physiological embryological and genetical.

The system was initiated by **Engler** and **Prantl** (1887-1899) in *Die Naturalischen Pflanzfamilien*. In phylogenetic system, flowering plants are placed in ascending series related to complexity of floral morphology. The phylogenetic system of classification are mainly the rearrangement of taxonomic characters in addition to the phylogenetic information.

Advantages

Advantages of phylogenetic system are given below :

- (i) Families and order in this system are of small size.
- (ii) This system is in conformation with the modern views of phylogeny.
- (iii) They use information from various sources to solve problems of classification. Such information become more important in the absence of supporting fossil evidences.

Disadvantage

Disadvantages of phylogenetic system are given below :

- (i) This is not helpful in plant identification.
- (ii) The classification is outdated as the habit is used as main basis of classification.

Types of Taxonomy

For the suitability in studies various categorisations has been done in taxonomy.

The important taxonomies are as follows

1. Numerical Taxonomy

It is carried out by quantitative assessment of similarities and differences in order to make objective assessment. It is now easily carried out using computers based on all observable characteristics. Number and codes are assigned to all the characters and the data are then processed. In this way, each character is given equal importance and at the same time hundreds of characters can be considered.

2. Cytotaxonomy

Cytotaxonomy is based on cytological information like chromosome number, structure, behaviour and type of chromosomes. For example,

- (i) Chromosome number is constant for a species, e.g., 46 in man, 48 in apes and potato, 20 in maize, 16 in onion and 8 in *Drosophila*.
- (ii) Behaviour of chromosomes was used by taxonomists during pairing and banding patterns to understand the relationships between species e.g., Origin of humans from apes, origin of wheat etc.

3. Chemotaxonomy

It is the system based on the evidences from chemical constituents (enzymes, hormones, proteins, amino acids, etc.), some specific chemicals (usually secondary metabolites) and the chemical nature of proteins have been utilised to establish similarities and relationships. For example, the presence of raphides has been found to be common in 35 families of plants.

Algae

Algae are chlorophyll-bearing, simple, talloid, autotrophic and largely aquatic (both freshwater and marine) organisms.

The important salient features of algae are given below :

(i) **Habitat**

These are found in both freshwater and marine habitats. Some algae forms are also found in moist habitats like wet rocks and soil, tree trunks. etc. Some of them are also found in close association with fungi (lichen) and animals (e.g., on sloth bear).

(ii) **Thallus Organisation**

The plant body (thallus) is without differentiation. The basis form and size of algae is highly variable, and ranges from filamentous (e.g., *Ulothrix* and *Spirogyra*) to colonial (e.g., *Volvox*). These are attached on the substratum with the help of holdfast.

(iii) **Structure of Algae Cell**

It has on eukaryotic plant cell structure. The cytoplasm contains membrane bound chloroplast, mitochondria, ER, Golgi bodies and other cell organelles. The cell wall contains cellulose. The nucleus contains nuclear membrane.

(iv) **Food Material**

Algae have main food reserve as starch. In brown algae, mannitol and laminarine are the main reserve food material, whereas in red algae floridean starch is the reserve food material.

(v) **Reproduction**

The algae reproduces vegetatively, asexually and sexually.

(i) **Vegetative Reproduction**

It occurs by fragmentation, tubes, stolons, adventitious branches, etc. Each fragment gets develop into a thallus.

(ii) **Asexual Reproduction**

It occurs by a number of accessory spores, such as zoospores aplanospores, akinetes, carpospores, etc. The most common being the zoospores, which are flagellated. The cells which produce spores are called sporangia. The sporangia may be the vegetative cells (e.g. in *Chlamydomonas* and *Ulothrix*) or modified vegetative cells (e.g. in *Vaucheria*).

(iii) **Sexual Reproduction**

It occurs by fusion of two gametes.

On the basis of morphology of reproductive cells, sexual reproduction is of two types :

- (a) **Isogamy** In this method, two morphologically similar gametes fuse to form a zygote, e.g., *Spirogyra*.
- (b) **Heterogamy** In this process, fusion occurs between morphologically as well as physiologically different gametes. It is of two types
 - ♦ **Anisogamy** It is the fusion of structurally dissimilar gametes, which differ in size and behaviour. Male gamete is more active and female gamete is less active and bigger in size, e.g., Some species of *Chlamydomonas*.
 - ♦ **Oogamy** In this process, the male gamete is motile, active, small and without reserve food. The female gamete is bigger, passive, non-motile and laden with food, e.g., *Volvox*, *Fucus*.

(vi) Embryo and Life Cycle

An embryo stage is non present. Life cycle is haplontic, diplontic, diplohaplontic haplohaplontic, etc. An alternation of generation occurs in diplohaplontic life cycle.

(vii) Economic Importance of Algae

- (i) Algae are responsible for carrying out about a half of the total carbon dioxide fixation on earth by the process of photosynthesis.
- (ii) some forms of marine brown and red algae produce large amount of hydrocolloids. These are **alginate** (brown algae) and **carrageenan** (red algae) which have many commercial uses.
- (iii) The algae *Gelidium* and *Gracilaria* are used to produce **agar**, which is used in preparation of ice creams and jellies.
- (iv) Some protein rich algae, like *Chlorella* and *Spirulina* are used as food supplements by sailors and space travellers.
- (v) About 70 species of marine algae are used as food, e.g., *Porphyra*, *Laminaria* and *Sargassum*.

Edujournal.in

Classification of Algae

Algae are divided into various classes based on pigmentation, stored food and flagellation. The main classes are Chlorophyceae, Phaeophyceae and Rhodophyceae.

i. **Class Chlorophyceae (Green Algae)**

The members of Chlorophyceae are commonly called **green algae**. There are about 7000 species in this class. The reserve food material is starch.

The characteristics feature of Chlorophyceae are discussed below :

- (a) **Habitat** These are mostly marine forms, only some are freshwater. *Chlorella* can tolerate moderately warm waters. Snow dwelling forms are called **cryophytes**, e.g., *Chlamydomonas nivalis*, *Scotiella*, etc.
- (b) **Cell Organisation** These are unicellular, colonial, coenocytic and multicellular forms. Cell walls contain cellulose (inner layer) and pectose (outer layer) in most of the green algae. The chloroplasts may be discoid, plate-like, reticulate, cup-shaped, spiral or ribbon shaped.
The chloroplasts contain pigments. Most of the members have one or more stage bodies called **pyrenoids** located in the chloroplasts. Pyrenoids contain protein besides starch. Some algae may store food in the form of oil droplets also.
- (c) **Thallus** Unicellular green algae can be flagellate, (*Chlamydomonas*), unicellular, non-flagellate (*Chlorella*). *Acetabularia* (umbrella plant) has unic upto 10 cm long with distinction nucleus containing rhizoid, elongated stem and umbrella like cap.
A colony for fixed number of individual unicells (*Volvox*) is called **coenobium**. Coenocytic or siphonaceous thallus occur in *Caulerpa*. Unbranched filamentary *Cladophora*, heterotrichous in *Stigecloniu* and parenchymatous in *Ulva*.
- (d) **Reproduction** The members Chlorophyceae reproduce vegetatively asexually and sexually by various methods.
 - ♦ Vegetative reproduction occurs through cell division (unicellular forms fragmentation, stolons, tubers, storage cells, etc).
 - ♦ Asexual reproduction occurs by zoospores, aplanospores, hyphospores, akinetes and daughter colonies.
 - ♦ Sexual reproduction may be isogamous, anisogamous or oogamous.
- (e) **Life Cycle** It can be haplontic, diplontic and diplohaplontic. In haplontic life cycle there is a single somatic phase, which is haploid. **Diploid** stage is represented by a single cell zygote, e.g., in *Spirogyra*.

ii. **Cell-Phaeophyceae (Brown Algae)**

The members of Phaeophyceae are fucoxanthin and phycocolloid rich multicellular eukaryotic algae. Its common members are seaweeds called **kelps**. This class is comprised of about 2000 species.

- (a) **Habitat** Brown algae are mostly marine. These are found mostly in colder seas or during cold seasons in tropical regions.
- (b) **Size** They range from simple branched, filamentous forms (*Ectocarpus*) to profusely branched forms as represented by kelps.
These represent largest algae. The largest kelps are *Macrocystis* (40-100m) and *Nereocystis* (20-30m).
- (c) **Cell Organisation** All members are multicellular. Cell wall is composed of cellulose, pectose and phycocolloids. The cellulosic wall of vegetative cells is usually covered on the outside by a gelatinous coating of **algin**.

- (d) **Thallus** It is heterotrichous filament with both prostrate and upright branches (*Ectocarpus*). The parenchymatous structure is found in higher forms.
The plant body of large forms often differentiated into **holdfast** (with which it usually attaches to the substratum) a stalk called the **stipe** and **lamina** (frond), which is photosynthetic. Conducting tubes or trumpet hyphae are present in larger brown algae or kelps. They help in conduction of food materials.
- (e) **Photosynthetic Pigments and Colour** These include chlorophyll-a, c and carotenoids. They range in colour from olive green to various shades of brown depending upon the amount of the xanthophyll pigment, fucoxanthin present in them.
- (f) **Food Reserve** It remains in the form of complex carbohydrates such as laminarin or mannitol.
- (g) **Flagellation** These contain heterokont flagellation with one smooth (whiplash) and one tinsel flagella.
- (h) **Reproduction** Vegetative reproduction occurs through fragmentation (e.g., *Sargassum*), adventitious branches and stolons (e.g., *Dictyota*). Asexual reproduction by biflagellate zoospores, which are pear-shaped having two unequal laterally attached flagella.
Sexual reproduction is performed by isogamy, anisogamy and oogamy. Union of gametes may take place in water or within the oogonium (oogamous species). The gametes are pyriform (pear-shaped) and have two laterally attached flagella.
- (i) **Life Cycle** Isomorphic alternation of generation is found in some brown algae, e.g., *Ectocarpus*, *Dictyota*. In many brown algae, the diploid generation or phase is dominant. The haploid phase is either microscopic or represented by gametes only (e.g., *Fucus*).

Economic Importance of Pheophyceae

- ♦ The brown algae that are used as food are *Laminaria*, *Nerocystis*, *Macrocystis*, *Alaria*, etc.
- ♦ Some brown algae like *Fucus*, *Sargassum*, *Laminaria*, *Macrocystis* are important fodder for cattle.
- ♦ Brown algae are collected from sea shores and used as manure. They improve mineral content of soils.
- ♦ Alginic acid is a phycocolloid obtained from a number of brown algae such as *Alaria*, *Macrocystis*, *Ascophyllum*, *Laminaria*. It is used as emulsifier, thickener, gelling agent in toothpastes, shaving creams, ice-creams, emulsion paints, shampoo, cosmetics, etc.
- ♦ *Fucus* and *Laminaria* are rich in iodine.

iii. Class-Rhodophyceae (Red Algae)

The members of Rhodophyceae are commonly called red algae because of the predominance of the red pigment, i.e., **r-phycoerythrin** in their body.

- (a) **Habitat** Most of the red algae are marine with greater concentrations found in the warmer areas. They are found in both well lighted regions close to the surface of water and also at great depths in oceans where relatively little light penetrates.
- (b) **Thallus** The red thalli of most of the algae are multicellular. Some of them have complex body organisation like *Asterocystis* is pseudofilamentous, *Porphyridium* is unicellular, *Porphyra* has parenchymatous sheets, *Chondrus* is ribbon like, *Gelidium* is a multicellular sea weed.
- (c) **Cell Wall** The cell wall contains cellulose, pectic compounds and certain mucopolysaccharides called phycocolloids, such as agar, carrageenin, etc. In many algae, cell wall contains pits.
- (d) **Photosynthetic Pigments** These include chlorophyll-a, carotenes, xanthophylls and phycobilins. Phycobilins are water soluble and are of two types, i.e., red-coloured **phycoerythrin** and blue-coloured **phycocyanin**.

- (e) **Reserve Food** It is floridean starch similar in constitution to glycogen and amylopectin.
- (f) **Reproduction** Vegetative reproduction occurs by fragmentation, regeneration of hold fast and gemmae. Asexual reproduction occurs by non-motile spores (carpospores, monospores, tetraspores and neutral spores).
Sexual reproduction occurs by non-motile gametes and is oogamous type. The male sex organs is called **spermatogonium** or **antheridium**. The male produced is non-flagellated, called as spermatium.
The female sex organ is called carpogonium. After fertilisation, a new structure called carposporophyte is produced. It remains attached to the parent alga.
- (g) **Life Cycle** Life Cycle has two or more phases such as haplohaplontic, haplohaplohaplontic, diplodiplohaplontic, etc.

Economic Importance of Red Algae

- ♦ The Red algae like *Porphyra*, *Chondrus*, *Rhodymenia*, *Centerella* and *Bostrychia* are used as food in various parts of the world.
- ♦ Agar yielding algae are called aerophytes, such as *Gelidium*, *Gracilaria*, *Gelidiella*, etc.
- ♦ Carrageenin a phycocolloid obtained from red algae like *Chondrus* and *Gigartina* is used in preparations of emulsions for ice cream, chocolates, sauces, toothpastes, cosmetics, etc. It is also used in clearing liqueurs and finishing leather, etc.
- ♦ Funori and adhesive phycocolloid is obtained from red alga *Gloiopeltis*. It is used in sizing textiles, paper and as glue.
- ♦ Some algae like *Corallina*, *Polysiphonia* have medicine properties.

Salient Feature of Major Classes of Algae			
Character	Chlorophyceae	Phaeophyceae	Rhodophyceae
Habitat	Freshwater, marine or terrestrial.	Only four genera are fresh water, rest marine.	Mostly marine; a few are terrestrial.
Thallus form	Unicellular, colonial, coenobial, filamentous, thalloid or siphonaceous.	Filamentous, branched usually heterotrichous.	Unicellular, filamentous pseudoparenchymatous
Cell Structure	Eukaryotic, well organized nucleus and other cell organelles present.	Eukaryotic	Eukaryotic
Cell wall	Cell wall is made up of cellulose and pectic material.	Outer mucilaginous layer of cell wall consists of alginic and fucinic acids and the inner layer is made up of cellulose.	Outer layer is pectic in nature, whereas inner layer is cellulosic.
Flagella	Motile cells with 2, 4 or more flagella; flagella anteriorly inserted equal in length and whiplash type.	2, laterally inserted, of unequal length; one of them whiplash and another of tinsel type.	Non-flagellate, motile cells are absent.
Dominant pigments	Chlorophyll-a, and b, α and β -carotene, xanthophyll.	Chlorophyll-a, and c, β -carotene, xanthophyll (violaxanthin, fucoxanthin)	Chlorophylla, and d β -carotene (lutein) xanthophylls,
Nucleus	Well organized nucleus with chromosomes and nucleolus is present.	As in Chlorophyceae.	As in Chlorophyceae.

Bryophytes

Bryophytes

Bryophytes include the various mosses and liverworts. These are non-vascular embryophytes, characterised by the presence of an independent gametophyte and parasitic sporophyte.

Habitat

Bryophytes commonly grow in moist, shaded areas in hills. These are also called amphibians of the plant kingdom because, these can live in soil but are dependent on water for sexual reproduction.

Rhizoids

These are attached to the substratum by unicellular multicellular rhizoids.

Sex Organs

The main plant body produces gametes, hence it is cal gametophyte. The sex organs in bryophytes multicellular, the male sex organs in bryophytes is cal antheridium (which produces biflagellate antherozoi and the flask-shaped female sex organ is cal archegonim (produces a single egg.).

Reproduction

It is of sexual type reproduction. The antherozoids : released into water they come in contact with archegonium.

Fertilisation

It occurs inside the archegonium. The egg secretes chemical which attracts spermatozoids. Sperms required thin film of water for swimming and reaching to dehiscid archegonium. One sperm fuses with an egg and produces a **diploid zygote**. Zygotes do not under reduction division immediately instead, they produce multicellular body called **sporophyte**.

The sporophyte is n free-living but attached to the photosynthet gametophyte deriving nourishment from it. Some cells the sporophyte undergo reduction division (meiosis) produce **haploid spores** (which germinate to produced gametophyte).

The sporophyte of bryophytes is called sporogonium because it is mainly dependent and mean for producing spores.

Life Cycle

Bryophytes have heteromorphic or heterologous alternation of generation. The gametophyte many be produced directly or first from a juvenile stage called protonema.

Economic Importance

Bryophytes in general are of little economic importance But, several species have some uses.

- (i) Some mosses provide food for herbaceous mammals birds and other animals.
- (ii) Species of Sphagnum (a moss), provides peat the have long been used as fuel. It has the capacity to hold water as packing material for transshipment of living material.
- (iii) Mosses along with lichens are the first organisms to colonise rocks. Hence, these help in biological succession. They decompose rocks making the substrate suitable for the growth of higher plants.
- (iv) Mosses from dense mats on the soil, they reduce the impact of falling rain and prevent soil erosion.
- (v) Marchantia has medicinal properties to cure lungs and liver infections. It also has antitumour properties.

Bryophytes do not attain heights. They may range of 0.4 to 70 cm and have thalloid body.

The reasons may be following

- (i) Root is absent.
- (ii) Vascular tissue are not present.
- (iii) Cuticle is not present on the plant body.
- (iv) Absence of mechanical tissue.
- (v) Male gametes need to swim upto the interior of dehiscent archegonia.

Types of Bryophytes

The bryophytes are generally classified into two main groups

i. Liverworts

Liverworts (Hepaticopsida) usually grow in moist, shady places such as banks of streams, marshy ground, damp soil, bark of trees and deep in the woods.

- (a) **Thallus** The plant body of a liverwort is thalloid, e.g., *Marchantia*. The thallus is dorsoventral and closely appressed to the substrate. The leafy members have tiny leaf like appendages in two rows on the stem like structure.
- (b) **Rhizoids** These are unicellular. These may be of two types in some liverworts, i.e., smooth walled and tuberculate.
- (c) **Reproduction** It may occur by sexual and asexual means.
 - ♦ **Asexual Reproduction** It occurs by fragmentation of thalli or by the formation of specialised structures called **gemmae** (sign. gemma). Gemmae are green, multicellular, asexual buds which develop in small receptacles called gemma cups located on the thalli. The gemmae become detached from the parent body and germinate, forming new individuals.
 - ♦ **Sexual Reproduction** During this male and female sex organs are produced either on the same or on different thalli. The sporophyte is differentiated into a foot, seta and capsule. After meiosis, spores are produced within the capsule.
 - ♦ These spores germinate to form free-living gametophytes, e.g., *Riccia*, *Marchantia*, *Pellia*, etc.

ii. Mosses

Mosses (Bryopsida) grow in dense mats over moist shady places, especially during rains. Some mosses grow in desert bogs and streams.

- (a) **Plant Body** The predominant stage of the life cycle of a moss is the gametophyte which consists of two stages the first stage is the **protonema stage** which develops directly from a spore. It is a creeping, green, branched and frequently filamentous stage; the second stage is the **leafy stage** which develops from the secondary protonema as a lateral bud. They consist of upright, slender axes bearing spirally arranged leaves. This stage bears sex cells.
- (b) **Rhizoids** These are long, multicellular branched structures with oblique septa. They take part in fixation and absorption of water. However, surface conduction through capillarity is an important mechanism of water supply to aerial parts.
- (c) **Reproduction** This can occur both by vegetative and sexual means.
 - ♦ **Vegetative Reproduction** It occurs by fragmentation and budding in the secondary protonema from exposed rhizoids and other parts (like gemmae, buds and tubes).
 - ♦ **Sexual Reproduction** Sex organs, **antheridia** and **archegonia** are produced the apex of the leafy surface. Male organs appear cup-shaped, while female organs are bud-like. After fertilisation, the zygote develops into a sporophyte, consisting of foot, seta and capsule (containing spores).

(d) **Sporophyte** The sporophyte in mosses is more elaborate than that in liverworts. Spores are formed after meiosis. The mosses have an elaborate mechanism of spore dispersal, e.g., *Funaria*, *Polytrichum* and *Sphagnum*.

Difference between Liverworts and Mosses

Liverworts

They are dorsiventral.
Plants are thalloid or foliose.
Midribs is not present in leaves.
Rhizoids are unicellular and unbranched.
Plants bear scales.
A conducting strand is absent.
Sporophyte is differentiated in foot, seta and capsule.
Capsule often possesses elaters.
Dehiscence occurs through elaters.
A protonema stage is absent.

Mosses

They have radial symmetry.
Mosses are foliose.
Leaves have unbranched midrib.
Rhizoids are multicellular and branched.
Scales are absent in plants.
A conducting strand is commonly present.
It is not differentiated in foot, seta and capsule
Elaters are absent.
Peristome perform these functions.
A juvenile protonema stage is present.

Pteridophytes

Pteridophytes are primitive seedless vascular plants called **cryptogams**. These have conspicuous sporophyte plant body, inconspicuous independent gametophyte with antheridia and partially embedded archegonia have 4-rowed necks.

The term Pteridophyte was coined by **Haeckel** (1866). There are about 13000 species of pteridophytes have been reported. They were perhaps the first land plants evolved during ordovician (450-500) million years ago) period.

The characteristics features of pteridophytes are as follow :

Habitat

The pteridophytes are found in cool, damp shady places though some may flourish well in sandy-soil conditions. Some members like *Azolla*, *Salvinia*, *Marsilea* species.

Plant Body

The size varies from a fraction of centimeter (e.g., *Azolla*) to 20 m in tree ferns (e.g., *Angiopteris*). The main plant body a sporophyte which is differentiated into true roots, and leaves. These organs possess well differentiated vascular tissues. The leaves are small (microphylls) as in *Selaginella* or large (macrophylls) as in **ferns**.

Vascular Tissues

These are xylem and phloem present throughout the body. Xylem consists of tracheids and phloem is made up of sieve cells and albuminous cells.

Sporophylls

The sporophytes bear sporangia that are subtended by leaf-like appendages called sporophylls. In some cases sporophylls may form distinct compact structures called **strobili** or **cones** (*Selaginella* and *Equisetum*). The sporangia produce spores by meiosis in spore mother cells.

Spores

The spores germinate to give rise to inconspicuous, small but multicellular free-living, mostly photosynthetic thalloid gametophytes called **prothallus**. In majority of the pteridophytes, all the spores are of similar kinds; such plants are called **homosporous**. Genera like *Selaginella* and *Salvinia* which produce two kinds of spores, macro (large) and micro (small) spores; such plants are called **heterosporous**.

Gametophyte

The thalloid gametophyte or prothallus requires cool, damp, shady places to grow. The megaspores and microspores germinate and give rise to female and male gametophytes respectively. The female gametophytes in these plants is retained on the parent sporophytes for viable periods. In most ferns, **prothallus** is green and autotrophic. In heterosporous ferns, the female gametophyte depends on food by the megaspore.

Sex Organs

The gametophytes bear male sex organs called **antheridia** and female sex organs called **archegonia**. Antheridium is sessile and surrounded by a single layered jacket. Archegonium is flask-shaped. It is partially embedded.

Fertilisation

Water is required for transfer of antherozoids. The male gametes released from the antheridia reach the mouth of archegonium. Fusion of male gamete with the egg present in the archegonium results in the formation of zygote. Zygote therefore, produces a multicellular, well differentiated **sporophyte**, which is the dominant phase of the pteridophytes.

Embryo

Fertilisation produces a zygote that undergoes division to produce embryo. The development of the zygote into young embryo takes place within the female gametophyte. This event is a precursor of the **seed habit** and considered as an important step in evolution, e.g., *Dryopteris*, *Selaginella*, *Adiantum*, *Equisetum* and *Salvinia*.

Economic Importance of Pteridophytes

- (i) Pteridophytes are a good source of food for animals. For example, sporocarps of *Marsilea* is edible. *Angiopteris* and *Alsophila* have starchy pith eaten by natives of Australia.
- (ii) Ferns protect soil from erosion by providing a good cover on the hill slopes and other fragile places.
- (iii) *Equisetum* stems have rough surfaces. They are used in scrubbing and polishing.
- (iv) *Azolla* a water fern has a symbiotic association with nitrogen fixing cyanobacterium *Anabaena azollae*. It is cultured in paddy fields to harbour nitrogen fixing bacterium. Thus, act as a **biofertiliser**.
- (v) Ferns are also grown as ornamental plants for their graceful plant body.

Classification of Pteridophytes

The pteridophytes are further classified into four classes

- (i) Psilopsida (*Psilotum*)
- (ii) Lycopsidea (*Selaginella* and *Lycopodium*)
- (iii) Sphenopsida (*Equisetum*)
- (iv) Pteropsida (*Dryopteris*, *Pteris* and *Adiantum*)

Difference between Bryophytes and Pteridophytes

Bryophytes

The main plant body is gametophyte.
These are non-vascular plants.
Sporophyte is parasitic over gametophyte.
Plant body can be thallus or foliose.
True stems and leaves are not present.
Roots are absent, rhizoids are present.
Sex organs are stalked.
The wall of archegonial neck is 5-6 rowed.

Pteridophytes

It is sporophyte.
These are vascular plants.
Sporophyte is independent of gametophyte.
It is differentiated into stem, leaves and roots
It has true stems and leaves
Roots are present
Sex organs are sessile.
The wall of archegonial neck is 4-rowed.

Gymnosperms

The gymnosperms (*Gymnos* = naked ; *sperma* = seeds) are plants in which the ovules are not enclosed by any ovary wall and remain exposed, both before and after fertilisation. These are small groups of seed plants which are represented by only 900 living species. Unlike bryophytes and pteridophytes, in gymnosperms the male and the female gametophytes do not have an independent free-living existence. They remain within the sporangia retained on the sporophytes.

Habitat

These plants are mostly found in colder parts of northern hemisphere, where they form extensive forests. A number of gymnosperms are now, known as ornamentals, e.g., *Ginkgo*, *Thuja*, *Araucaria*, etc.

Morphology

Gymnosperms include, medium-sized trees or tall trees and shrubs. The giant red wood tree *Sequoia* is one of the forest tree up to 100 m. Species of *Gnetum* are woody climbers. The smallest gymnosperm is *Zamia pygmaea* which reaches a height of 25 cm. Many of the gymnosperms live for more than 400 years, e.g., Pine (*Pinus*), redwood (*Sequoia*).

External Features

The plant body is sporophyte and differentiated into root, stem and leaves.

Plant Body

Tap roots are present for proper anchorage to heavy plant. Roots in some genera have fungal association in the form of **mycorrhiza** (*Pinus*), while in some other (*Cycas*) small specialised roots called **coralloid roots** are associated with N₂-fixing cyanobacteria such as *Anabaena*, *Nostoc*, etc.

The stems are branched (*Pinus*, *Cedrus*) or unbranched (*Cycas*). The leaves may be simple or compound.

In *Cycas*, the pinnate leaves persist for few years. The leaves in gymnosperms are well-adapted to withstand extremes of temperature, humidity and wind. While, in conifers, the needle-like leaves reduce the surface area. Their thick cuticle and sunken stomata also help to reduce water loss.

Internal Features

Plants possess vascular tissues i.e., *xylem* and *phloem*. They are arranged in vascular bundles.

Vascular Tissues

The conducting part of xylem consists of tracheid. Vessels occur in *Gnetum* and its relatives. Phloem has sieve cells, but are arranged in rows in *Gnetum*.

Reproduction

The gymnosperms are heterosporous, i.e., they produce haploid microspores and megaspore.

Cones

The two kinds of spores are produced within sporangia that are borne on sporophylls, which are arranged along an axis to form compact strobili or cones. They are produced in different types of sporangia called microsporangia or microsporophylls and ovules or megasporangia respectively.

The strobili bearing microsporangia are called **microsporangiate** or **male strobilia** and the strobili, bearing megasporophylls with ovules or megasporangia are called **macrosporangiate** or **female strobili**.

Pollen Grains

The microspores develop into a male gametophytic generation which is highly reduced and is confined only to a limited number of cells, known as pollen grain. The development of pollen grains occurs within the microsporangia.

Ovules

The megaspore mother cell is differentiated from one of the cells of the nucellus, which is protected by envelopes and the composite structure called ovule. The ovules are borne on the megasporophylls which may be clustered together forming female cones.

Archegonia

The megaspore mother cell thus, undergo meiotic division forming four megaspores. Out of which one is enclosed within the megasporangium (nucellus) and develops into a multicellular female gametophyte, bearing two or more archegonia of female sex organs.

The multicellular female gametophyte is also retained within megasporangium.

Fertilisation

Air current required for transport of male gametes. The male gametes are carried to the archegonia. i.e., gamete in the ovule by means of a tube called **pollen tube** and discharge their contents near the mouth of archegonia. This is called siphonogamy.

Seeds

Following fertilisation, zygote develops into an embryo and the ovules into seeds. The seeds contain food laden tissue called **endosperm**. It lies naked or exposed. Endosperm provides nourishment for growth of seedling at the time of seed germination.

Examples *Cycas*, *Pinus*, *Ginkgo*, *Ephedra*, *Gnetum*, *Vaucheria*, *Cedrus*, *Abies*, etc.

Difference between Microsporophyll and Megasporophyll

Microsporophyll

It bears microsporangia.
It contains numerous microspores.
Microspores are released.

Megasporophyll

It bears megasporangia.
It usually contains one megaspore.
Megaspore are retains inside the megasporangium.

Classification of Gymnosperms

Gymnosperms further include three main classes

Cycadopsida (*cycads*), Coniferopsida (*conifers*) Gnetopsida (*Gnetum*).

Difference between Male Gametophyte of Pteridophyte and Gymnosperms

Male Gametophyte of Pteridophyte

A distinct male gametophyte may not be present.
It contains an antheridium.
Male gametes are flagellate.
Male gametes reach female gametes by swimming in water.

Male Gametophyte of Gymnosperms

A distinct male gametophyte is present.
Antheridium is not present.
Male gametes can be flagellate or non-flagellate.
Male gametes reach the female gamete through a pollen tube.

Economic Importance of Gymnosperms

- (i) Seeds of *Pinus gerardiana* (chilgoza) are used a food after roasting. Some other edible gymnosperms plant part are endosperm of *Ginkgo*, seed kernel of *Cycas* and *Gnetum*, sego grains from stems of *Cycas* etc.
- (ii) Gymnosperms provide softwood for construction, plywood and paper industry.
- (iii) Branches of gymnosperm trees ae used as food.
- (iv) Saw dust of conifers is used in making plastic and linoleum.

- (v) Resin is a semifluid secreted by special tubes of number of conifers. It helps in sealing female cones after pollination, scale leaves around leaf bases and apical buds. Resin is distilled to obtain **turpentine** and **resin**. Turpentine is used for thinning paints, varnishes, etc., and resin is used for sealing joints, wheel green, preparation of writing paper, oil clothes, etc.
- (vi) **Ephedrine** an antibiotic is obtained from Ephedra. It is used for curing respiratory problems and asthma. **Taxol** is an anticancer drug obtained from taxus.

Edujournal.in

Angiosperms

Angiosperms are seed bearing plants or flowering plants. Unlike gymnosperms where the ovules are naked, in angiosperms the sporophyll are organised into flowers and the seeds are produced inside fruits. There are about 250000 species of angiosperms in the nature. They evolved about 130-160 million years ago.

Habitat

Angiosperm are found in wide range of habitats, from the land to upto 6000 m in Himalayas or Antarctica and Tundra, dry hot deserts, cold deserts, tropics, fresh water up to 60⁰ C, underground, over other plants as parasites, saprophyte, etc. *Zostera* is a marine angiosperm.

Plnat Body

These plant are sporophytic, in the form of herbs, shrubs, trees, climber creepers, etc. The smallest angiosperm is water plant *Wolffia* and tallest is *Eucalyptus regnans* (100 m and above). Primary root develops from radicle. It forms tap root system. In many angiosperms roots develop from places other than radicle, these are **adventitious roots**. Stem develops from **plumule**.

Leaves

These are simple or compound. The leaves bear axillary buds which can grow into stem branches.

Vascular Tissues

Angiosperms have vessels in xylem. Phloem contains sieve tubes and companion cells in regard to gymnosperms which do not have companion cells.

Flowers

Flowers are the reproductive structures formed by the union of one or both types of sporophylls (microsporophylls or stamens and megasporophylls or carpels).

Microsporophylls or Male Sex Organs

Stamens are considered as the male sex organs of a flower. Each stamen has two main parts. i.e., a slender **filament** with an **anther** (at the tip). An anther contains four microsporangia where microspore mother cells become differentiated to form four microspores, each developing into a pollen grain.

Megasporophylls or Female Sex Organs

Carpel or pistil is called the female sex organ of the flower. Each carpel has three parts (i.e., an ovary, style and stigma). A megaspore mother cell is differentiated in the nucellus and undergoes meiosis, ultimately on functional megaspore gets enlarged and forms the female gametophyte known as embryo-sac.

Each cell of an embryo sac is haploid.

Embryo sac

Each embryosac contain a three called egg apparatus consisting of one egg cell and two synergids, three antipodal cells (at the opposite end) and two polar nuclei (in the central cell). The polar nuclei eventually fuses and forms diploid secondary nucleus.

Pollination

Pollen grains after dispersal from the anthers are carried by various ways such as wind, water or by various other agencies to the stigma of the pistil.

Fertilisation and Development of a Seed

Each pollen grain germinates on the stigma forming a pollen tube that carries two male gametes to the embryo sac, growing through the tissues of stigma and style.

One of the male gametes fuses with the egg cell to form a zygote (syngamy).

The other male gamete fuses with the diploid secondary nucleus to produce the triploid Primary Endosperm Nucleus (PEN). Because of the involvement of above mentioned two events, it is known as **double fertilisation**. After fertilisation, synergids and antipodal cells degenerate.

Endosperm

The zygote develops into embryo and the primary endosperm nucleus develops into an endosperm.

Fruit

The ovule gradually transforms into a seed and the ovary becomes the fruit. A fruit is actually a ripened ovary. They not only protect the seed but also help in their dispersal.

Difference between Gymnosperm and Angiosperm

Gymnosperms	Angiosperms
Vessel or trachea is absent in xylem.	Trachea is present in xylem.
Phloem contains sieve cells.	Phloem tissue contains sieve tube and companion cells.
Sporophyll are aggregated to form cones.	Sporophyll are aggregated to produce flowers.
Sepals and petals absent.	Sepals and petals present.
Cones are generally unisexual.	Flowers are generally bisexual.
Microsporophyll are broad and leaf like.	Microsporophyll have a stalk and terminal anthers.
Megasporophylls are woody and there is not rolling of megasporophylls.	The megasporophyll are softer and rolled into ovary, style and stigma.
Ovules are exposed.	Ovules are enclosed in ovary.
Archegonia is present.	Archegonia absent.
Pollen enters microphyte directly.	Pollen germinates on stigma, pollen tube passes through style to enter ovary.
Single fertilisation.	Double fertilisation.
Endosperm is haploid and formed before fertilisation.	Endosperm is triploid and formed after fertilisation.
Seeds exposed because there is no ovary, so no fruit formation.	Seeds are enclosed in a fruit which forms after fertilisation.

Classification of Angiosperms

George Bentham and Joseph Dalton Hooker presented the system of classification of angiosperms published in *Genera Plantarum* (1862-1883) which appeared in three volumes.

This system of classification is used by most of the well known Herbaria of the world. The details of this system are not described here. Conveniently, on the basis of the number of cotyledons angiosperm are classified in two broad groups i.e., Monocotyledonae and Dicotyledonae.

Difference between Monocots and Dicots

Monocots	Dicots
They contain one cotyledon.	They contain two cotyledons.
Leaves have parallel venation.	Leaves have reticulate venation.
Fibrous root system is present.	Tap root system is present.
Stomata are dumb-bell shaped.	Stomata are kidney-shaped.
Vascular bundles are scattered.	Vascular bundles are arranged in rings.
Cambium is absent.	Cambium is present.
Stems do not have concentric arrangement of tissue. A ground tissue is present.	Stems have concentric arrangement of tissue epidermis cortex, endodermis, pericycle, pith, etc.
Secondary growth is absent with some exception in stems.	These show secondary growth in stem.
Root has pith in its centre.	Root is generally devoid of pith.
Secondary growth is absent in roots with some exception.	Secondary growth occurs in roots.

Economic Importance of Angiosperm

- (i) The angiosperms are major source of food, fibres, and beverages.
- (ii) They also provide valuable timber and medicines.
- (iii) These also add beauty to our environment as well.

Alternation Generation

Life cycle of an organism is a sequence of events that occur from birth to death of an organism. In plants, both haploid and diploid cells can divide by mitosis.

This feature leads to the formation of different plant bodies, haploid and diploid. The haploid plant body produces gametes by mitosis. This plant body represents gametophyte.

After fertilisation, the zygote also divides by mitosis to produce a diploid sporophyte plant body. Haploid spores are produced by this plant body by meiosis. The spores, in turn, divide by mitosis to form a haploid plant body once again.

Thus, during the life cycle of a sexual reproducing plant, there is an alternation of generation between gamete-producing haploid gametophyte and spore-producing diploid sporophyte.

Plant Life Cycles


Different plant groups and individual different features in their life cycle

Haplontic

The dominant photosynthetic phase in gametophyte produced by haploid sporophyte. The gametophyte produces gametes by mitosis.

The gametes fuse and produce a diploid zygote, that represents sporophyte generation. There are no free-living sporophytes. Meiosis in the zygote results in the formation of haploid spores. This kind of life cycle is called **haplontic**.


Many algae such as Volvox, Spirogyra and Chlamydomonas represent this pattern of life cycle.




Diplontic

In this type, the diploid sporophyte is the dominant. The multicellular diploid phase is called **sporophyte**. The gametophytic phase is represented by the single-celled haploid gametophyte.

This kind of life cycle is termed as diplontic. All seed-bearing plants, gymnosperms and angiosperms follow this pattern of life cycle. *Fucus*, an alga, is diplontic.





Haplodiplontic

In this type, there are two distinct multicellular phases, diploid sporophyte and haploid gametophyte are present. Both phases are multicellular. However, they differ in their dominant phases.

- (i) A dominant, independent, photosynthetic, thalloid or erect phase is represented by a haploid gametophyte. It alternates with the short lived multicellular sporophyte totally, partially or dependent on the gametophyte for its anchorage and nutrition. All bryophytes represent this pattern.
- (ii) The diploid sporophyte is represented by a dominant independent photosynthetic vascular plant body. It alternates with multicellular saprophytic/autotrophic, independent but short lived haploid gametophyte. This pattern is called haplodiplontic life cycle.

All pteridophytes demonstrate this pattern.

However, most algal genera are haplontic, some of them such as Ectocarpus, Polysiphonia and kelps are haplodiplontic.